

Umberto Marconi

Some conditions under which a uniform space is fine

Commentationes Mathematicae Universitatis Carolinae, Vol. 34 (1993), No. 3, 543--547

Persistent URL: <http://dml.cz/dmlcz/118611>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Some conditions under which a uniform space is fine

UMBERTO MARCONI

Abstract. Let X be a uniform space of uniform weight μ . It is shown that if every open covering, of power at most μ , is uniform, then X is fine. Furthermore, an ω_μ -metric space is fine, provided that every finite open covering is uniform.

Keywords: uniform space, uniform weight, fine uniformity, uniformly locally finite, ω_μ -additive space, ω_μ -metric space

Classification: Primary 54E15; Secondary 54A25, 54A35

0. A recurrent problem about uniform spaces is to see whether a uniformity is the finest one compatible with the topology. Isiwata and Atsuji solved this problem in metric spaces [2], [6]. The following theorem summarizes some equivalent conditions of Theorem 1 of [2].

Theorem 1. *The following conditions on a metric space X are equivalent:*

- (1) *every open covering is uniform;*
- (2) *every countable open covering is uniform;*
- (3) *every open covering consisting of two elements is uniform;*
- (4) *the subset K of limit points is compact and, for every uniform covering \mathcal{U} , the subspace $X \setminus \text{St}(K, \mathcal{U})$ is uniformly discrete.*

(The star $\text{St}(K, \mathcal{U})$ of K with respect to \mathcal{U} is the union of all elements of \mathcal{U} which have a non-empty intersection with K).

It is interesting to see if a suitable version of Theorem 1 also holds in uniform spaces. Metric spaces are uniform spaces with a countable base for the uniformity; this base can be assumed to be well ordered by star-refinement. In uniform spaces, the existence of a well ordered base is a very strong property.

We will prove that an equivalence analogous to $1 \Leftrightarrow 2$ holds for general uniform spaces and depends only on cardinal properties, while the equivalence $1 \Leftrightarrow 3$ can be generalized to ω_μ -metric spaces (= uniform spaces which admit a base of uniform coverings well ordered by a regular cardinal ω_μ). In ω_μ -metric spaces we will provide a suitable formulation of the condition 4 (obviously in uniform spaces $3 \neq 1$ and $1 \neq 4$).

1. Unless otherwise specified, the space X is a uniform space, with the uniform topology.

The following lemma is useful for working with cardinal properties of locally finite families. The proof is easy to check (for example, see [7]).

Lemma 1. *Let \mathcal{G} be a locally finite family of subsets of X . If the power of \mathcal{G} is at most μ , then there exists an open covering \mathcal{B} , of power at most μ , such that every element of \mathcal{B} meets only finitely many elements of \mathcal{G} .*

We recall that the uniform weight of X is the smallest cardinal number of a base for the uniformity.

The following theorem is the uniform analogue of $2 \Rightarrow 1$ in Theorem 1.

Theorem 2. *Let μ be the uniform weight of X . If every open covering of power at most μ is uniform, then every open covering is uniform.*

We shall prove Theorem 2 in three steps (the statements of each step hold under the hypotheses of Theorem 2).

Let $\{\mathcal{U}_\alpha : \alpha < \mu\}$ be a base for the uniformity, consisting of open uniform coverings.

Step 1. *X is a paracompact topological space.*

PROOF: Let \mathcal{A} be an open covering. For every $x \in X$, choose $\alpha(x) < \mu$ such that $\text{St}(x, \mathcal{U}_{\alpha(x)})$ is contained in some element of \mathcal{A} . For every α , let

$$A_\alpha = \bigcup_{\alpha(x)=\alpha} \text{St}(x, \mathcal{U}_{\alpha(x)}).$$

By assumption, the covering $\{A_\alpha : \alpha < \mu\}$ is uniform.

The covering

$$\mathcal{B} = \{A_\alpha \cap \text{St}(x, \mathcal{U}_{\alpha(x)}) : \alpha < \mu, \alpha(x) = \alpha\}$$

is an open refinement of \mathcal{A} .

Since uniform coverings are normal coverings (in the sense of Tukey), by the condition (g) in [9, Theorem 1.2] it follows that the covering \mathcal{B} has an open star-refinement.

□

A family \mathcal{F} of subsets of X is said to be uniformly locally finite if there exists a uniform covering \mathcal{B} such that every element of \mathcal{B} meets \mathcal{F} in only a finite number of elements.

Step 2. *Every locally finite family \mathcal{F} is uniformly locally finite.*

PROOF: We proceed by contradiction. Suppose that for every α there exist an element $U_\alpha \in \mathcal{U}_\alpha$ and a countable subfamily \mathcal{F}_α of \mathcal{F} such that $U_\alpha \cap F \neq \emptyset$ for every $F \in \mathcal{F}_\alpha$.

Let \mathcal{G} be the union of all subfamilies \mathcal{F}_α . \mathcal{G} is a locally finite family of subsets and the power of \mathcal{G} is at most μ . By Lemma 1, there exists an open covering \mathcal{B} of power at most μ such that every element of \mathcal{B} meets \mathcal{G} only in a finite number of elements. \mathcal{B} is an open covering of power at most μ , which cannot be uniform because the family \mathcal{G} is not uniformly locally finite. This is a contradiction with the hypothesis of Theorem 2. □

Step 3. Every locally finite open covering \mathcal{A} is uniform.

PROOF: By Step 2, \mathcal{A} is uniformly locally finite. For every α , by possibly refining coverings \mathcal{U}_α , we can assume that every element of \mathcal{U}_α meets \mathcal{A} only for a finite number of elements.

If \mathcal{A} is not a uniform covering, then for every α there exists $U_\alpha \in \mathcal{U}_\alpha$ such that $U_\alpha \setminus A \neq \emptyset$ for every $A \in \mathcal{A}$.

Let $\mathcal{A}_\alpha = \{A \in \mathcal{A} : A \cap U_\alpha \neq \emptyset\}$ and let $\mathcal{C} = \bigcup_\alpha \mathcal{A}_\alpha$.

Every \mathcal{A}_α is a finite family, thus the power of \mathcal{C} is at most μ . Therefore we have a contradiction, because the open covering $\mathcal{C} \cup \{\bigcup(\mathcal{A} \setminus \mathcal{C})\}$, of power at most μ , cannot be uniform. In fact, for every α , $U_\alpha \cap (\bigcup(\mathcal{A} \setminus \mathcal{C})) = \emptyset$ and $U_\alpha \setminus A \neq \emptyset$ for every $A \in \mathcal{C}$. \square

The conclusion of Theorem 2 follows from Step 1 and Step 3.

Remark. One might conjecture that every open covering is uniform, provided that the open coverings of power less than μ are uniform coverings. This, however, is not the case. For a counterexample, let X be the space of ordinals less than ω_1 , equipped with the unique (precompact) uniformity (an open covering is uniform iff it has a finite subcovering).

By countable compactness of X , every countable open covering is uniform and it is easy to verify that the uniform weight of X is ω_1 . Furthermore, X is not a paracompact topological space [4, p. 380].

2. Denote by \mathbf{C}^* the weak uniformity of continuous bounded real functions on a completely regular Hausdorff space X .

X is a normal space iff every open covering consisting of two elements belongs to \mathbf{C}^* .

It is an interesting question to see when a uniformity finer than \mathbf{C}^* is fine. For example, the implication $3 \Rightarrow 1$ of Theorem 1 says that metric uniformities finer than \mathbf{C}^* are fine. Another example of uniform space with this property are sequentially uniform spaces [3].

In the next theorem, we extend the equivalences $1 \Leftrightarrow 3 \Leftrightarrow 4$ of Theorem 1 to ω_μ -metric spaces. Notice that the proof of this theorem follows from ordinal properties.

An ω_μ -metric space is a uniform space which admits a base of uniform coverings

$$\mathbf{B} = \{\mathcal{U}_\alpha : \alpha < \omega_\mu\}$$

well ordered by refinement (hence by star-refinement) by a regular cardinal ω_μ .

An ω_μ -metric space is paracompact (ultra-paracompact if $\mu > 0$) [1].

Let λ be a cardinal number. A topological space is said to be λ -compact if every open covering has a subcovering of power less than λ . A weakly paracompact space X is λ -compact iff the power of every discrete closed subset of X is less than λ (as one can prove by mimicking the proof of [4, Theorem 5.3.2]).

In the proof of Theorem 3, the base \mathbf{B} is assumed well ordered by star-refinement. The equivalence $1 \Leftrightarrow 2$ has been already proved in [8].

Theorem 3. Let X be an ω_μ -metric space. The following conditions are equivalent:

- (1) every open covering is uniform;
- (2) the set K of limit points is ω_μ -compact and for every α the subspace $X \setminus \text{St}(K, \mathcal{U}_\alpha)$ is uniformly discrete;
- (3) every finite open covering is uniform.

PROOF: $1 \Rightarrow 3$ Obvious.

$3 \Rightarrow 2$ By way of contradiction, assume that there exists a closed discrete subset $D = \{x_\alpha : \alpha < \omega_\mu\}$ of pairwise distinct limit points.

We shall prove that, for every α , one can choose $\beta(\alpha) \geq \alpha$ such that the collection $\mathcal{F} = \{\text{St}(x_\alpha, \mathcal{U}_{\beta(\alpha)}) : \alpha < \omega_\mu\}$ consists of pairwise disjoint subsets. We proceed by transfinite induction. Choose $\beta(0) \geq 0$ such that $\overline{\text{St}(x_0, \mathcal{U}_{\beta(0)})}$ is disjoint from $D \setminus \{x_0\}$. Let $\alpha > 0$ and $C_\alpha = \bigcup_{\gamma < \alpha} \overline{\text{St}(x_\gamma, \mathcal{U}_{\beta(\gamma)})}$. The set C_α is closed, because X is an ω_μ -additive topological space [1]. The set $C = C_\alpha \cup \{x_\gamma : \gamma > \alpha\}$ is closed and therefore there exists $\beta(\alpha) \geq \alpha$ such that the subset $\overline{\text{St}(x_\alpha, \mathcal{U}_{\beta(\alpha)})}$ is disjoint from C .

Choose $y_\alpha \in \text{St}(x_\alpha, \mathcal{U}_{\beta(\alpha)})$, $y_\alpha \neq x_\alpha$. The subset $F = \{y_\alpha : \alpha < \omega_\mu\}$ has no limit points and the open covering $\{X \setminus F, X \setminus D\}$ cannot be uniform, because the subsets F and D cannot be separated by a uniform covering.

What we still need to prove is that for every $\mathcal{U} \in \mathbf{B}$ the subspace $Y = X \setminus \text{St}(K, \mathcal{U})$ is uniformly discrete (notice that every subset of Y is closed in X). We argue by way of contradiction. Again using transfinite induction, it is easy to choose elements x_α, y_α such that $y_\alpha \in \text{St}(x_\alpha, \mathcal{U}_\alpha)$ and $x_\alpha \neq y_\beta$ for every α, β . Thus the open covering consisting of $X \setminus \{x_\alpha : \alpha < \omega_\mu\}$ and $X \setminus \{y_\alpha : \alpha < \omega_\mu\}$ cannot be uniform. This contradiction concludes the proof.

$2 \Rightarrow 1$

It is enough to prove that the trace on K of every open covering is a uniform covering of K . This trace can be refined by a covering of the form $\{\text{St}(\text{St}(x, \mathcal{U}_{\alpha(x)}), \mathcal{U}_{\alpha(x)}) : x \in K\}$. As K is ω_μ -compact, the covering $\{\text{St}(x, \mathcal{U}_{\alpha(x)}) : x \in K\}$ has a subcovering of power less than ω_μ , say $\{\text{St}(x_i, \mathcal{U}_{\alpha(x_i)}) : i < \delta\}$ for a suitable $\delta < \omega_\mu$. Thus $\{\text{St}(x, \mathcal{U}_\gamma) : x \in K\}$ where $\gamma = \sup\{\alpha(x_i) : i < \delta\}$ is the required uniform refinement (see [4, Theorem 4.3.31]). \square

Remark. It follows from the proof that the condition 2 of the above theorem can be strengthened as follows:

- (2') the set K of limit points is ω_μ -compact and every closed discrete subset of X is uniformly discrete.

Remark. It is well-known that the fine uniformity on a metrizable topological space X is a metric uniformity iff the set of limit points is compact (see for example [10]).

We can see that an analogous result holds for ω_μ -metrizable spaces: precisely, if the subset K of limit points of an ω_μ -metrizable space X is ω_μ -compact, then the fine uniformity is an ω_μ -metric uniformity.

Let

$$\mathbf{B} = \{\mathcal{U}_\alpha : \alpha < \omega_\mu\}$$

be a well ordered base for a compatible uniformity. For every α , consider the open covering

$$\mathcal{V}_\alpha = \{\{x\}, U : U \in \mathcal{U}_\alpha, U \cap K \neq \emptyset, x \in X \setminus \text{St}(K, \mathcal{U}_\alpha)\}.$$

It is easy to check that $\mathbf{C} = \{\mathcal{V}_\alpha : \alpha < \omega_\mu\}$ is a well ordered (by refinement) base which induces the fine uniformity (see Theorem 3).

REFERENCES

- [1] Artico G. and Moresco R., ω_μ -additive topological spaces, *Rend. Sem. Mat. Univ. Padova* **67** (1982), 131–141.
- [2] Atsuji M., Uniform continuity of continuous functions of metric spaces, *Pacific J. Math.* **8** (1958), 11–16.
- [3] Di Concilio A., Naimpally S.A., Uniform continuity in sequentially uniform spaces, *Acta Mathematica Hungarica* **61** 3–4 (1993) (to appear).
- [4] Engelking R., *General Topology*, Polish Scientific Publishers, Warsaw, 1977.
- [5] Isbell J.R., *Uniform Spaces*, Mathematical Surveys nr 12 AMS, Providence, Rhode Island, 1964.
- [6] Iwaiwa T., On uniform continuity of $C(X)$ (Japanese), *Sugaku Kenku Roku of Tokyo Kyoiku Daigaku* **2** (1955), 36–45.
- [7] Marconi U., On the uniform paracompactness, *Rend. Sem. Mat. Univ. Padova* **72** (1984), 101–105.
- [8] ———, On uniform paracompactness of the ω_μ -metric spaces, *Rend. Accad. Naz. Lincei* **75** (1983), 102–105.
- [9] Morita K., Paracompactness and product spaces, *Fund. Math.* **50** (1962), 223–236.
- [10] Rainwater J., Spaces whose finest uniformity is metric, *Pacific J. Math.* **9** (1959), 567–570.

DIPARTIMENTO DI MATEMATICA PURA E APPLICATA, UNIVERSITÀ DEGLI STUDI DI PADOVA,
I-35131 PADOVA, ITALY

(Received September 23, 1992)