

Kulumani M. Rangaswamy  
A property of  $B_2$ -groups

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 35 (1994), No. 4, 627--631

Persistent URL: <http://dml.cz/dmlcz/118704>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A property of $B_2$ -groups

K.M. RANGASWAMY

*Abstract.* It is shown, under ZFC, that a  $B_2$ -group has the interesting property of being  $\aleph_0$ -prebalanced in every torsion-free abelian group in which it is a pure subgroup. As a consequence, we obtain alternate proofs of some well-known theorems on  $B_2$ -groups.

*Keywords:* torsion-free abelian groups, Butler groups,  $B_2$ -groups,  $\aleph_0$ -prebalanced subgroups, completely decomposable groups, separative subgroups

*Classification:* Primary 20K20

### Introduction

All groups considered here, unless otherwise stated, are additively written torsion-free abelian groups. For unexplained terminology and notations, we refer to Fuchs [F-1]. A torsion-free abelian group  $G$  of infinite rank is called a  $B_2$ -group if, for some ordinal  $\tau$ ,  $G$  is the union of a continuous well-ordered ascending chain of pure subgroups,

$$(*) \quad 0 = G_0 \subset G_1 \subset \cdots \subset G_\alpha \subset \dots, \quad (\alpha < \tau) \dots$$

such that, for each  $\alpha < \tau$ ,  $G_{\alpha+1} = G_\alpha + B_\alpha$ , where  $B_\alpha$  is a finite rank pure subgroup of a completely decomposable group. Such groups  $B_\alpha$  are also called Butler groups. Recently Fuchs [F-2] made striking advances in the study of  $B_2$ -groups by employing the concept of  $\aleph_0$ -prebalancedness introduced in [BF]. In this note we prove that a  $B_2$ -group has the interesting property of being  $\aleph_0$ -prebalanced in every torsion-free group in which it is a pure subgroup. A noteworthy corollary is that a  $B_2$ -group  $A$  is a pure subgroup of index  $\leq \aleph_1$  in a  $B_1$ -group  $G$ , then  $G$  itself becomes a  $B_2$ -group. Taking  $A = 0$  leads to a well-known theorem ([DHR]) that a  $B_1$ -group of cardinality  $\leq \aleph_1$  is a  $B_2$ -group. An adaptation of our methods also leads to a direct and simple proof of a theorem of Hill and Megibben ([HM]) that completely decomposable groups are absolutely separative.

### Preliminaries

A torsion-free group  $G$  is called a  $B_1$ -group if  $\text{Bext}^1(G, T) = 0$  for all torsion groups  $T$ . (Here  $\text{Bext}^1$  denotes the subfunctor of  $\text{Ext}^1$  consisting of all the balanced extensions.) The chain of subgroups  $(*)$  defined above for a  $B_2$ -group  $G$  is called a  $B_2$ -filtration of  $G$ . Let  $A$  be a pure subgroup of a torsion-free group  $G$ .  $A$  is called decent (prebalanced) in  $G$  if whenever  $L/A$  is a finite rank (rank one) pure subgroup of  $G/A$ , then  $L = A + B$ , for some finite rank Butler group  $B$ .

$A$  is a TEP subgroup of  $G$  if, for any torsion group  $T$ , every homomorphism from  $A$  to  $T$  extends to a homomorphism from  $G$  to  $T$ .  $A$  is said to be  $\aleph_0$ -prebalanced ([BF]) in  $G$  if, for each  $g \in G \setminus A$  there is a countable subset  $\{a_1, a_2, \dots\} \subset A$  such that for each  $a \in A$ , there is an  $n < \omega$  with  $t(g+a) \leq \sup\{t(g+a_1), \dots, t(g+a_n)\}$  where  $t(x)$  denotes the type of  $x$ . In the last definition, if  $A$  satisfies the stronger condition that  $\chi(g+a) \leq \chi(g+a_i)$  for some  $i < \omega$ , then  $A$  is said to be separative (or in the terminology of [HM], separable) in  $G$ , where, as usual,  $\chi(x)$  denotes the characteristic of  $x$ . An  $\aleph_0$ -prebalanced chain for a group  $G$  is a continuous well-ordered ascending chain of  $\aleph_0$ -prebalanced subgroups

$$0 = G_0 \subset G_1 \subset \dots \subset G_\alpha \subset \dots G_\tau = G \quad (\text{for some ordinal } \tau)$$

where all the factors  $G_{\alpha+1}/G_\alpha$  are of rank one. A key result of Fuchs ([F-2, Corollary 2.4]) is that if  $G$  has an  $\aleph_0$ -prebalanced chain, then  $G$  is of the form  $G = C/K$ , where  $C$  is completely decomposable and  $K$  is a balanced  $B_2$ -subgroup. Another useful idea that we need from [BF] is the balanced-projective resolution of a group  $G$  relative to a pure subgroup  $A$ . To form this, consider all the rank-1 pure subgroups  $J_\alpha$  in  $G \setminus A$  and let  $C$  be the direct sum of all these  $J_\alpha$ 's. Then the map  $C \rightarrow B$  induced by the inclusion of the  $J_\alpha$  in  $G$  together with the inclusion of  $A$  in  $G$  induces a balanced exact sequence

$$0 \longrightarrow K \longrightarrow A \oplus C \longrightarrow G \longrightarrow 0$$

which is called the balanced-projective resolution of  $G$  relative to  $A$ . An important result of Bican-Fuchs ([BF, Theorem 3.2]) that we shall be using asserts that if  $G/A$  is countable, then  $A$  is  $\aleph_0$ -prebalanced in  $G$  exactly when  $K$  is a  $B_2$ -group. We shall also need a result from [R] that if  $A$  is a TEP subgroup of  $B$  and if both  $A$  and  $B$  are  $B_2$ -groups, then so is  $B/A$ . The reader is referred to [BF], [F-2] and [R] for background details.

**The results**

We shall begin with the following simple lemma.

**Lemma 1.** *Let  $A$  and  $S$  be subgroups of a torsion-free group  $G$ . If  $A \cap S$  is pure and decent in  $A$ , then  $S$  is pure and decent in  $A + S$ .*

PROOF: We first show that given any finite subset  $X$  of  $A + S$ , there is a finite rank Butler subgroup  $B$  such that  $B + S$  is pure in  $A + S$  and contains  $X$ . Without loss of generality, we may assume that  $X \subset A$ . By the decency of  $A \cap S$ , there is a finite rank Butler subgroup  $B$  of  $A$  such that  $B + (A \cap S)$  is pure in  $A$  and contains  $X$ . It is then readily seen that both  $B + S$  and  $S$  are pure in  $A + S$ . From this the decency of  $S$  follows. □

Bican and Fuchs [BF] showed, under  $V = L$ , that every  $B_1$ -group is “absolutely  $\aleph_0$ -prebalanced”, that is, it is an  $\aleph_0$ -prebalanced subgroup of every group in which it is a pure subgroup. The next theorem says that this holds for any  $B_2$ -group and we prove this under ZFC.

**Theorem 2.** *Every  $B_2$ -group is absolutely  $\aleph_0$ -prebalanced.*

PROOF: Let  $A$  be a  $B_2$ -group with an axiom-3 family  $\mathbb{C}$  of pure decent subgroups so chosen that for each  $Y \in \mathbb{C}$ ,  $A/Y$  is again a  $B_2$ -group (see [AH] for the construction of  $\mathbb{C}$ ). Suppose  $A$  is a pure subgroup of a torsion-free group  $B$  with  $B/A$  countable. Then  $B = A + S$ , where  $S$  is a countable pure subgroup. Moreover, by the usual back and-forth argument, we could assume that  $A \cap S = Y \in \mathbb{C}$ . By Lemma 1,  $S$  is decent and pure in  $B$ . Moreover,  $B/S \cong A/Y$  is a  $B_2$ -group. Since  $S$  is decent and countable, the pre-image of a  $B_2$ -filtration of  $B/S$  gives rise to an  $\aleph_0$ -prebalanced chain in  $B$ . In order to show that  $A$  is  $\aleph_0$ -prebalanced in  $B$ , consider a relative balanced-projective resolution (as explained in the Preliminaries)

$$0 \longrightarrow K \longrightarrow A \oplus X \longrightarrow B \longrightarrow 0$$

where  $X$  is completely decomposable. Let

$$0 \longrightarrow M \longrightarrow X' \longrightarrow A \longrightarrow 0$$

be a balanced-projective resolution of  $A$  with  $X'$  completely decomposable. Then the obvious epimorphism  $X' \oplus X \rightarrow A \oplus X$  induces the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & M & \xlongequal{\quad} & M & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & L & \longrightarrow & X' \oplus X & \longrightarrow & B \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & K & \longrightarrow & A \oplus X & \longrightarrow & B \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 
 \end{array}$$

Here all the rows and columns are balanced exact. Since  $B$  has an  $\aleph_0$ -prebalanced chain, Corollary 2.4 of [F-2] implies that  $L$  is a  $B_2$ -group. Since  $A \oplus X$  is a  $B_2$ -group, the middle column is TEP exact and, moreover, by [F-2] and [R],  $M$  is a  $B_2$ -group. Clearly the first column is now TEP exact and Theorem 3 of [R] then yields that  $K$  is also a  $B_2$ -group. An appeal to Theorem 3.2 of [BF] (alluded to in the Preliminaries) leads to the conclusion that  $A$  is  $\aleph_0$ -prebalanced in  $B$ .

□

**Corollary 3.** *Suppose  $A$  is a  $B_2$ -group which is a pure subgroup of a torsion-free group  $B$  with  $B/A$  having cardinality  $\leq \aleph_1$ . Then*

- (a)  $B$  has an  $\aleph_0$ -prebalanced chain and  $\text{Bext}^2(B, T) = 0$  for all torsion groups  $T$ .
- (b) If  $B$  is a  $B_1$ -group, then  $B$  is also a  $B_2$ -group.

PROOF: (a) Now  $B$  is a union of a smooth ascending chain of pure subgroups

$$(1) \quad A = A_0 \subset A_1 \subset \dots \subset A_\alpha \subset \dots, \alpha < \omega_1, \dots$$

where, for each  $\alpha$ ,  $A_{\alpha+1}/A_\alpha$  is countable. Since a countable extension of an absolutely  $\aleph_0$ -prebalanced subgroup is again absolutely  $\aleph_0$ -prebalanced, the chain (1) gives rise to a  $\aleph_0$ -prebalanced chain for  $B$ . By Corollary 2.3 of [F-2],  $\text{Bext}^2(B, T) = 0$ .

(b) Follows from the fact (Theorem 4.1 of [F-2]) that a  $B_1$ -group with an  $\aleph_0$ -prebalanced chain is a  $B_2$ -group. □

In Corollary 3 (b) if we take  $A = 0$ , then we obtain the following

**Corollary 4** ([DHR]). *A  $B_1$ -group of cardinality  $\leq \aleph_1$  is a  $B_2$ -group.*

**Corollary 5.** *If  $A$  is a pure  $B_2$ -subgroup of a finitely Butler group  $B$  with  $B/A$  countable, then  $B$  itself is a  $B_2$ -group.*

PROOF: Since  $B$  is finitely Butler, the countable subgroup  $S$  in the first part of the proof of Theorem 2 is Butler and decent in  $B$  with  $B/S$  a  $B_2$ -group. Clearly  $B$  is then a  $B_2$ -group. □

**Note:** The group  $\Pi Z$ , the direct product of  $\aleph_0$  copies of the group  $Z$  of integers, shows that Corollary 5 is false if  $B/A$  is uncountable.

If  $A$  is a completely decomposable group, then the subgroup  $S$  in the proof of Theorem 2 can actually be a direct summand, as the following lemma shows.

**Lemma 6.** *Suppose  $A$  is a completely decomposable group and is a pure subgroup of a torsion-free group  $B$  with  $B/A$  countable. Then  $B = A' \oplus S$ , where  $A' \subset A$  and  $S$  is countable.*

PROOF: Now  $B = A + X$ , where  $X$  is a suitable countable pure subgroup of  $B$ . Then we can write  $A = A' \oplus Y$ , where  $Y$  is countable and  $X \cap A \subset Y$ . If  $S = Y + X$ , then clearly  $B = A' + S$ . Moreover,  $A' \cap S = A' \cap A \cap S \subset A' \cap Y = 0$ , so that  $B = A' \oplus S$ . □

As an application we get a direct and simpler proof of theorem of Hill and Megibben ([HM]) that completely decomposable are absolutely separative.

**Theorem 7** ([HM]). *A completely decomposable group  $A$  is separative in every torsion-free group containing  $A$  as a pure subgroup.*

PROOF: Let  $A$  be a pure subgroup of a torsion-free group  $G$ . Let  $g \in G \setminus A$ . If  $B = \langle A, g \rangle^*$ , the pure subgroup generated by  $A$  and  $g$ , then by Lemma 6  $B = A' \oplus S$ ,  $A = A' \oplus C$ , where  $S$  is countable and  $C = A \cap S$ . Write  $g = a' + s$ , where  $a' \in A'$  and  $s \in S$ . Clearly,  $H = \{-a' + c : c \in C\}$  is a countable subset of  $A$ . We claim that for any  $a \in A$ , there is an  $h \in H$  such that  $\chi(g+a) \leq \chi(g+h)$ . Indeed if  $a = x + y$ , with  $x \in A'$  and  $y \in C$ , then we have  $\chi(g+a) = \chi(a' + s + x + y) = \chi((a' + x) + (s + y)) \leq \chi(s + y) = \chi(g + h)$ , where  $h = -a' + y \in L$ . Thus  $A$  is separative in  $G$ .  $\square$

#### REFERENCES

- [AH] Albrecht U., Hill P., *Butler groups of infinite rank and Axiom-3*, Czech. Math. J. **37** (1987), 293–309.
- [BF] Bican L., Fuchs L., *Subgroups of Butler groups*, to appear.
- [DHR] Dugas M., Hill P., Rangaswamy K.M., *Butler groups of infinite rank*, Trans. Amer. Math. Soc. **320** (1990), 643–664.
- [F-1] Fuchs L., *Infinite Abelian Groups, vol. 2*, Academic Press, New York, 1973.
- [F-2] ———, *Butler Groups of Infinite Rank*, to appear.
- [HM] Hill P., Megibben C., *Pure subgroups of torsion-free groups*, Trans. Amer. Math. Soc. **303** (1987), 765–778.
- [R] Rangaswamy K.M., *A homological characterization of abelian  $B_2$ -groups*, Proc. Amer. Math. Soc., to appear.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, COLORADO SPRINGS,  
CO 80933–7150, USA

(Received February 21, 1994)