

Peg Daniels

$\omega^\omega$ -directedness and a question of E. Michael

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 36 (1995), No. 1, 115--121

Persistent URL: <http://dml.cz/dmlcz/118737>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## $\omega^\omega$ -directedness and a question of E. Michael

PEG DANIELS

*Abstract.* We define  $\omega^\omega$ -directedness, investigate various properties to determine whether they have this property or not, and use our results to obtain easier proofs of theorems due to Laurence and Alster concerning the existence of a Michael space, i.e. a Lindelöf space whose product with the irrationals is not Lindelöf.

*Keywords:* Michael space, Lindelöf

*Classification:* Primary 54A35, 03E35; Secondary 54G20

### 1. Introduction

E. Michael used CH to construct a Lindelöf space  $Z$  whose product with the irrationals  $\omega^\omega$  is not Lindelöf. He asked if such an example can be found in ZFC [M].

Suppose there is such a (ZFC) space  $Z$ , and let  $X = Z \times \omega^\omega$ . Then  $X$  can be written as a union,  $X = \bigcup_{f \in \omega^\omega} X_f$ , where each  $X_f$  is a closed Lindelöf space, and  $f \leq g$  implies  $X_f \subset X_g$ . To do this, let  $X_f = Z \times \{g \in \omega^\omega : g \leq f\}$  (where  $g \leq f$  means  $\forall n(g(n) \leq f(n))$ ). Each  $\{g \in \omega^\omega : g \leq f\}$  is compact, so each  $X_f$  is Lindelöf.

Let us call a property  $P$   $\omega^\omega$ -directed if a space  $X$  has property  $P$  whenever  $X = \bigcup_{f \in \omega^\omega} X_f$ , where  $X_f$  is a closed subspace of  $X$  having property  $P$ , and  $f \leq g$  implies  $X_f \subset X_g$ .

In this paper we investigate various properties to determine whether they are  $\omega^\omega$ -directed, and use our results to obtain easier proofs of the following theorems due to Laurence and Alster.

**Theorem** (Laurence). *If  $\mathfrak{b} > \omega_1$ , there is no concentrated Michael space.*

**Theorem** (Alster). *If  $\mathfrak{b} > \omega_1$ , and  $X$  is Lindelöf, then  $X \times \omega^\omega$  is  $\omega_1$ -compact; furthermore if  $X \times \omega^\omega$  is metalindelöf, then  $X \times \omega^\omega$  is Lindelöf.*

### 2. $\omega^\omega$ -directedness and Michael’s question

Clearly the property of having cardinality  $\leq \mathfrak{c}$  is  $\omega^\omega$ -directed. What about the property of being countable? Suppose  $X = \bigcup_{f \in \omega^\omega} X_f$ , each  $X_f$  closed in  $X$  and  $|X_f| \leq \omega$ . Suppose  $X$  is not countable. Let  $\{x_\alpha : \alpha < \omega_1\} \subset X$ , and for each  $\alpha < \omega_1$ , let  $f_\alpha \in \omega^\omega$  be such that  $x_\alpha \in X_{f_\alpha}$ . Now note that if  $\langle f_\alpha : \alpha < \omega_1 \rangle$  is bounded in  $\leq^*$ , i.e. there is an  $f \in \omega^\omega$  such that  $\forall \alpha < \omega_1 \{n : f_\alpha > f(n)\}$  finite,

then  $\forall \alpha < \omega_1 \exists g_\alpha =^* f$  (i.e.  $\{n : g_\alpha(n) \neq f(n)\}$  is finite) with  $f_\alpha \leq g_\alpha$ ; there are only countably many such  $g_\alpha$ , so we would have a  $g \in \omega^\omega$  and an uncountable set  $S \subset \omega_1$  such that  $\forall \alpha \in S, f_\alpha \leq d$ ; then  $\{x_\alpha : \alpha \in S\} \subset \bigcup_{\alpha \in S} X_{f_\alpha} \subset X_g$ , but as  $X_g$  is countable we would have a contradiction. Thus if every  $\omega_1$ -sequence from  $\omega^\omega$  is bounded, the property of being countable is  $\omega^\omega$ -directed. We shall see momentarily that the converse is also true.

**Definition 2.1.** For  $f, g \in \omega^\omega$ ,  $f \leq^* g$  if and only if  $\{n : f(n) > g(n)\}$  is finite. □

**Definition 2.2.** A sequence  $\langle f_\alpha : \alpha < \kappa \rangle$  from  $\omega^\omega$  is *unbounded* if and only if there is no  $f \in \omega^\omega$  such that  $\forall \alpha < \kappa (f_\alpha \leq^* f)$ ; if a sequence is not unbounded, we say it is *bounded*. □

**Definition 2.3.**  $\mathfrak{b}$  is the least cardinal  $\kappa$  such that there is an unbounded sequence from  $\omega^\omega$  of cardinality  $\kappa$ . □

$\mathfrak{b}$  is a regular cardinal.

**Proposition 2.4.** *The property of being countable is  $\omega^\omega$ -directed if and only if  $\mathfrak{b} > \omega_1$ .*

PROOF: We have seen in previous paragraphs that if  $\mathfrak{b} > \omega_1$ , then the property of being countable is  $\omega^\omega$ -directed.

Suppose that  $\mathfrak{b} = \omega_1$ . We construct a space  $X = \bigcup_{f \in \omega^\omega} X_f$ , where each  $X_f$  is closed and countable, but  $X$  is uncountable.

Let  $\langle g_\alpha : \alpha < \omega_1 \rangle$  be unbounded. Without loss of generality,  $\alpha < \beta$  implies that  $g_\alpha <^* g_\beta$  [vD]. Our space  $X$  is  $\{g_\alpha : \alpha < \omega_1\}$ , with the subspace topology of  $\omega^\omega$ .  $\forall f \in \omega^\omega$ , let  $G_f = \{g \in X : g \leq f\}$ . We first show  $G_f$  is countable. Let  $\alpha < \omega_1$  be such that  $g_\alpha \not\leq^* f$ . Suppose  $\beta > \alpha$ .  $g_\alpha <^* g_\beta$ , so if  $g_\beta \leq f$ , then  $g_\alpha \leq^* f$ , a contradiction. We have shown that if  $\beta > \alpha$ , then  $g_\beta \not\leq f$ , that is, that  $G_f \subset \{g_\beta : \beta \leq \alpha\}$ . Thus  $G_f$  is countable. Finally we show each  $G_f$  is closed. Suppose  $g_\alpha$  is a limit point of  $G_f$ , not in  $G_f$ .  $g_\alpha \not\leq f$ . Let  $n$  be such that  $g_\alpha(n) > f(n)$ . Let  $g_\beta \in G_f \cap [g_\alpha]_{n+1}$ .  $g_\beta \in G_f$  implies  $g_\beta(n) \leq f(n)$ , but  $g_\beta \in [g_\alpha]_{n+1}$  implies  $g_\beta(n) = g_\alpha(n) > f(n)$ , a contradiction. □

By a similar argument, we have

**Theorem 2.5.** *For  $cf\kappa > \omega$ , the property of having cardinality  $< \kappa$  is  $\omega^\omega$ -directed if and only if  $\mathfrak{b} > \kappa$ .* □

As we mentioned in the introduction, Michael showed under CH that there is a Lindelöf  $Z$  such that  $Z \times \omega^\omega$  is not Lindelöf. For this example, we need some terminology. A space  $X$  is *concentrated about*  $A \subset X$  if each neighborhood of  $A$  contains all but countably many points of  $X$ . Under CH, there is an uncountable  $B \subset \omega^\omega$  such that  $B \cup \mathbb{Q}$ , as a subspace of  $\mathbb{R}$ , is concentrated about  $\mathbb{Q}$ .

Let  $Z = B \cup \mathbb{Q}$ , with the points of  $B$  isolated. Then  $X_f = Z \times \{g : g \leq f\}$  is a closed Lindelöf subspace of  $Z \times \omega^\omega$ , but  $Z \times \omega^\omega$  is not Lindelöf because  $\{(g, g) : g \in B\}$  is a closed discrete subset.

In fact, there is such a  $B$  if and only if  $\mathfrak{b} = \omega_1$ .

Laurence [L] defines a *concentrated Michael space* to be a Lindelöf space  $X$  that is concentrated about a closed subset  $A$ , and having the property that  $A \times \omega^\omega$  is Lindelöf (or equivalently, normal), but  $X \times \omega^\omega$  is not.

He constructs such a space assuming  $\mathfrak{b} = \omega_1$ , and shows there is no such space if  $\mathfrak{b} > \omega_1$ . Here we show his latter result to be a corollary to Proposition 2.4.

**Corollary 2.6** ( $\mathfrak{b} > \omega_1$ ). *There is no concentrated Michael space.*

PROOF: Suppose  $X$  is a Lindelöf space concentrated about a closed subset  $A$  such that  $A \times \omega^\omega$  is Lindelöf. We show  $X \times \omega^\omega$  is Lindelöf, assuming  $\mathfrak{b} > \omega_1$ . Suppose  $\mathcal{U}$  is an open cover of  $X \times \omega^\omega$ . Let  $\mathcal{V} \subset \mathcal{U}$  be countable, with  $\cup \mathcal{V}$  covering  $A \times \omega^\omega$ . Let  $H = (X \times \omega^\omega) \setminus \cup \mathcal{V}$ .

**Claim.**  $\pi_2(H)$  is countable.

Suppose that the claim holds. Then  $H$  is the countable union of Lindelöf sets, hence Lindelöf, and so also may be covered by a countable subcollection of  $\mathcal{U}$ .

PROOF OF CLAIM: Suppose  $\pi_2(H)$  is uncountable. For each  $i \in \pi_2(H)$ , let  $x_i \in X \setminus A$  be such that  $(x_i, i) \in H$ . Let  $K = \overline{\{(x_i, i) : i \in \pi_2(H)\}}$ .  $K \cap (A \times \omega^\omega) = \emptyset$ . We now show  $|\pi_2(H) \cap \mathbb{P}_g| \leq \omega$ , for each  $g \in \omega^\omega$ , where  $\mathbb{P}_g = \{f \in \omega^\omega : f \leq g\}$ . Suppose  $|\pi_2(H) \cap \mathbb{P}_g| \geq \omega_1$ . For each  $a \in A$ , let  $\mathcal{W}_a = \{U_{n,\alpha} \times V_{n,\alpha} : n \leq m_a\}$  be a finite basic open cover of  $\{a\} \times \mathbb{P}_g$ , each element of which is disjoint from  $K$ .  $\mathcal{W} = \{\bigcap_{n \leq m_a} U_{n,a} : a \in A\}$  covers  $A$ . For each  $\alpha \in \omega_1$ , let  $i_\alpha \in \pi_2(H) \cap \mathbb{P}_g$ . Since  $\mathcal{W}$  must cover all but countably many points of  $X$ , let  $\alpha \in \omega_1$  be such that  $x_{i_\alpha} \in \cup \mathcal{W}$ . Let  $a_\alpha$  be such that  $x_{i_\alpha} \in \bigcap_{n \leq m_{a_\alpha}} U_{n,a_\alpha}$ .  $(a_\alpha, i_\alpha) \in \cup \mathcal{W}_{a_\alpha}$ , so let  $n \leq m_{a_\alpha}$  be such that  $(a_\alpha, i_\alpha) \in U_{n,a_\alpha} \times V_{n,a_\alpha}$ . Then  $(x_{i_\alpha}, i_\alpha)$  is in  $(U_{n,a_\alpha} \times V_{n,a_\alpha}) \cap K$ , a contradiction. So  $\pi_2(H) \cap \mathbb{P}_g$  is countable. By Proposition 2.4, being countable is  $\omega^\omega$ -directed under  $\mathfrak{b} > \omega_1$ , and so  $\pi_2(H)$  is countable.  $\square$

**Theorem 2.7.** *For  $cf \kappa > \omega$ , the property of being  $\kappa$ -compact is  $\omega^\omega$ -directed if and only if  $\mathfrak{b} > \kappa$ .*

PROOF: Since not being  $\kappa$ -compact means there is a subset  $\{x_\alpha : \alpha < \kappa\}$  with no limit point, it is easy to see how to produce an argument similar to that in Theorem 2.6 if  $\mathfrak{b} > \kappa$ .  $\square$

A lemma we will find useful for the converse, as well as in other applications is:

**Lemma 2.8.** *Let  $\langle f_\alpha : \alpha < \mathfrak{b} \rangle$  be unbounded. For each  $f \in \omega^\omega$ , let  $\alpha(f)$  be the least  $\alpha$  such that  $f_\alpha \not\prec^* f$ . Then  $f \leq g$  implies  $\alpha(f) \leq \alpha(g)$ , and  $\sup\{\alpha(f) : f \in \omega^\omega\} = \mathfrak{b}$ .*

PROOF OF LEMMA 2.8: Suppose  $f \leq g$  and  $\alpha(f) > \alpha(g)$ . By definition of  $\alpha(f)$ ,  $f_{\alpha(g)} <^* f$ . But then  $f_{\alpha(g)} <^* g$ , a contradiction.  $\square$

PROOF OF THEOREM 2.7, CONTINUED: Suppose  $\mathfrak{b} = \kappa$ . Let  $\langle f_\alpha : \alpha < \mathfrak{b} \rangle$  be unbounded and define  $\alpha(f)$  for each  $f \in \omega^\omega$  as in the lemma; let  $X = \kappa$  and let  $X_f = \alpha(f) + 1$ . Clearly  $X$  witnesses the non- $\omega^\omega$ -directedness of  $\kappa$ -compactness.  $\square$

Alster [A] has shown that (a)  $\mathfrak{b} > \omega_1$  implies that if  $X$  is Lindelöf, then  $X \times \omega^\omega$  does not contain an uncountable closed discrete set, and hence that (b) if  $\mathfrak{b} > \omega_1$ ,  $X$  is Lindelöf, and  $X \times \omega^\omega$  is metalindelöf, then  $X \times \omega^\omega$  is Lindelöf. His result (a) follows easily from our work:

**Corollary 2.9** ( $\mathfrak{b} > \omega_1$ ). *If  $X$  is Lindelöf, then  $X \times \omega^\omega$  does not contain an uncountable closed discrete set, i.e.  $X \times \omega^\omega$  is  $\omega_1$ -compact.*

PROOF: Suppose  $X$  is Lindelöf. As each  $X \times \{g \in \omega^\omega : g \leq f\}$  is Lindelöf, each is  $\omega_1$ -compact; this property is  $\omega^\omega$ -directed under  $\mathfrak{b} > \omega_1$  (Theorem 2.7), so  $X \times \omega^\omega = \bigcup_{f \in \omega^\omega} (X \times \{g \in \omega^\omega : g \leq f\})$  is  $\omega_1$ -compact.  $\square$

### 3. Other $\omega^\omega$ -directed properties

Very few properties seem to be  $\omega^\omega$ -directed, even consistently. It is easy to check that the properties of being  $T_0$ ,  $T_1$ , or connected are  $\omega^\omega$ -directed. Besides having a cardinality  $\leq \kappa$  and being  $\kappa$ -compact, the properties of being hereditarily Lindelöf and hereditarily of density  $\kappa$  also depend on the size of  $\mathfrak{b}$ .

**Theorem 3.1.** *For  $cf\kappa > \omega$ , the property of being hereditarily  $k$ -Lindelöf is  $\omega^\omega$ -directed if and only if  $\mathfrak{b} > \kappa$ .*

PROOF: Since a space  $X$  is not hereditarily  $\kappa$ -Lindelöf if and only if there is a subset  $\{x_\alpha : \alpha < \kappa\} \subset X$  such that for each  $\alpha < \kappa$ ,  $x_\alpha \notin \overline{\{x_\beta : \beta > \alpha\}}$ , the argument proceeds as before for  $\mathfrak{b} > \kappa$ . For  $\mathfrak{b} = \kappa$ , the  $X = \kappa$  of Theorem 2.7 again works.  $\square$

**Theorem 3.2.** *For  $cf\kappa > \omega$ , the property of being hereditarily of density  $\leq \kappa$  is  $\omega^\omega$ -directed if and only if  $\mathfrak{b} > \kappa$ .*

PROOF: Note that a space  $X$  is not hereditarily of density  $\leq \kappa$  if and only if there is a set  $\{x_\alpha : \alpha < \kappa\}$  such that for each  $\alpha < \kappa$ ,  $x_\alpha \notin \overline{\{x_\beta : \beta < \alpha\}}$ .  $\square$

### 4. ZFC counterexamples

We found easy counterexamples to many properties, although the problem may be more difficult if we require higher separation axioms. We first present our easy counterexamples, and then give a counterexample for separability.

Let  $\alpha(\mathfrak{b})$  be the set  $\alpha$  with the topology as follows:  $U$  is open in  $\alpha(\mathfrak{b})$  if and only if  $|\alpha \setminus U| < \mathfrak{b}$ . If  $\alpha < \mathfrak{b}$ , then  $\alpha(\mathfrak{b})$  is discrete. Let  $X = \mathfrak{b}(\mathfrak{b})$  and  $X_f = \alpha(f)(\mathfrak{b})$ . Then  $X$  witnesses that  $T_2$  is not  $\omega^\omega$ -directed. It also gives a  $T_1$

counterexample showing the hereditary disconnectedness, 0-dimensionality, and strong 0-dimensionality properties are not  $\omega^\omega$ -directed.

$PR_{\leq 2}(\mathfrak{b}(\mathfrak{b}))$  denotes the Pixley-Roy topology on  $[\mathfrak{b}(\mathfrak{b})]^{\leq 2}$ , i.e. the basis is sets of the form  $[A, U] = \{B \in [\mathfrak{b}(\mathfrak{b})]^{\leq 2} : A \subset B \subset U\}$ , where  $A \in [\mathfrak{b}(\mathfrak{b})]^{\leq 2}$  and  $U$  is open in  $\mathfrak{b}(\mathfrak{b})$ . This space is normal, since the singletons can be mutually separated. With this as our space  $X$  and  $X_f = PR_{\leq 2}(\alpha(f)(\mathfrak{b}))$ , we have a counterexample showing that the Fréchet, sequential,  $k$ -space, closed sets are  $G_\delta$ , metric, extreme disconnectedness, character  $\leq \mathfrak{b}$ , and density  $\leq \mathfrak{b}$  properties are not  $\omega^\omega$ -directed.

Now let  $X = \mathfrak{b} + 1$ , with the usual topology on  $\mathfrak{b}$ , and a basic open set about  $\mathfrak{b}$  having the form  $(\alpha, \mathfrak{b}] \setminus \{\gamma > \alpha : \gamma \text{ is a limit ordinal}\}$ .  $X_f = (\alpha(f) + 1) \cup \{\mathfrak{b}\}$ .  $X$  witnesses that  $T_3$  is not  $\omega^\omega$ -directed.

Let  $X$  be  $(\mathfrak{b} + 1) \times (\omega + 1) \setminus \{(\mathfrak{b}, \omega)\}$ . (A ‘‘Tychonoff plank’’ counterexample.)  $X$  is  $T_{3\frac{1}{2}}$ .  $X$  witnesses that  $T_4$ , Lindelöf, paracompactness, metacompactness, subparacompactness, and collectionwise normality properties are not  $\omega^\omega$ -directed.

If we let  $X$  be as above, except isolate all the points not on the edges, we have a  $T_{3\frac{1}{2}}$  counterexample witnessing that the countable paracompactness, collectionwise Hausdorffness, and hereditary normality properties are not  $\omega^\omega$ -directed.

The irrationals,  $\omega^\omega$ , with  $X_f = \{g \in \omega^\omega : g \leq f\}$ , show that compactness properties (including local, countable, pseudo-, and sequential) are not  $\omega^\omega$ -directed.

For our counterexample for separability we need the following definition:

**Definition 4.1.** If  $\mathcal{A} \subset \mathcal{P}(\omega)$ ,  $\mathcal{A}$  is an independent family (i.f.) if and only if, whenever  $m, n \in \omega$  and  $a_1, \dots, a_m, b_1, \dots, b_n$  are distinct members of  $\mathcal{A}$ ,  $|a_1 \cap \dots \cap a_m \cap (\omega \setminus b_1) \cap \dots \cap (\omega \setminus b_n)| = \omega$ .

Clearly ‘‘ $\omega$ ’’ can be replaced by any countable set. Note that there is always an i.f. of size  $\mathfrak{c}$  (see [K]), and hence of size  $\mathfrak{b}$ .

**Example 4.2.** For each limit  $\lambda < \mathfrak{b}$ , let  $\lambda^* = (\lambda + \omega) \setminus \lambda$ . Let  $\{F_{\lambda, \alpha} : \alpha < \mathfrak{b}\}$  be an i.f. of subsets of  $\lambda^*$ . For each  $\alpha < \mathfrak{b}$ , let  $\lambda_\alpha < \mathfrak{b}$  be such that  $\alpha \in \lambda_\alpha^*$ ; let  $U(\alpha) = \{\alpha\} \cup \bigcup \{F_{\tau, \alpha} : \tau \geq \lambda_\alpha + \omega\}$ . Let  $\{U(\alpha) \setminus U(\beta) : \alpha \neq \beta\}$  be a subbasis for a topology on  $\mathfrak{b}$ . This is our space  $X$ . Note that  $\alpha \in U(\alpha) \setminus U(\beta)$  for any  $\beta > \alpha$ . Also, if  $\gamma \notin U(\alpha) \setminus U(\beta)$ , then either (1)  $\gamma \notin U(\alpha)$ , in which case  $U(\gamma) \setminus U(\alpha)$  is an open set about  $\gamma$  missing  $U(\alpha) \setminus U(\beta)$ , or (2)  $\gamma \in U(\alpha) \cap U(\beta)$ , in which case for a sufficiently large  $\delta$ ,  $[U(\alpha)] \cap [U(\beta) \setminus U(\delta)]$  is an open set about  $\gamma$  missing  $U(\alpha) \setminus U(\beta)$ ; thus the elements of the subbasis are closed, so  $X$  is completely regular.

Let  $\langle f_\alpha : \alpha < \mathfrak{b} \rangle$  be an unbounded family in  $\omega^\omega$ , and  $\alpha(f)$  defined as before for  $f \in \omega^\omega$ . Let  $X_f = \alpha(f) + \omega$ .  $X_f$  is closed since if  $\beta \geq \alpha(f) + \omega$ ,  $U(\beta)$  misses  $X_f$ .

We wish to show that each  $X_f$  is separable. Let  $\lambda$  be the greatest limit ordinal  $\leq \alpha(f)$ . We claim  $\lambda^*$  is dense in  $X_f$ .

Consider a  $\bigcap_{i \leq n} [U(\alpha_i) \setminus U(\beta_i)] = U$ , such that  $U \cap X_f \neq \emptyset$ . (Note each  $\alpha_i \in X_f$ .) We wish to show  $U \cap \lambda^* \neq \emptyset$ . We first show  $\lambda \geq \lambda_{\alpha_i} + \omega$  for each  $i$ . Let  $\gamma$  be a point in  $U \cap X_f$ . If  $\gamma \in \lambda^*$ , we are done, so assume  $\gamma < \lambda$ . Then  $\lambda_\gamma < \lambda$ . If  $\gamma = \alpha_i$ , then  $\lambda_\gamma = \lambda_{\alpha_i}$ , so  $\lambda_{\alpha_i} + \omega = \lambda_\gamma + \omega \leq \lambda$ . On the other hand, if  $\gamma \neq \alpha_i$ , then since  $\gamma \in U(\alpha_i)$  there is a  $\tau \geq \lambda_{\alpha_i} + \omega$  with  $\gamma \in F_{\tau, \alpha_i} \subset \tau^*$ , so  $\lambda_\gamma = \tau$ .  $\lambda_{\alpha_i} + \omega \leq \tau < \tau + \omega = \lambda_\gamma + \omega \leq \lambda$ . In every case,  $\lambda \geq \lambda_{\alpha_i} + \omega$ . This means that for each  $i$ ,  $F_{\lambda, \alpha_i} \subset U(\alpha_i)$ . It is easy to see then that  $F_{\lambda, \alpha_i} \setminus [F_{\lambda, \beta_i} \cup \{\beta_i\}] \subset U(\alpha_i) \setminus U(\beta_i)$ . Since  $|\bigcap_{i \leq n} [F_{\lambda, \alpha_i} \setminus F_{\lambda, \beta_i}]| = \omega$  by the i.f. property,  $(\bigcap_{i \leq n} [F_{\lambda, \alpha_i} \setminus F_{\lambda, \beta_i}]) \setminus \{\beta_i : i \leq n\} \neq \emptyset$ , and an element of this set is in  $U \cap \lambda^*$ . So  $X_f$  is separable.

$X$ , however, is not separable, since if  $C$  is a countable subset of  $X$  (or even of cardinality  $< \mathfrak{b}$ ), there is a limit  $\lambda < \mathfrak{b}$  such that  $C \subset \lambda$ , and for  $\alpha(f) \geq \lambda$ ,  $\bar{C} \subset X_f$ .  $\square$

### 5. $\mathfrak{b} = \omega_1$

If  $\mathfrak{b} = \omega_1$ ,  $\omega_1$  itself is a simpler counterexample to separability, where  $X_f = \alpha(f) + 1$ . In this case it also provides a counterexample for perfect normality, having a countable base, Lindelöfness (a normal example), hereditary Lindelöfness, paracompactness and metacompactness (a normal example), and hereditary separability.

$PR_{<\omega}(\omega_1(\omega))$  (with underlying set  $[\omega_1]^{<\omega}$ ) provides a completely regular counterexample to hereditary paracompactness, under  $\mathfrak{b} = \omega_1$ .

Bing's example  $H$  provides a normal counterexample to collectionwise normality under  $\mathfrak{b} = \omega_1$ . Here,  $X \subseteq 2^{\mathcal{P}(\omega_1)}$  is defined as follows. For each  $\alpha < \omega_1$ , let  $f_\alpha \in 2^{\mathcal{P}(\omega_1)}$  be defined by  $f_\alpha(A) = 1$  if and only if  $\alpha \in A$ . Let  $F = \{f_\alpha : \alpha < \omega_1\}$ . Recalling that the support of an  $x \in 2^{\mathcal{P}(\omega_1)}$  is defined to be  $\{A \in \mathcal{P}(\omega_1) : x(A) = 1\}$ , let  $Y = \{x \in 2^{\mathcal{P}(\omega_1)} \setminus F : |\text{supp}(x)| < \omega\}$ . Let  $X = Y \cup F$ , where the points of  $Y$  are isolated, and the points of  $F$  have their usual (subspace) neighborhoods in  $2^{\mathcal{P}(\omega_1)}$ .  $F$  is close discrete in  $X$ , and by standard arguments, cannot be separated. For each  $f \in \omega^\omega$ , let  $X_f = \{f_\alpha : \alpha < \alpha(f)\} \cup \{x \in Y : \text{supp}(x) \subset \alpha(f)\}$ .  $X_f$  is closed, since if  $f_\beta$  is a limit point,  $x \in X_f \cap [f_\beta \mid \{\beta\}]$  implies  $x(\beta) = 1$ , which implies  $\beta \in \text{supp}(x)$ , and since  $x \in X_f$  implies  $\text{supp}(x) \subset \alpha(f)$ , we have  $\beta < \alpha(f)$  and so  $f_\beta \in X_f$ .  $X_f$  is collectionwise normal, since any closed discrete subset of  $\{f_\alpha : \alpha < \alpha(f)\}$  is countable.

### REFERENCES

- [A] Alster K., *On the product of a Lindelöf space with the space of irrationals under Martin's Axiom*, Proc. Amer. Math. Soc. **110** (1990), 543–547.
- [K] Kunen K., *Set Theory*, North-Holland, Amsterdam, 1980.
- [L] Laurence L.B., *The influence of a small cardinal on the product of a Lindelöf space and the irrationals*, Proc. Amer. Math. Soc. **110** (1990), 535–542.

- [M] Michael E., *Paracompactness and the Lindelöf property in finite and countable Cartesian products*, *Comp. Math.* **23** (1971), 199–214.
- [vD] van Douwen E.K., *The integers and topology*, in *Handbook of Set-Theoretic Topology*, ed. K. Kunen and J.E. Vaughan, North Holland, Amsterdam, 1984.

DEPARTMENT OF MATHEMATICS, AUBURN UNIVERSITY, ALABAMA 36849-5310, USA

(Received August 2, 1994)