

Donal O'Regan

Differential equations at resonance

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 36 (1995), No. 4, 673--694

Persistent URL: <http://dml.cz/dmlcz/118795>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Differential equations at resonance

DONAL O'REGAN

*Abstract.* New existence results are presented for the two point singular “resonant” boundary value problem  $\frac{1}{p}(py')' + ry + \lambda_m qy = f(t, y, py')$  a.e. on  $[0, 1]$  with  $y$  satisfying Sturm Liouville or Periodic boundary conditions. Here  $\lambda_m$  is the  $(m + 1)^{st}$  eigenvalue of  $\frac{1}{pq}[(pu')' + rpu] + \lambda u = 0$  a.e. on  $[0, 1]$  with  $u$  satisfying Sturm Liouville or Periodic boundary data.

*Keywords:* boundary value problems, resonance, existence

*Classification:* 34B15

### 1. Introduction

In this paper we derive some existence results for the second order equation

$$(1.1) \quad \frac{1}{p(t)}(p(t)y'(t))' + r(t)y(t) + \lambda_m q(t)y(t) = f(t, y(t), p(t)y'(t)) \quad \text{a.e. on } [0, 1]$$

with  $y$  satisfying either

(i) (Sturm Liouville)

$$(SL) \quad \begin{cases} -\alpha y(0) + \beta \lim_{t \rightarrow 0^+} p(t)y'(t) = 0, & \alpha \geq 0, \beta \geq 0, \alpha^2 + \beta^2 > 0 \\ \alpha y(1) + b \lim_{t \rightarrow 1^-} p(t)y'(t) = 0, & a \geq 0, b \geq 0, a^2 + b^2 > 0 \end{cases}$$

or

(ii) (Periodic)

$$(P) \quad \begin{cases} y(0) = y(1) \\ \lim_{t \rightarrow 0^+} p(t)y'(t) = \lim_{t \rightarrow 1^-} p(t)y'(t). \end{cases}$$

*Remarks.* (i)  $\lambda_m$  will be described later.

(ii) The Neumann condition  $\lim_{t \rightarrow 0^+} p(t)y'(t) = \lim_{t \rightarrow 1^-} p(t)y'(t) = 0$  is included in (SL) with  $\alpha = a = 0$ .

(iii) If a function  $u \in C[0, 1] \cap C^1(0, 1)$  with  $pu' \in C[0, 1]$  satisfies boundary condition (i) we write  $u \in (SL)$ . A similar remark applies for the boundary condition (ii).

Throughout the paper  $p \in C[0, 1] \cap C^1(0, 1)$  together with  $p > 0$  on  $(0, 1)$ . Also  $pf : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  is an  $L^1$ -Carathéodory function. By this we mean:

- (i)  $t \rightarrow p(t)f(t, y, q)$  is measurable for all  $(y, q) \in \mathbf{R}^2$ ;
- (ii)  $(y, q) \rightarrow p(t)f(t, y, q)$  is continuous for a.e.  $t \in [0, 1]$ ;

- (iii) for any  $r > 0$  there exists  $h_r \in L^1[0, 1]$  such that  $|p(t)f(t, y, q)| \leq h_r(t)$  for a.e.  $t \in [0, 1]$  and for all  $|y| \leq r, |q| \leq r$ .

For notational purposes let  $w$  be a weight function. By  $L^1_w[0, 1]$  we mean the space of functions  $u$  such that  $\int_0^1 w(t)|u(t)| dt < \infty$ .  $L^2_w[0, 1]$  denotes the space of functions  $u$  such that  $\int_0^1 w(t)|u(t)|^2 dt < \infty$ ; also for  $u, v \in L^2_w[0, 1]$  define  $\langle u, v \rangle = \int_0^1 w(t)u(t)\overline{v(t)} dt$ . Let  $AC[0, 1]$  be the space of functions which are absolutely continuous on  $[0, 1]$ .

Before we discuss the boundary value problem (1.1) and its appropriate literature we first gather together some facts on second order differential equations ([12], [16]). Consider the linear equation

$$(1.2) \quad \begin{cases} \frac{1}{p}(py')' + \tau y = g(t) & \text{a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}). \end{cases}$$

By a solution to (1.2) we mean a function  $y \in C[0, 1] \cap C^1(0, 1)$  with  $py' \in AC[0, 1]$  which satisfies the differential equation in (1.2) a.e. on  $[0, 1]$  and the stated boundary conditions.

**Theorem 1.1.** *Suppose*

$$(1.3) \quad p \in C[0, 1] \cap C^1(0, 1) \text{ with } p > 0 \text{ on } (0, 1) \text{ and } \int_0^1 \frac{ds}{p(s)} < \infty$$

and

$$(1.4) \quad \tau, g \in L^1_p[0, 1]$$

are satisfied. If

$$(1.5) \quad \begin{cases} \frac{1}{p}(py')' + \tau y = 0 & \text{a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

has only the trivial solution, then (1.2) has exactly one solution  $y$  given by

$$y(t) = d_0u_1(t) + d_1u_2(t) + \int_0^t \frac{[u_2(t)u_1(s) - u_1(t)u_2(s)]}{W(s)}g(s) ds$$

where  $u_1$  is the unique solution to

$$\begin{cases} \frac{1}{p}(pu')' + \tau u = 0 \text{ a.e. on } [0, 1] \\ u(0) = 1, \lim_{t \rightarrow 0^+} p(t)u'(t) = 0 \end{cases}$$

and  $u_2$  is the unique solution to

$$\begin{cases} \frac{1}{p}(pu')' + \tau u = 0 \text{ a.e. on } [0, 1] \\ u(0) = 0, \lim_{t \rightarrow 0^+} p(t)u'(t) = 1 \end{cases}$$

and  $d_0$  and  $d_1$  are uniquely determined from the boundary condition;  $W$  of course denotes the Wronskian. In fact

$$y(t) = \int_0^1 G(t, s)g(s) ds$$

with

$$G(t, s) = \begin{cases} \frac{y_1(s)y_2(t)}{W(s)}, & 0 < s \leq t \\ \frac{y_1(t)y_2(s)}{W(s)}, & t \leq s < 1 \end{cases}$$

where  $y_1$  and  $y_2$  are the two “usual” linearly independent solutions i.e. choose  $y_1 \neq 0, y_2 \neq 0$  so that  $y_1, y_2$  satisfy  $\frac{1}{p}(py')' + \tau y = 0$  a.e. on  $[0, 1]$  with  $y_1$  satisfying the first boundary condition and  $y_2$  satisfying the second boundary condition.

We now state an existence principle ([16]), which was established using fixed point methods, for the second order nonresonant boundary value problem

$$(1.6) \quad \begin{cases} \frac{1}{p}(py')' + \tau y = f(t, y, py') \text{ a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}). \end{cases}$$

**Theorem 1.2.** Let  $pf : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  be an  $L^1$ -Carathéodory function and assume (1.3) and

$$(1.7) \quad \tau \in L^1_p[0, 1]$$

hold. In addition suppose (1.5) has only the trivial solution. Now assume there is a constant  $M_0$ , independent of  $\lambda$ , with

$$\|y\|_* = \max\{\sup_{[0,1]} |y(t)|, \sup_{(0,1)} |p(t)y'(t)|\} \leq M_0$$

for any solution  $y$  to

$$\begin{cases} \frac{1}{p}(py')' + \tau y = \lambda f(t, y, py') \text{ a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

for each  $\lambda \in (0, 1)$ . Then (1.6) has at least one solution  $u \in C[0, 1] \cap C^1(0, 1)$  with  $pu' \in AC[0, 1]$ .

Next we gather together some results on the Sturm Liouville eigenvalue problem

$$(1.8) \quad \begin{cases} Lu = \lambda u & \text{a.e. on } [0, 1] \\ u \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

where  $Lu = -\frac{1}{pq(t)}[(pu')' + r(t)pu]$ . Assume (1.3) and

$$(1.9) \quad r, q \in L^1_p[0, 1] \text{ with } q > 0 \text{ a.e. on } [0, 1]$$

hold. Let

$$D(L) = \{w \in C[0, 1] : w, pw' \in AC[0, 1] \text{ with } w \in (\text{SL}) \text{ or } (\text{P})\}.$$

Then  $L$  has a countably infinite number ([1], [12], [16]) of real eigenvalues  $\lambda_i$  with corresponding eigenfunctions  $\psi_i \in D(L)$ . The eigenfunctions  $\psi_i$  may be chosen so that they form an orthonormal set and we may also arrange the eigenvalues so that

$$(1.10) \quad \lambda_0 < \lambda_1 < \lambda_2 < \dots$$

*Remark.* The  $\lambda_i$ 's may be estimated numerically ([2]) using SLEIGN.

In addition the set of eigenfunctions  $\psi_i$  form a basis for  $L^2_{pq}[0, 1]$  and if  $h \in L^2_{pq}[0, 1]$  then  $h$  has a Fourier series representation and  $h$  satisfies Parseval's equality i.e.

$$h = \sum_{i=0}^{\infty} \langle h, \psi_i \rangle \psi_i \quad \text{and} \quad \int_0^1 pq|h|^2 dt = \sum_{i=0}^{\infty} |\langle h, \psi_i \rangle|^2.$$

We are concerned with existence results for the nonlinear second order equation

$$(1.11) \quad \begin{cases} \frac{1}{p}(py')' + ry + \lambda_m qy = f(t, y, py') & \text{a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

where  $\lambda_m$  is the  $(m+1)^{st}$  eigenvalue of (1.8). In recent years several authors ([4], [7]–[9], [11], [13], [18]–[19]) have examined the boundary value problems

$$\begin{cases} y'' + n^2\pi^2y = f(t, y) & \text{a.e. on } [0, 1] \\ y(0) = y(1) = 0 \end{cases}$$

and

$$\begin{cases} y'' + m^2\pi^2y = f(t, y) & \text{a.e. on } [0, 1] \\ y(0) = y(1), y'(0) = y'(1) \end{cases}$$

where  $n \geq 1, m \geq 0$  are integers. Most of the papers in the literature ([3], [7], [11], [18]–[19]) concentrate on the first eigenvalue ( $n = 1$  or  $m = 0$ ). However over the last ten years or so ([6], [10]) the case when  $n > 1$  or  $m > 0$  has been discussed. This paper continues this study for the more general problem (1.11); also it provides a new approach to studying the above resonant type problems. We refer the reader to [6]–[9] for many of the motivating ideas in this paper. Finally it is of interest to note that in previous studies ([6], [8], [11]) the nonlinearity  $f$  is required to grow no more than linearly in  $y$  as  $|y| \rightarrow \infty$  whereas in this paper solutions will exist provided  $f$  grows fast enough e.g.  $yf(t, y, z) \geq A|y|^{\theta+1}$  for some  $A > 0$  and  $\theta > 0$ .

**2. Existence**

Existence theory is developed for the second order boundary value problem

$$(2.1) \quad \begin{cases} \frac{1}{p}(py')' + ry + \lambda_m qy = f(t, y, py') & \text{a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

where  $\lambda_m$  is the  $(m + 1)^{st}$  eigenvalue of

$$(2.2) \quad \begin{cases} Lu = \lambda u & \text{a.e. on } [0, 1] \\ u \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

and  $Lu = -\frac{1}{pq(t)}[(pu')' + r(t)pu]$ .

Two types of existence results are presented, the first examines the problem on the “left” of the eigenvalue whereas the second discusses the problem on the “right” of the eigenvalue.

**Existence theory I.**

Throughout this subsection let

$$H_{\alpha_0, \theta}(u_1) = \begin{cases} |u_1|^{\theta+1}, & |u_1| \leq 1 \\ |u_1|^{\alpha_0+1}, & |u_1| > 1. \end{cases}$$

**Theorem 2.1.** *Let  $pf : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  be an  $L^1$ -Carathéodory function with (1.3) and (1.9) satisfied. Suppose  $f$  has the decomposition  $f(t, u_1, u_2) = g(t, u_1, u_2) + h(t, u_1, u_2)$  with  $pg, ph : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$   $L^1$ -Carathéodory functions and*

$$(2.3) \quad \begin{cases} \text{there exist constants } A > 0, 0 < \alpha_0 < 1 \text{ and a function} \\ \phi \in L^1_p[0, 1], \phi > 0 \text{ a.e. on } [0, 1] \text{ with } u_1 g(t, u_1, u_2) \geq A\phi(t)H_{\alpha_0, \theta}(u_1) \\ \text{for a.e. } t \in [0, 1]; \text{ here } \alpha_0 \leq \theta \end{cases}$$

$$(2.4) \quad \left\{ \begin{array}{l} \text{there exist } \phi_i \in L_p^1[0, 1], i = 1, 2, 3 \text{ and constants } \beta_0 \text{ and} \\ \sigma \text{ with } |h(t, u_1, u_2)| \leq \phi_1(t) + \phi_2(t)|u_1|^{\beta_0} + \phi_3(t)|u_2|^\sigma \text{ for} \\ \text{a.e. } t \in [0, 1]; \text{ here } \beta_0 < \alpha_0 \text{ and } 0 \leq \sigma < \frac{\alpha_0}{2} \text{ and} \\ \phi_3 > 0 \text{ a.e. on } [0, 1] \text{ or } \phi_3 \equiv 0 \text{ on } [0, 1] \end{array} \right.$$

$$(2.5) \quad \left\{ \begin{array}{l} \text{there exist } \phi_i \in L_p^1[0, 1], i = 4, 5 \text{ and a constant } \gamma \leq \alpha_0 \text{ with} \\ |g(t, u_1, u_2)| \leq \phi_4(t) + \phi_5(t)|u_1|^\gamma \text{ for a.e. } t \in [0, 1] \end{array} \right.$$

$$(2.6) \quad \left\{ \begin{array}{l} \phi_4^2 q^{-1} \in L_p^1[0, 1], \phi_1^2 q^{-1} \in L_p^1[0, 1], \\ \left( \phi_5^{2(\alpha_0+1)} q^{-(\alpha_0+1)} \phi^{-2\gamma} \right)^{\frac{1}{\alpha_0+1-2\gamma}} \in L_p^1[0, 1], \\ \left( \phi_2^{2(\alpha_0+1)} q^{-(\alpha_0+1)} \phi^{-2\beta_0} \right)^{\frac{1}{\alpha_0+1-2\beta_0}} \in L_p^1[0, 1] \text{ and} \\ \left( \phi^2 q^{-(\alpha_0+1)} \right)^{\frac{1}{1-\alpha_0}} \in L_p^1[0, 1] \end{array} \right.$$

and

$$(2.7) \quad \left\{ \begin{array}{l} \left( \phi_1^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} \in L_p^1[0, 1], \left( \phi_2^{\alpha_0+1} \phi^{-(\beta_0+1)} \right)^{\frac{1}{\alpha_0-\beta_0}} \in L_p^1[0, 1], \\ \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} \in L_p^1[0, 1], \\ \left( \phi_5^{\alpha_0+1} \phi^{-\gamma} \right)^{\frac{1}{\alpha_0+1-\gamma}} \in L_p^1[0, 1] \text{ and} \\ \left( q^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} \in L_p^1[0, 1] \end{array} \right.$$

holding. Then (2.1) has at least one solution  $y \in C[0, 1] \cap C^1(0, 1)$  with  $py' \in AC[0, 1]$ .

*Remark.* Typical examples where (2.3) is satisfied are say (i)  $g(t, u_1, u_2) = u_1^{\frac{m}{n}}$ ,  $m$  odd and  $n$  odd or (ii)  $g(t, u_1, u_2) = u_1^{\frac{1}{2}}$ ,  $u_1 \geq 0$  with  $g(t, u_1, u_2) = -|u_1|^{\frac{1}{2}}$ ,  $u_1 < 0$ .

PROOF: Consider the family of problems

$$(2.8)_\lambda \quad \left\{ \begin{array}{l} \frac{1}{p}(py')' + ry + \mu qy = \lambda[f(t, y, py') + (\mu - \lambda_m)qy] \text{ a.e. on } [0, 1] \\ y \in (SL) \text{ or } (P) \end{array} \right.$$

where  $0 < \lambda < 1$  and  $\lambda_{m-1} < \mu < \lambda_m$ ; here  $\lambda_{-1} = -\infty$  (for notational purposes) with  $\lambda_i$  as described in (1.10).

Notice  $L_{pq}^2[0, 1] = \Omega \oplus \Omega^\perp$  where  $\Omega = span\{\psi_0, \psi_1, \dots, \psi_{m-1}\}$ ; here  $\psi_i$  are the eigenfunctions corresponding to the eigenvalues  $\lambda_i$  (see Section 1).

Let  $y$  be any solution to  $(2.8)_\lambda$ . Then  $y = u + w$  where  $u \in \Omega$  and  $w \in \Omega^\perp$ . Multiply  $(2.8)_\lambda$  by  $w - u$  and integrate from 0 to 1 to obtain

$$\begin{aligned} \int_0^1 (w - u)(py')' dt + \int_0^1 pr[w^2 - u^2] dt + \mu \int_0^1 pq[w^2 - u^2] dt \\ = \lambda \int_0^1 (w - u)pf(t, y, py') dt + \lambda(\mu - \lambda_m) \int_0^1 pq[w^2 - u^2] dt. \end{aligned}$$

Integration by parts yields

$$\int_0^1 (w - u)(py')' dt = Q_0 - \int_0^1 p(w')^2 dt + \int_0^1 p(u')^2 dt$$

where

$$Q_0 = \begin{cases} -\frac{a}{b}[w^2(1) - u^2(1)] - \frac{a}{\beta}[w^2(0) - u^2(0)] & \text{if } y \in (SL) \\ 0 & \text{if } y \in (P); \end{cases}$$

here  $y(0) = 0$  means  $u(0) + w(0) = 0$  and so  $u(0) = w(0) = 0$ . Thus we have

$$\begin{aligned} (2.9) \quad Q_0 + \int_0^1 [-p(w')^2 + prw^2 + \mu pqw^2] dt + \int_0^1 [p(u')^2 - pru^2 - \mu pq u^2] dt \\ = \lambda \int_0^1 (w - u)pf(t, y, py') dt + \lambda(\mu - \lambda_m) \int_0^1 pqw^2 dt \\ - \lambda(\mu - \lambda_m) \int_0^1 pq u^2 dt. \end{aligned}$$

Now since  $u \in \Omega$ ,  $w \in \Omega^\perp$  and  $y = u + w$  we have

$$u = \sum_{i=0}^{m-1} c_i \psi_i \quad \text{and} \quad w = \sum_{i=m}^{\infty} c_i \psi_i \quad \text{where} \quad c_i = \langle y, \psi_i \rangle;$$

note  $u = 0$  if  $m = 0$ . Then since  $(p\psi'_i)' + r p\psi_i + \lambda_i p q\psi_i = 0$  we have

$$\begin{aligned} Q_0 + \int_0^1 [-p(w')^2 + prw^2 + \mu pqw^2] dt + \int_0^1 [p(u')^2 - pru^2 - \mu pq u^2] dt \\ = \sum_{i=m}^{\infty} (\mu - \lambda_i) c_i^2 \int_0^1 pq\psi_i^2 dt + \sum_{i=0}^{m-1} (\lambda_i - \mu) c_i^2 \int_0^1 pq\psi_i^2 dt \\ \leq (\mu - \lambda_m) \int_0^1 pqw^2 dt + (\lambda_{m-1} - \mu) \int_0^1 pq u^2 dt. \end{aligned}$$



Put this into (2.9) to obtain

$$\begin{aligned} & \lambda \int_0^1 (w - u)pg(t, y, py') dt + (1 - \lambda)(\lambda_m - \mu) \int_0^1 pqw^2 dt \\ & + (\mu - \lambda_{m-1}) \int_0^1 pqu^2 dt + \lambda(\lambda_m - \mu) \int_0^1 pqu^2 dt \\ & \leq -\lambda \int_0^1 (w - u)ph(t, y, py') dt. \end{aligned}$$

Consequently

$$(2.10) \quad \begin{aligned} & \int_0^1 pyg(t, y, py') dt + (\lambda_m - \mu) \int_0^1 pqu^2 dt \leq 2 \int_0^1 pug(t, y, py') dt \\ & + \int_0^1 p|y||h(t, y, py')| dt + 2 \int_0^1 p|u||h(t, y, py')| dt. \end{aligned}$$

Assumption (2.3) yields

$$\begin{aligned} \int_0^1 pyg(t, y, py') dt & \geq A \int_0^1 p\phi H_{\alpha_0, \theta}(y) dt \\ & = A \int_0^1 p\phi|y|^{\alpha_0+1} dt + A \int_{\{t:|y(t)| \leq 1\}} p\phi[|y|^{\theta+1} - |y|^{\alpha_0+1}] dt \\ & \geq A \int_0^1 p\phi|y|^{\alpha_0+1} dt - A \int_0^1 p\phi dt \end{aligned}$$

and put this into (2.10), and use (2.4) and (2.5), to obtain

$$(2.11) \quad \begin{aligned} & A \int_0^1 p\phi|y|^{\alpha_0+1} dt + (\lambda_m - \mu) \int_0^1 pqu^2 dt \leq A \int_0^1 p\phi dt + 2 \int_0^1 p\phi_4|u| dt \\ & + 2 \int_0^1 p\phi_5|u||y|^\gamma dt + \int_0^1 p\phi_1|y| dt \\ & + \int_0^1 p\phi_2|y|^{\beta_0+1} dt + \int_0^1 p\phi_3|y||py'|^\sigma dt \\ & + 2 \int_0^1 p\phi_1|u| dt + 2 \int_0^1 p\phi_2|u||y|^{\beta_0} dt \\ & + 2 \int_0^1 p\phi_3|u||py'|^\sigma dt. \end{aligned}$$

For the remainder of the proof we assume without loss of generality that  $\sigma > 0$  and  $\phi_3 \not\equiv 0$  on  $[0, 1]$ . Let  $\epsilon > 0$  be given. Hölder's inequality together with assumption (2.6) immediately yields the following inequalities:

$$2 \int_0^1 p\phi_4|u| dt \leq 2Q_1 \left( \int_0^1 pqu^2 dt \right)^{\frac{1}{2}} \leq \epsilon \int_0^1 pqu^2 dt + \frac{Q_1}{\epsilon};$$

$$\begin{aligned}
 2 \int_0^1 p\phi_1|u| dt &\leq \epsilon \int_0^1 pqu^2 dt + \frac{Q_2}{\epsilon}; \\
 2 \int_0^1 p\phi_5|u||y|^\gamma dt &\leq 2Q_3 \left( \int_0^1 pqu^2 dt \right)^{\frac{1}{2}} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\gamma}{\alpha_0+1}} \\
 &\leq \epsilon Q_3 \int_0^1 pqu^2 dt + \frac{Q_3}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\gamma}{\alpha_0+1}}; \\
 2 \int_0^1 p\phi_2|u||y|^{\beta_0} dt &\leq \epsilon Q_4 \int_0^1 pqu^2 dt + \frac{Q_4}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\beta_0}{\alpha_0+1}}; \\
 \int_0^1 p\phi_1|y| dt &\leq Q_5 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}}; \\
 \int_0^1 p\phi_2|y|^{\beta_0+1} dt &\leq Q_6 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0+1}{\alpha_0+1}}; \\
 \int_0^1 p\phi_3|y||py'|^\sigma dt &\leq \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} \\
 &\quad \times \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}}; \\
 2 \int_0^1 p\phi_3|u||py'|^\sigma dt &\leq 2Q_7 \left( \int_0^1 pqu^2 dt \right)^{\frac{1}{2}} \\
 &\quad \times \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}} \\
 &\leq \epsilon Q_7 \int_0^1 pqu^2 dt \\
 &\quad + \frac{Q_7}{\epsilon} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\alpha_0}{\alpha_0+1}}
 \end{aligned}$$

for some constants  $Q_1, \dots, Q_7$ . Put these into (2.11) to obtain

$$\begin{aligned}
 &A \int_0^1 p\phi|y|^{\alpha_0+1} dt + (\lambda_m - \mu - 2\epsilon - \epsilon Q_3 - \epsilon Q_4 - \epsilon Q_7) \int_0^1 pqu^2 dt \\
 &\leq Q_8 + \frac{Q_3}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\gamma}{\alpha_0+1}} + \frac{Q_4}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\beta_0}{\alpha_0+1}} \\
 &+ Q_5 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} + Q_6 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0+1}{\alpha_0+1}} \\
 &+ \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}}
 \end{aligned}$$

$$+ \frac{Q_7}{\epsilon} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\alpha_0}{\alpha_0+1}}$$

for some constant  $Q_8$ . We may choose  $\epsilon$  so that  $\lambda_m - \mu - 2\epsilon - \epsilon Q_3 - \epsilon Q_4 - \epsilon Q_7 > 0$  and we have

$$\begin{aligned}
 A \int_0^1 p\phi|y|^{\alpha_0+1} dt &\leq Q_8 + \frac{Q_3}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\gamma}{\alpha_0+1}} \\
 &+ \frac{Q_4}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\beta_0}{\alpha_0+1}} + Q_5 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} \\
 (2.12) \quad &+ Q_6 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0+1}{\alpha_0+1}} \\
 &+ \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}} \\
 &+ \frac{Q_7}{\epsilon} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\alpha_0}{\alpha_0+1}}.
 \end{aligned}$$

We now consider two cases  $\int_0^1 p\phi|y|^{\alpha_0+1} dt > 1$  and  $\int_0^1 p\phi|y|^{\alpha_0+1} dt \leq 1$  separately.

Case (i).  $\int_0^1 p\phi|y|^{\alpha_0+1} dt > 1$ .

Divide (2.12) by  $\left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}}$  and use  $\int_0^1 p\phi|y|^{\alpha_0+1} dt > 1$  to obtain

$$\begin{aligned}
 A \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\alpha_0}{\alpha_0+1}} &\leq Q_8 + \frac{Q_3}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\gamma-1}{\alpha_0+1}} \\
 &+ \frac{Q_4}{\epsilon} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{2\beta_0-1}{\alpha_0+1}} \\
 &+ Q_5 + Q_6 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0}{\alpha_0+1}} \\
 &+ \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}} \\
 &+ \frac{Q_7}{\epsilon} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\alpha_0}{\alpha_0+1}}.
 \end{aligned}$$

Now since  $\max\{2\gamma - 1, 2\beta_0 - 1, \beta_0\} < \alpha_0$  there exist constants  $Q_9, Q_{10}$  and  $Q_{11}$

with

$$\begin{aligned} \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{\alpha_0}{\alpha_0+1}} &\leq Q_9 + Q_{10} \left(\int_0^1 p\left(\phi_3^{\alpha_0+1}\phi^{-1}\right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt\right)^{\frac{\alpha_0}{\alpha_0+1}} \\ &\quad + Q_{11} \left(\int_0^1 p\left(\phi_3^{\alpha_0+1}\phi^{-1}\right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt\right)^{\frac{2\alpha_0}{\alpha_0+1}}. \end{aligned}$$

Using the inequality  $(a + b)^c \leq 2^c(a^c + b^c)$  for  $a \geq 0, b \geq 0, c \geq 0$  we see that there exist constants  $Q_{12}$  and  $Q_{13}$  with

$$(2.13) \quad \int_0^1 p\phi|y|^{\alpha_0+1} dt \leq Q_{12} + Q_{13} \left(\int_0^1 p\left(\phi_3^{\alpha_0+1}\phi^{-1}\right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt\right)^2.$$

Case (ii).  $\int_0^1 p\phi|y|^{\alpha_0+1} dt \leq 1$ .

In this case (2.13) is clearly true with  $Q_{12} = 1$ .

Consequently in all cases (2.13) is true. Returning to (2.8) $_\lambda$  we have

$$(2.14) \quad y(t) = \lambda \int_0^1 G(t, s)[f(s, y(s), p(s)y'(s)) + (\mu - \lambda_m)q(s)y(s)] ds$$

and

$$(2.15) \quad p(t)y'(t) = \lambda \int_0^1 p(t)G_t(t, s)[f(s, y(s), p(s)y'(s)) + (\mu - \lambda_m)q(s)y(s)] ds$$

where  $G(t, s)$  is the Green's function associated with  $\frac{1}{p}(pv')' + rv + \mu qv = 0$  a.e. on  $[0, 1]$ ,  $v \in (\text{SL})$  or  $(\text{P})$ .

Notice ([16], [17]) that  $\sup_{t \in [0, 1]} |p(t)G_t(t, s)| \leq Q_{14}p(s)$  for some constant  $Q_{14}$ . Now (2.15) together with (2.4) and (2.5) imply for  $t \in (0, 1)$  that

$$\begin{aligned} |p(t)y'(t)| &\leq Q_{15} \int_0^1 p\phi_1 ds + Q_{15} \int_0^1 p\phi_2|y|^{\beta_0} ds + Q_{15} \int_0^1 p\phi_3|py'|^\sigma ds \\ &\quad + Q_{15} \int_0^1 p\phi_4 ds + Q_{15} \int_0^1 p\phi_5|y|^\gamma ds + Q_{16} \int_0^1 pq|y| ds \end{aligned}$$

for some constants  $Q_{15}$  and  $Q_{16}$ . Hölder's inequality together with (2.6) implies

$$\begin{aligned} |p(t)y'(t)| &\leq Q_{17} + Q_{18} \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{\beta_0}{\alpha_0+1}} \\ &\quad + Q_{19} \left(\int_0^1 p\left(\phi_3^{\alpha_0+1}\phi^{-1}\right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt\right)^{\frac{\alpha_0}{\alpha_0+1}} \\ &\quad + Q_{20} \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{\gamma}{\alpha_0+1}} + Q_{21} \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{1}{\alpha_0+1}} \end{aligned}$$

for some constants  $Q_{17}, \dots, Q_{21}$ . Thus for  $t \in (0, 1)$  we have

$$\begin{aligned}
 |p(t)y'(t)|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} &\leq Q_{22} + Q_{23} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\sigma\beta_0}{\alpha_0}} \\
 &+ Q_{24} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^\sigma \\
 &+ Q_{25} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\sigma\gamma}{\alpha_0}} \\
 &+ Q_{26} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\sigma}{\alpha_0}}
 \end{aligned}
 \tag{2.16}$$

for some constants  $Q_{22}, \dots, Q_{26}$ . This together with (2.13) implies

$$\begin{aligned}
 &\int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \\
 &\leq Q_{27} + Q_{28} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\sigma\beta_0}{\alpha_0}} \\
 &+ Q_{29} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^\sigma \\
 &+ Q_{30} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\sigma\gamma}{\alpha_0}} \\
 &+ Q_{31} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{2\sigma}{\alpha_0}}
 \end{aligned}$$

for some constants  $Q_{27}, \dots, Q_{31}$ . Finally since  $\max\{\frac{2\sigma\beta_0}{\alpha_0}, \sigma, \frac{2\sigma\gamma}{\alpha_0}, \frac{2\sigma}{\alpha_0}\} < 1$  there exists a constant  $Q_{32}$  with

$$\int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \leq Q_{32}
 \tag{2.17}$$

and this together with (2.13) implies that there exists a constant  $Q_{33}$  with

$$\int_0^1 p\phi|y|^{\alpha_0+1} dt \leq Q_{33}.
 \tag{2.18}$$

Putting these inequalities into (2.16) establishes the existence of a constant  $Q_{34}$  with

$$\sup_{t \in (0,1)} |p(t)y'(t)| \leq Q_{34}.
 \tag{2.19}$$

Now (2.14) together ([16], [17]) with  $\sup_{t \in [0,1]} |G(t, s)| \leq Q_{35}p(s)$ , for some constant  $Q_{35}$ , and Hölder's inequality implies for  $t \in [0, 1]$  that

$$\begin{aligned}
 |y(t)| \leq & Q_{36} + Q_{37} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0}{\alpha_0+1}} \\
 & + Q_{38} \left( \int_0^1 p \left( \phi_3^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} |py'|^{\frac{\sigma(\alpha_0+1)}{\alpha_0}} dt \right)^{\frac{\alpha_0}{\alpha_0+1}} \\
 & + Q_{39} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\gamma}{\alpha_0+1}} + Q_{40} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}}
 \end{aligned}$$

for some constants  $Q_{36}, \dots, Q_{40}$ . This together with (2.17) and (2.18) implies that there is a constant  $Q_{41}$  with

$$(2.20) \quad \sup_{t \in [0,1]} |y(t)| \leq Q_{41}.$$

Now (2.19), (2.20) together with Theorem 1.2 establish the result. □

**Example.** Theorem 2.1 (here  $H_{\alpha_0, \theta}(u) = H_{\frac{1}{3}, \frac{1}{3}}(u)$ ) immediately guarantees that

$$\begin{cases}
 y'' + n^2\pi^2 y = y^{\frac{1}{3}} + [y']^{\frac{1}{7}} + 1 \quad \text{a.e. on } [0, 1] \\
 y(0) = y(1) = 0, \quad n \in \{1, 2, \dots\}
 \end{cases}$$

has a solution.

One can improve considerably the above theorem if  $m = 0$  (at the first eigenvalue). In particular the condition  $0 < \alpha_0 < 1$  is replaced by  $\alpha_0 > 0$  in this case; also condition (2.5) can be improved and the condition  $\sigma < \frac{\alpha_0}{2}$  can be relaxed. We present two existence results.

Consider

$$(2.21) \quad \begin{cases} \frac{1}{p}(py')' + ry + \lambda_0 qy = f(t, y, py') \quad \text{a.e. on } [0, 1] \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

where  $\lambda_0$  is the first eigenvalue of (2.2).

**Theorem 2.2.** *Let  $pf : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  be an  $L^1$ -Carathéodory function with (1.3) and (1.9) satisfied. Suppose  $f$  has the decomposition  $f(t, u_1, u_2) = g(t, u_1, u_2) + h(t, u_1, u_2)$  with  $pg, ph : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$   $L^1$ -Carathéodory functions and*

$$(2.22) \quad \begin{cases} \text{there exist constants } A > 0, \alpha_0 > 0 \text{ and a function } \phi \in L^1_p[0, 1], \\ \phi > 0 \text{ a.e. on } [0, 1] \text{ with } u_1g(t, u_1, u_2) \geq A\phi(t)H_{\alpha_0, \theta}(u_1) \\ \text{for a.e. } t \in [0, 1]; \text{ here } \alpha_0 \leq \theta \end{cases}$$

$$(2.23) \quad \begin{cases} \text{there exist } \phi_i \in L_p^1[0, 1], i = 1, 2, 3 \text{ and constants } \beta_0 \text{ and} \\ \sigma \text{ with } |h(t, u_1, u_2)| \leq \phi_1(t) + \phi_2(t)|u_1|^{\beta_0} + \phi_3(t)|u_2|^\sigma \text{ for} \\ \text{a.e. } t \in [0, 1]; \text{ here } \beta_0 < \alpha_0 \text{ and } \phi_3 > 0 \\ \text{a.e. on } [0, 1] \text{ or } \phi_3 \equiv 0 \text{ on } [0, 1] \end{cases}$$

$$(2.24) \quad \begin{cases} \text{there exist } \phi_i \in L_p^1[0, 1], i = 4, 5, 6 \text{ and constants } \gamma \leq \alpha_0, \tau > \sigma \\ \text{with } |g(t, u_1, u_2)| \leq \phi_4(t) + \phi_5(t)|u_1|^\gamma + \phi_6(t)|u_2|^\tau \\ \text{for a.e. } t \in [0, 1]; \\ \text{here } \phi_6 > 0 \text{ a.e. on } [0, 1] \text{ or } \phi_6 \equiv 0 \text{ on } [0, 1] \end{cases}$$

$$(2.25) \quad \sigma < \min\{1, \frac{\alpha_0}{\gamma}, \alpha_0\} \text{ and } \tau < 1$$

$$(2.26) \quad \begin{cases} \left( \phi_1^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} \in L_p^1[0, 1], \left( \phi_2^{\alpha_0+1} \phi^{-(\beta_0+1)} \right)^{\frac{1}{\alpha_0-\beta_0}} \in L_p^1[0, 1], \\ \left( \phi_5^{\alpha_0+1} \phi^{-\gamma} \right)^{\frac{1}{\alpha_0+1-\gamma}} \in L_p^1[0, 1] \text{ and } \left( q^{\alpha_0+1} \phi^{-1} \right)^{\frac{1}{\alpha_0}} \in L_p^1[0, 1] \end{cases}$$

and

$$(2.27) \quad \begin{cases} \text{with } \kappa = \max\{\frac{\alpha_0+1}{\alpha_0}, 2\}, \left( \phi_3 \phi^{-\frac{1}{\alpha_0+1}} \right)^\kappa \in L_p^1[0, 1]. \text{ Also need} \\ \phi_6^\kappa \in L_p^1[0, 1] \text{ and } \left( \phi_3 \phi^{-\frac{1}{\alpha_0+1}} \right)^{\frac{\kappa\tau}{\tau-\sigma}} (\phi_6)^{-\kappa \left( \frac{\sigma}{\tau-\sigma} \right)} \in L_p^1[0, 1] \\ \text{if } \phi_6 > 0 \text{ a.e. on } [0, 1] \end{cases}$$

holding. Then (2.21) has at least one solution  $y \in C[0, 1] \cap C^1(0, 1)$  with  $py' \in AC[0, 1]$ .

PROOF: Let  $y$  be a solution of (2.8) $_\lambda$  with  $m = 0$ . Following the ideas of Theorem 2.1 with  $u = 0$  and  $y = w$  we obtain the analogue of (2.11), namely

$$(2.28) \quad A \int_0^1 p\phi|y|^{\alpha_0+1} dt \leq A \int_0^1 p\phi dt + \int_0^1 p\phi_1|y| dt + \int_0^1 p\phi_2|y|^{\beta_0+1} dt + \int_0^1 p\phi_3|y||py'|^\sigma dt.$$

Hölder's inequality implies

$$(2.29) \quad A \int_0^1 p\phi|y|^{\alpha_0+1} dt \leq N_0 + N_1 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} + N_2 \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0+1}{\alpha_0+1}} + \int_0^1 p\phi_3|y||py'|^\sigma dt$$

for some constants  $N_0, N_1$  and  $N_2$ . Let  $\kappa = \max\{2, \frac{\alpha_0+1}{\alpha_0}\}$ . Hölder's inequality together with assumption (2.27) implies

$$\int_0^1 p\phi_3|y||py'|^\sigma dt \leq \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{1}{\alpha_0+1}} \left(\int_0^1 p\left(\phi_3\phi^{-\frac{1}{\alpha_0+1}}\right)^\kappa |py'|^{\sigma\kappa} dt\right)^{\frac{1}{\kappa}}$$

if  $\kappa = \frac{\alpha_0+1}{\alpha_0}$  whereas

$$\int_0^1 p\phi_3|y||py'|^\sigma dt \leq \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{1}{\alpha_0+1}} \times \left(\int_0^1 p\left(\phi_3\phi^{-\frac{1}{\alpha_0+1}}\right)^\kappa |py'|^{\sigma\kappa} dt\right)^{\frac{1}{\kappa}} \left(\int_0^1 p(t) dt\right)^{\frac{\alpha_0-1}{2(\alpha_0+1)}}$$

if  $\kappa = 2$ . Put this into (2.29) and essentially the same reasoning as in Theorem 2.1 establishes the existence of constants  $N_3$  and  $N_4$  with

$$(2.30) \quad \int_0^1 p\phi|y|^{\alpha_0+1} dt \leq N_3 + N_4 \left(\int_0^1 p\left(\phi_3\phi^{-\frac{1}{\alpha_0+1}}\right)^\kappa |py'|^{\sigma\kappa} dt\right)^{\frac{\alpha_0+1}{\kappa\alpha_0}}.$$

Also (2.15) implies (as in Theorem 2.1) for  $t \in (0, 1)$  that

$$(2.31) \quad \begin{aligned} |p(t)y'(t)| \leq & N_5 + N_6 \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{\beta_0}{\alpha_0+1}} + N_7 \int_0^1 p\phi_3|py'|^\sigma dt \\ & + N_8 \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{\gamma}{\alpha_0+1}} + N_9 \int_0^1 p\phi_6|py'|^\tau dt \\ & + N_{10} \left(\int_0^1 p\phi|y|^{\alpha_0+1} dt\right)^{\frac{1}{\alpha_0+1}} \end{aligned}$$

for some constants  $N_5, \dots, N_{10}$ . Again with  $\kappa = \max\{2, \frac{\alpha_0+1}{\alpha_0}\}$  we have

$$\begin{aligned} \int_0^1 p\phi_3|py'|^\sigma dt & \leq \left(\int_0^1 p\left(\phi_3\phi^{-\frac{1}{\alpha_0+1}}\right)^\kappa |py'|^{\sigma\kappa} dt\right)^{\frac{1}{\kappa}} \left(\int_0^1 p\phi dt\right)^{\frac{1}{\alpha_0+1}} \\ & \text{if } \kappa = \frac{\alpha_0+1}{\alpha_0} \\ \int_0^1 p\phi_3|py'|^\sigma dt & \leq \left(\int_0^1 p\left(\phi_3\phi^{-\frac{1}{\alpha_0+1}}\right)^\kappa |py'|^{\sigma\kappa} dt\right)^{\frac{1}{\kappa}} \left(\int_0^1 p\phi^{\frac{2}{\alpha_0+1}} dt\right)^{\frac{1}{2}} \\ & \text{if } \kappa = 2 \\ \int_0^1 p\phi_6|py'|^\tau dt & \leq \left(\int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt\right)^{\frac{1}{\kappa}} \left(\int_0^1 p(t) dt\right)^{1-\frac{1}{\kappa}}. \end{aligned}$$



There are two cases to consider, namely  $\phi_6 > 0$  a.e. on  $[0, 1]$  or  $\phi_6 \equiv 0$  on  $[0, 1]$ .

Case (i).  $\phi_6 > 0$  a.e. on  $[0, 1]$ .

Putting the above into (2.31) and using (2.30) leads to

$$\begin{aligned}
 \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt &\leq N_{11} + N_{12} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\beta_0\tau}{\alpha_0}} \\
 &+ N_{13} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^\tau \\
 (2.32) \quad &+ N_{14} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\gamma\tau}{\alpha_0}} \\
 &+ N_{15} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^\tau \\
 &+ N_{16} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\tau}{\alpha_0}}
 \end{aligned}$$

for some constants  $N_{11}, \dots, N_{16}$ . Also Hölder's inequality implies

$$\begin{aligned}
 &\int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \\
 &\leq \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^{\frac{\sigma}{\tau}} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^{\frac{\kappa\tau}{\tau-\sigma}} (\phi_6)^{-\kappa\left(\frac{\sigma}{\tau-\sigma}\right)} dt \right)^{\frac{\tau-\sigma}{\tau}}
 \end{aligned}$$

and putting this into (2.32) yields

$$\begin{aligned}
 \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt &\leq N_{17} + N_{18} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^{\frac{\beta_0\sigma}{\alpha_0}} \\
 &+ N_{19} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^\sigma + N_{20} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^{\frac{\gamma\sigma}{\alpha_0}} \\
 &+ N_{21} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^\tau + N_{22} \left( \int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \right)^{\frac{\sigma}{\alpha_0}}
 \end{aligned}$$

for some constants  $N_{17}, \dots, N_{22}$ . Now since  $\max\{\frac{\sigma\beta_0}{\alpha_0}, \sigma, \frac{\sigma\gamma}{\alpha_0}, \tau, \frac{\sigma}{\alpha_0}\} < 1$  then there exists a constant  $N_{23}$  with

$$\int_0^1 p\phi_6^\kappa |py'|^{\tau\kappa} dt \leq N_{23}$$

and essentially the same reasoning as in Theorem 2.1 establishes the result.

Case (ii).  $\phi_6 \equiv 0$  on  $[0, 1]$ .

We may assume without loss of generality that  $\sigma > 0$  and  $\phi_3 > 0$  a.e. on  $[0, 1]$ ; otherwise the result is easy. Then (2.31) for  $t \in (0, 1)$  becomes

$$\begin{aligned} |p(t)y'(t)| &\leq N_{23} + N_{24} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\beta_0}{\alpha_0+1}} \\ &\quad + N_{25} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{1}{\kappa}} \\ &\quad + N_{26} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{\gamma}{\alpha_0+1}} + N_{27} \left( \int_0^1 p\phi|y|^{\alpha_0+1} dt \right)^{\frac{1}{\alpha_0+1}} \end{aligned}$$

for some constants  $N_{23}, \dots, N_{27}$ . This together with (2.30) leads to

$$\begin{aligned} &\int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \\ &\leq N_{28} + N_{29} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\beta_0\sigma}{\alpha_0}} \\ &\quad + N_{30} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^\sigma \\ &\quad + N_{31} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\gamma\sigma}{\alpha_0}} \\ &\quad + N_{32} \left( \int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \right)^{\frac{\sigma}{\alpha_0}} \end{aligned}$$

for some constants  $N_{28}, \dots, N_{32}$ . Thus there exists a constant  $N_{33}$  with

$$\int_0^1 p \left( \phi_3\phi^- - \frac{1}{\alpha_0+1} \right)^\kappa |py'|^{\sigma\kappa} dt \leq N_{33}$$

and the result follows as in Theorem 2.1. □

The next theorem establishes the existence of a nonnegative solution to

$$(2.33) \quad \begin{cases} \frac{1}{p}(py')' + \lambda_0 qy = \psi(t)f(t, y, py'), & 0 < t < 1 \\ y \in (\text{SL}) \text{ or } (\text{P}) \end{cases}$$

where  $\lambda_0$  is the first eigenvalue of (2.2) with  $r \equiv 0$  and  $q, \psi$  satisfies

$$(2.34) \quad q, \psi \in L^1_p[0, 1] \text{ with } q, \psi > 0 \text{ on } (0, 1).$$

Let

$$H^*_{\alpha_0, \theta}(u_1) = \begin{cases} u_1^{\theta+1}, & 0 \leq u_1 \leq 1 \\ u_1^{\alpha_0+1}, & 1 < u_1 < \infty. \end{cases}$$

**Theorem 2.3.** Let  $f : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  be continuous with (1.3), (2.34) and

$$(2.35) \quad f(t, 0, 0) \leq 0$$

holding. Suppose  $\psi f$  has the decomposition  $\psi(t)f(t, u_1, u_2) = g(t, u_1, u_2) + h(t, u_1, u_2)$  with  $pg, ph : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$   $L^1$ -Carathéodory functions and

$$(2.36) \quad \left\{ \begin{array}{l} \text{there exist constants } A > 0, \alpha_0 > 0 \text{ and a function } \phi \in L^1_p[0, 1], \\ \phi > 0 \text{ on } (0, 1) \text{ with } u_1g(t, u_1, u_2) \geq A\phi(t)H^*_{\alpha_0, \theta}(u_1) \\ \text{for } t \in (0, 1), u_1 \geq 0 \text{ and } u_2 \in \mathbf{R}; \text{ here } \alpha_0 \leq \theta \end{array} \right.$$

$$(2.37) \quad \left\{ \begin{array}{l} \text{there exist } \phi_i \in L^1_p[0, 1], i = 1, 2, 3 \text{ and constants } \beta_0 \text{ and} \\ \sigma \text{ with } |h(t, u_1, u_2)| \leq \phi_1(t) + \phi_2(t)u_1^{\beta_0} + \phi_3(t)|u_2|^\sigma \text{ for} \\ t \in (0, 1), u_1 \geq 0 \text{ and } u_2 \in \mathbf{R}; \text{ here } \beta_0 < \alpha_0 \\ \text{and } \phi_3 > 0 \text{ on } (0, 1) \text{ or } \phi_3 \equiv 0 \end{array} \right.$$

and

$$(2.38) \quad \left\{ \begin{array}{l} \text{there exist } \phi_i \in L^1_p[0, 1], i = 4, 5, 6 \text{ and constants } \gamma \leq \alpha_0, \tau > \sigma \\ \text{with } |g(t, u_1, u_2)| \leq \phi_4(t) + \phi_5(t)u_1^\gamma + \phi_6(t)|u_2|^\tau \text{ for } t \in (0, 1), u_1 \geq 0 \\ \text{and } u_2 \in \mathbf{R} \text{ and } \phi_6 > 0 \text{ on } (0, 1) \text{ or } \phi_6 \equiv 0 \end{array} \right.$$

hold. Finally suppose (2.25) and (2.26) are satisfied. Then (2.33) has at least one nonnegative solution  $y \in C[0, 1] \cap C^1(0, 1)$  with  $py' \in AC[0, 1]$ .

PROOF: Consider the family of problems

$$(2.39)_\lambda \quad \left\{ \begin{array}{l} \frac{1}{p}(py')' + \mu qy = \lambda f^*(t, y, py'), \quad 0 < t < 1 \\ y \in (SL) \text{ or } (P) \end{array} \right.$$

where  $0 < \lambda < 1$  and

$$\mu = \begin{cases} 0 & \text{if } y \in (SL) \text{ and } \alpha^2 + a^2 > 0 \\ -1 & \text{if } y \in (P) \text{ or } y \in (SL) \text{ with } \alpha = a = 0. \end{cases}$$

Also

$$f^*(t, u_1, u_2) = \begin{cases} \psi(t)f(t, u_1, u_2) + (\mu - \lambda_0)qu_1, & u_1 \geq 0 \\ \psi(t)f(t, 0, u_2) + (\mu + 1)qu_1, & u_1 < 0. \end{cases}$$

Notice  $pf^* : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  is an  $L^1$ -Carathéodory function.

Let  $y$  be a solution to (2.39) $_\lambda$  for some  $0 < \lambda < 1$ . We **claim** that  $y \geq 0$  on  $[0, 1]$ . If not then  $y$  would have a negative absolute minimum somewhere on  $[0, 1]$ ,

say at  $t_0$ . If  $t_0 \in (0, 1)$  then  $y'(t_0) = 0$  and this together with the differential equation and (2.35) yields

$$y''(t_0) = \frac{1}{p(t_0)}(p(t_0)y'(t_0))' \\ = \lambda(\psi(t_0)f(t_0, 0, 0) + q(t_0)y(t_0)) + (\lambda - 1)\mu q(t_0)y(t_0) < 0,$$

a contradiction. Next suppose the negative absolute minimum were to occur at  $t_0 = 0$ . Consider first the Sturm Liouville boundary condition. Of course we need only consider  $\beta \neq 0$ . If  $\alpha \neq 0$  as well then

$$y(0) \lim_{t \rightarrow 0^+} p(t)y'(t) = \frac{\alpha}{\beta}y^2(0) > 0,$$

which implies  $y^2(t)$  is an increasing function near 0, a contradiction. So it remains to consider the case  $\alpha = 0$  and  $\beta \neq 0$ . The boundary condition is  $\lim_{t \rightarrow 0^+} p(t)y'(t) = 0$ . Now  $f(0, 0, 0) \leq 0$  and this together with the differential equation and (2.34) implies there exists  $\delta > 0$  with  $(p(t)y'(t))' < 0$  for  $t \in (0, \delta)$ . Thus the boundary condition implies  $p(t)y'(t) < 0$  for  $t \in (0, \delta)$ , a contradiction. Consequently  $t_0 \neq 0$ . A similar argument shows  $t_0 \neq 1$ . Thus our claim is established for Sturm Liouville boundary data.

Consider now Periodic boundary data. If the absolute minimum of  $y$  occurs at  $t_0 = 0$  then, since  $y(0) = y(1)$ , it must also occur at 1. Thus  $\lim_{t \rightarrow 0^+} p(t)y'(t) \geq 0$  and  $\lim_{t \rightarrow 1^-} p(t)y'(t) \leq 0$ . Consequently

$$\lim_{t \rightarrow 0^+} p(t)y'(t) = \lim_{t \rightarrow 1^-} p(t)y'(t) = 0$$

because of the second boundary condition. As above there exists  $\delta > 0$  with  $(p(t)y'(t))' < 0$  for  $t \in (0, \delta)$  and so  $p(t)y'(t) < 0$  for  $t \in (0, \delta)$ , a contradiction.

Thus  $y \geq 0$  on  $[0, 1]$  for any solution  $y$  to (2.39) $_{\lambda}$ . Consequently  $y$  satisfies

$$\frac{1}{p}(py')' + \mu qy = \lambda(\psi(t)f(t, y, py') + (\mu - \lambda_0)qy), \quad 0 < t < 1.$$

Essentially the same reasoning as in Theorem 2.2 (in this case we look at  $\int_0^1 p\phi y^{\alpha_0+1} dt$ ) guarantees the existence of a solution  $y$  to (2.39) $_1$ . Of course  $y$  is automatically a solution of (2.33) since  $y \geq 0$  on  $[0, 1]$ . □

**Existence theory II.**

In this subsection we examine the resonant problem (2.1) on the “right” of the eigenvalue.

**Theorem 2.4.** *Let  $pf : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$  be an  $L^1$ -Carathéodory function with (1.3) and (1.9) holding. Suppose  $f$  has the decomposition  $f(t, u_1, u_2) =$*

$g(t, u_1, u_2) + h(t, u_1, u_2)$  with  $pg, ph : [0, 1] \times \mathbf{R}^2 \rightarrow \mathbf{R}$   $L^1$ -Carathéodory functions and assume

$$(2.40) \quad \begin{cases} \text{there exist constants } A > 0, 0 < \alpha_0 < 1 \text{ and a function} \\ \phi \in L^1_p[0, 1], \phi > 0 \text{ a.e. on } [0, 1] \text{ with} \\ u_1g(t, u_1, u_2) \leq -A\phi(t)H_{\alpha_0, \theta}(u_1) \text{ for a.e. } t \in [0, 1]; \text{ here } \alpha_0 \leq \theta \end{cases}$$

holds. In addition assume (2.4), (2.5), (2.6) and (2.7) are satisfied. Then (2.1) has at least one solution  $y \in C[0, 1] \cap C^1(0, 1)$  with  $py' \in AC[0, 1]$ .

PROOF: Consider the family of problems

$$(2.41)_\lambda \quad \begin{cases} \frac{1}{p}(py')' + ry + \mu qy = \lambda[f(t, y, py') + (\mu - \lambda_m)qy] \text{ a.e. on } [0, 1] \\ y \in (SL) \text{ or } (P) \end{cases}$$

where  $0 < \lambda < 1$  and  $\lambda_m < \mu < \lambda_{m+1}$ .

Notice  $L^2_{pq}[0, 1] = \Gamma \oplus \Gamma^\perp$  where  $\Gamma = span \{\psi_0, \psi_1, \dots, \psi_m\}$ . Multiply (2.41) $_\lambda$  by  $w - u$  and integrate from 0 to 1 to obtain as in Theorem 2.1 ( $Q_0$  is as in Theorem 2.1)

$$(2.42) \quad \begin{aligned} & Q_0 + \int_0^1 [-p(w')^2 + prw^2 + \mu pqw^2] dt + \int_0^1 [p(u')^2 - pr u^2 - \mu pq u^2] dt \\ &= \lambda \int_0^1 (w - u)pf(t, y, py') dt + \lambda(\mu - \lambda_m) \int_0^1 pqw^2 dt \\ &\quad - \lambda(\mu - \lambda_m) \int_0^1 pq u^2 dt. \end{aligned}$$

Now since  $u \in \Gamma, w \in \Gamma^\perp$  and  $y = u + w$  we have

$$u = \sum_{i=0}^m c_i \psi_i \text{ and } w = \sum_{i=m+1}^\infty c_i \psi_i \text{ where } c_i = \langle y, \psi_i \rangle.$$

Also as before

$$\begin{aligned} & Q_0 + \int_0^1 [-p(w')^2 + prw^2 + \mu pqw^2] dt + \int_0^1 [p(u')^2 - pr u^2 - \mu pq u^2] dt \\ &\leq (\mu - \lambda_{m+1}) \int_0^1 pqw^2 dt + (\lambda_m - \mu) \int_0^1 pq u^2 dt \end{aligned}$$

so putting this into (2.42) yields

$$\begin{aligned} & \lambda \int_0^1 (w - u)pg(t, y, py') dt + (1 - \lambda)(\mu - \lambda_m) \int_0^1 pq u^2 dt \\ &\quad + (\lambda_{m+1} - \mu) \int_0^1 pqw^2 dt + \lambda(\mu - \lambda_m) \int_0^1 pqw^2 dt \\ &\leq -\lambda \int_0^1 (w - u)ph(t, y, py') dt. \end{aligned}$$

Now  $w - u = -y + 2w$  and  $-yg(t, y, py') \geq A\phi(t)H_{\alpha_0, \theta}(y)$  for a.e.  $t \in [0, 1]$  so with the above we have

$$\begin{aligned} A \int_0^1 p\phi H_{\alpha_0, \theta}(y) dt + (\mu - \lambda_m) \int_0^1 pqw^2 dt &\leq -2 \int_0^1 pwg(t, y, py') dt \\ &+ \int_0^1 p|y||h(t, y, py')| dt \\ &+ 2 \int_0^1 p|w||h(t, y, py')| dt. \end{aligned}$$

Essentially the same reasoning as in Theorem 2.1 (the only difference is that we use  $\int_0^1 pqw^2 dt$  in place of  $\int_0^1 pqu^2 dt$ ) establishes the result.  $\square$

#### REFERENCES

- [1] Atkinson F.V., *Discrete and continuous boundary problems*, Academic Press, New York, 1964.
- [2] Atkinson F.V., Everitt W.N., Zettl A., *Regularization of a Sturm Liouville problem with an interior singularity*, *Diff. Int. Eq.* **1** (1988), 213–221.
- [3] Bobisud L.E., O'Regan D., *Positive solutions for a class of nonlinear singular boundary value problems at resonance*, *Jour. Math. Anal. Appl.* **184** (1994), 263–284.
- [4] Cesari L., Kannan R., *Existence of solutions of a nonlinear differential equation*, *Proc. Amer. Math. Soc.* **88** (1983), 605–613.
- [5] Fonda A., Mawhin J., *Quadratic forms, weighted eigenfunctions and boundary value problems for nonlinear second order ordinary differential equations*, *Proc. Royal Soc. Edinburgh* **112A** (1989), 145–153.
- [6] Iannacci R., Nkashama M.N., *Unbounded perturbations of forced second order ordinary differential equations at resonance*, *Jour. Diff. Eq.* **69** (1987), 289–309.
- [7] ———, *Nonlinear two point boundary value problems at resonance without Landesman-Lazer conditions*, *Proc. Amer. Math. Soc.* **106** (1989), 943–952.
- [8] Mawhin J., Ward J.R., *Periodic solutions of some forced Lienard differential equations at resonance*, *Archiv der Math.* **41** (1983), 337–351.
- [9] Mawhin J., Ward J.R., Willem M., *Necessary and sufficient conditions for the solvability of a nonlinear two point boundary value problem*, *Proc. Amer. Math. Soc.* **93** (1985), 667–674.
- [10] Mawhin J., Willem M., *Critical point theory and Hamiltonian systems*, Springer Verlag, New York, 1989.
- [11] Mawhin J., Omano W., *Two point boundary value problems for nonlinear perturbations of some singular linear differential equations at resonance*, *Comment. Math. Univ. Carolinae* **30** (1989), 537–550.
- [12] Naimark M.A., *Linear differential operators Part II*, Ungar Publ. Co., London, 1968.
- [13] Nkashama M.N., Santanilla J., *Existence of multiple solutions of some nonlinear boundary value problems*, *Jour. Diff. Eq.* **84** (1990), 148–164.
- [14] O'Regan D., *Singular Sturm Liouville problems and existence of solutions to singular nonlinear boundary value problems*, *Nonlinear Anal.* **20** (1993), 767–779.
- [15] ———, *Solvability of some two point boundary value problems of Dirichlet, Neumann or Periodic type*, *Dynamic Systems and Appl.* **2** (1993), 163–182.
- [16] ———, *Existence principles for second order nonresonant boundary value problems*, *Jour. Appl. Math. Stoch. Anal.* **7** (1994), 487–507.
- [17] ———, *Carathéodory theory of nonresonant second order boundary value problems*, to appear.

- [18] Ramas M., Sanchez L., *Variational elliptic problems involving noncoercive functionals*, Proc. Roy. Soc. Edinburgh **112A** (1989), 177–185.
- [19] Sanchez L., *Positive solutions for a class of semilinear two point boundary value problems*, Bull. Aust. Math. Soc. **45** (1992), 439–451.

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE GALWAY, GALWAY, IRELAND

(Received December 19, 1994, revised August 1, 1995)