

Sehie Park

On a problem of Gulevich on nonexpansive maps in uniformly convex Banach spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 37 (1996), No. 2, 263--268

Persistent URL: <http://dml.cz/dmlcz/118831>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On a problem of Gulevich on nonexpansive maps in uniformly convex Banach spaces

SEHIE PARK

Abstract. Let X be a uniformly convex Banach space, $D \subset X$, $f : D \rightarrow X$ a nonexpansive map, and K a closed bounded subset such that $\overline{\text{co}}K \subset D$. If (1) $f|_K$ is weakly inward and K is star-shaped or (2) $f|_K$ satisfies the Leray-Schauder boundary condition, then f has a fixed point in $\overline{\text{co}}K$. This is closely related to a problem of Gulevich [Gu]. Some of our main results are generalizations of theorems due to Kirk and Ray [KR] and others.

Keywords: uniformly convex, Banach space, Hilbert space, contraction, nonexpansive map, weakly inward map, demi-closed, Rothe condition, Leray-Schauder condition, (KR)-bounded, Opial's condition

Classification: Primary 47H10; Secondary 54H25

The well-known theorem of Browder-Göhde-Kirk assures existence of a fixed point for nonexpansive maps $f : K \rightarrow K$ where K is a *bounded* closed convex subset of a uniformly convex Banach space X . In [KR], Kirk and Ray showed that f can be replaced by a weakly inward nonexpansive map $f : K \rightarrow X$ while the boundedness of K can be replaced by that of the geometric estimator

$$G(x, fx) = \{z \in K : \|z - x\| \geq \|z - fx\|\}$$

for some $x \in K$ or more general sets. Note that any fixed point of f is contained in $G(x, fx)$.

On the other hand, Gulevich [Gu] considered the situation as follows: H is a Hilbert space, K is a nonempty bounded closed (not necessarily convex) subset of H , and $f : D \subset H \rightarrow H$ is a nonexpansive map, where $\overline{\text{co}}K \subset D$. Gulevich's basic theorem [Gu, Theorem 1] states that f has a fixed point in $\overline{\text{co}}K$ if f satisfies the Rothe condition $f(\text{Bd } K) \subset K$. He also raised as a problem whether H can be replaced by a uniformly convex Banach space.

In the present paper, we obtain some fixed point theorems on nonexpansive maps defined on closed (not necessarily bounded or convex) subsets of a Banach space. Our results are closely related to Gulevich's theorem and extend some known results of Kirk and Ray [KR], Goebel and Kuczumow [Go], and Browder [B1], [B2]. Moreover, we adopt more general boundary conditions on those nonexpansive maps. In fact, the weakly inwardness or the so-called Leray-Schauder condition is used in our results instead of the Rothe condition used in [Gu].

Recall that $f : K \rightarrow X$ is a *contraction* if there exists a $k \in [0, 1)$ such that

$$\|fx - fy\| \leq k\|x - y\| \quad \text{for all } x, y \in K;$$

and a *nonexpansive map* if

$$\|fx - fy\| \leq \|x - y\| \quad \text{for all } x, y \in K.$$

We say that f is *weakly inward* if $fx \in \overline{I_K(x)}$ for any $x \in \text{Bd } K$ (equivalently, for any $x \in K$), where $\overline{}$, Bd , and Int denote the closure, boundary, and interior, respectively, and

$$I_K(x) = \{x + c(y - x) : y \in K, c \geq 1\}.$$

Note that any map satisfying the Rothe condition is weakly inward.

We begin with the following:

Theorem 0. *Let K be a closed subset of a Banach space X and $f : K \rightarrow X$ a contraction satisfying one of the following:*

- (i) $f(\text{Bd } K) \subset K$.
- (ii) f is weakly inward.
- (iii) $0 \in \text{Int } K$ and $fx \neq mx$ for all $x \in \text{Bd } K$ and $m > 1$.

Then f has a unique fixed point.

Note that Theorem 0(i) is a particular case of Assad and Kirk [AK, Theorem 1], Theorem 0(ii) is due to Martinez-Yanez [M, Theorem] or, in a more general form, to Zhang [Z, Theorem 3.3], and Theorem 0(iii) to Gatica and Kirk [GK, Theorem 2.1]. There are more general results than Theorem 0. However, Theorem 0 is sufficient for our purpose. Note also that (iii) can be replaced by the following:

(iii)' there exists a $w \in \text{Int } K$ such that

$$fx - w \neq m(x - w) \quad \text{for all } x \in \text{Bd } K \quad \text{and } m > 1.$$

Moreover, (i) \Rightarrow (ii) and, whenever K is convex and $0 \in \text{Int } K$, we have (ii) \Rightarrow (iii).

A subset K of a vector space is said to be *star-shaped* if there exists a given point $x_0 \in K$ such that $tx_0 + (1 - t)x \in K$ for any $t \in (0, 1)$ and $x \in K$, where x_0 is called a *center* of K .

For the K and f in Theorem 0, we say that K is (KR)-*bounded* or bounded in the sense of Kirk-Ray [KR] if, for some bounded set $A \subset K$, the set

$$G(A) = \bigcap_{u \in A} G(u, fu)$$

is either empty or bounded.

The following is a generalization of the almost fixed point property of bounded closed subsets of a Banach space for nonexpansive maps.

Theorem 1. *Let X be a Banach space, K a closed subset of X , and $f : K \rightarrow X$ a nonexpansive map such that K is (KR)-bounded and one of the following holds:*

- (i) K is star-shaped and $f(\text{Bd } K) \subset K$.
- (ii) K is star-shaped and f is weakly inward.
- (iii) $0 \in \text{Int } K$ and $fx \neq mx$ for all $x \in \text{Bd } K$ and $m > 1$.

Then there exists a bounded sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$ as $n \rightarrow \infty$.

PROOF: For cases (i) and (ii) we may without loss of generality assume that 0 is the center. For $\alpha \in (0, 1)$, define $f_\alpha : K \rightarrow X$ by $f_\alpha x = \alpha fx$ for $x \in K$. Then clearly f_α is a contraction. We show that one of (i)–(iii) in Theorem 0 holds for f_α :

(i) Since K is star-shaped at center 0 , we have $\alpha K \subset K$. Since $f(\text{Bd } K) \subset K$, for $x \in \text{Bd } K$, we have $f_\alpha x = \alpha fx \in \alpha K \subset K$. Therefore, $f_\alpha(\text{Bd } K) \subset K$.

(ii) From $fx \in \bar{I}_K(x)$, we have $f_\alpha x = \alpha fx \in \alpha \bar{I}_K(x) \subset \bar{I}_K(x)$ since $I_K(x)$ is a star-shaped set with center 0 . See Zhang [Z, Theorem 1.2]. Note that (i) \Rightarrow (ii).

(iii) Suppose that $f_\alpha x = mx$ for some $x \in \text{Bd } K$ and $m > 1$, then $fx = \alpha^{-1} f_\alpha x = (\alpha^{-1} m)x$ and $\alpha^{-1} m > 1$, which contradicts our assumption.

Therefore, by Theorem 0, f_α has a fixed point $x_\alpha \in K$. Suppose that the set $\{x_\alpha : \alpha \in (0, 1)\}$ is not bounded. Then it is possible to choose $\alpha \in (0, 1)$ so that

$$\sup_{u \in A} \|fu\| \leq \inf_{u \in A} \|x_\alpha - u\|$$

and in addition, if $G(A) \neq \emptyset$, then α may also be chosen so that

$$\|x_\alpha\| > \sup\{\|x\| : x \in G(A)\}.$$

Therefore, for each $u \in A$,

$$\begin{aligned} \|x_\alpha - fu\| &= \|\alpha fx_\alpha - fu\| \leq \alpha \|fx_\alpha - fu\| + (1 - \alpha)\|fu\| \\ &\leq \alpha \|x_\alpha - u\| + (1 - \alpha)\|x_\alpha - u\| = \|x_\alpha - u\|. \end{aligned}$$

This implies $x_\alpha \in G(A)$, which is a contradiction. Thus $M = \sup\{\|x_\alpha\| : \alpha \in (0, 1)\} < \infty$ and we have

$$\|x_\alpha - fx_\alpha\| = (\alpha^{-1} - 1)\|x_\alpha\| \leq (\alpha^{-1} - 1)M,$$

yielding $\|x_\alpha - fx_\alpha\| \rightarrow 0$ as $\alpha \rightarrow 1$. This completes our proof. □

Note that Kirk and Ray [KR, Theorem 2.3] obtained Theorem 1(ii) for the case K is convex. In the second half of the proof of Theorem 1, we followed that of [KR, Theorem 2.3]. Note that Theorem 1(i) generalizes Dotson [D, Theorem 1].

A Banach space X is said to satisfy *Opial's condition* if, whenever a sequence $\{x_n\}$ converges weakly to $x_0 \in X$, then

$$\liminf_{n \rightarrow \infty} \|x_n - x\| > \liminf_{n \rightarrow \infty} \|x_n - x_0\|$$

for all $x \in X, x \neq x_0$. Opial [O] showed that if C is a weakly compact subset of a Banach space X satisfying this condition and $f : C \rightarrow X$ is nonexpansive, then $I - f$ is demi-closed ([B2], [Gö]); that is, if $\{x_n\} \subset C$ satisfies $x_n \rightarrow x$ weakly while $(I - f)x_n \rightarrow y$ strongly, then $(I - f)x = y$, where I is the identity map on C .

Examples of spaces satisfying Opial's condition are Hilbert spaces, l^p ($1 \leq p < \infty$), and uniformly convex Banach spaces with weakly continuous duality maps.

From Theorem 1, we have the following:

Theorem 2. *Let X be a Banach space, K a weakly compact subset of X , and $f : K \rightarrow X$ a nonexpansive map satisfying one of (i)–(iii) in Theorem 1.*

- (a) *If $I - f$ is demi-closed on K , then f has a fixed point.*
- (b) *If X satisfies Opial's condition, then f has a fixed point.*

PROOF: Since K is closed and bounded, f satisfies all the requirements of Theorem 1. Hence, there exists a sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since K is weakly compact, we may assume that $x_n \rightarrow x$ weakly to some $x \in K$. Since $x_n - fx_n \rightarrow 0$ strongly, $x_n - fx_n = (I - f)x_n$, and $I - f$ is demi-closed, we conclude that $(I - f)x = 0$ and hence $x = fx$. This completes our proof. \square

Note that Zhang [Z, Theorem 3.8 and Corollaries 3.10, 3.11] obtained the multi-valued version of Theorem 2(ii), with different proof, and that Theorem 2(i) generalizes Dotson [D, Theorem 2]. Note also that if K is compact, then f has a fixed point in Theorem 2 without assuming the demi-closedness of $I - f$.

From Theorem 1, we also have the following:

Theorem 3. *Let X be a uniformly convex Banach space, D a subset of X , and $f : D \rightarrow X$ a nonexpansive map. Let K be a closed (KR) -bounded subset of X such that $\overline{\text{co}} K \subset D$ and one of (i)–(iii) of Theorem 1 holds for $f|_K$. Then f has a fixed point in $\overline{\text{co}} K$.*

PROOF: Since $f|_K$ satisfies all the requirements of Theorem 1, there exists a bounded sequence $\{x_n\}$ in K such that $\|x_n - fx_n\| \rightarrow 0$. Since $\{x_n\}$ is contained in a bounded closed convex subset $L \subset \overline{\text{co}} K$ and L is weakly compact, we may assume $x_n \rightarrow x_0$ weakly to some $x_0 \in L$. Since $I - f$ is demi-closed on L ([B2], [Gö]) and $(I - f)x_n \rightarrow 0$ strongly, we conclude that $(I - f)x = 0$, and hence $x = fx$. This completes our proof. \square

If we can eliminate the star-shapedness in (i), then Theorem 3(i) will be the required affirmative answer to Gulevich's problem. Moreover, Gulevich [Gu] noted that, for case (i) of Theorem 3 in a Hilbert space H , f has a fixed point in K .

In case $0 \in \text{Int } K$, Theorem 3(iii) generalizes [Gu, Theorem 1].

Note that the set $A \subset K$ for the (KR)-boundedness can be chosen so that $A \subset D$ and $f(A) \subset K$. See the proof of Theorem 1. Therefore, Theorem 3 generalizes Ray [R, Lemma 1].

For $D = K = \overline{\text{co}} K$, Theorem 3 reduces to the following:

Theorem 4. *Let X be a uniformly convex Banach space, K a closed convex subset, and $f : K \rightarrow X$ a nonexpansive map such that K is (KR)-bounded and one of (i)–(iii) in Theorem 0 holds. Then f has a fixed point.*

Note that Theorem 4(i) and (ii) are due to Kirk and Ray [KR, Theorem 2.3], which extends Goebel and Kuczumow [Go, Theorem 6]. Also note that Theorem 4(iii) extends Browder [B2, Theorem 1] for nonexpansive maps. For a Hilbert space X and a closed ball in X , Theorem 4(i) is due to Browder [B1, Theorem 2], which was used to show existence of periodic solutions for nonlinear equations of evolution.

Recently Canetti, Marino, and Pietramala [CMP] obtained multi-valued versions of Theorem 4(ii) and, under the stronger assumption of convexity, some other results similar to Theorems 1–3 for case (ii).

Finally, we note that the so-called Rothe condition (i) was first adopted by Knaster, Kuratowski, and Mazurkiewicz [KKM]. Also, the origin of the so-called Leray-Schauder condition (iii) seems to be Schaefer [S], and the following are well-known examples of that condition:

$$(A) \quad \|fx - x\|^2 \geq \|fx\|^2 - \|x\|^2 \text{ for } x \in \text{Bd } K.$$

$$(K) \quad \text{Re} \langle fx, x \rangle = \|x\|^2 \text{ for } x \in \text{Bd } K, x \neq 0, \text{ in a Hilbert space } H.$$

Or more generally,

$$(P) \quad (fx, Jx) \leq (x, Jx) \text{ for } x \in \text{Bd } K, 0 \in \text{Int } K, \text{ where } J \text{ is any duality map of } X \text{ into } 2^{X^*}.$$

Condition (A) is due to Altman [A], (K) to Krasnosel'skii [K] and Shinbrot [Sh], and (P) to Petryshyn [P].

Acknowledgement. This work was supported in part by KOSEF-951-0102-006-2.

REFERENCES

- [A] Altman M., *A fixed point theorem for completely continuous operators in Banach spaces*, Bull. Acad. Polon. Sci. **3** (1955), 409–413.
- [AK] Assad N.A., Kirk W.A., *Fixed point theorems for set-valued mappings of contractive type*, Pac. J. Math. **43** (1972), 553–562.
- [B1] Browder F.E., *Existence of periodic solutions for nonlinear equations of evolution*, Proc. Nat. Acad. Sci. USA **53** (1965), 1100–1103.
- [B2] Browder F.E., *Semicontractions and semiaccretive nonlinear mappings in Banach spaces*, Bull. Amer. Math. Soc. **74** (1968), 660–665.
- [CMP] Canetti A., Marino G., Pietramala P., *Fixed point theorems for multivalued mappings in Banach spaces*, Nonlinear Anal. TMA **17** (1991), 11–20.

- [D] Dotson W.G., *Fixed point theorems for non-expansive mappings on star-shaped subsets of Banach spaces*, J. London Math. Soc. (2) **4** (1972), 408–410.
- [GK] Gatica J.A., Kirk W.A., *Fixed point theorems for contraction mappings with applications to nonexpansive and pseudo-contractive mappings*, Rocky Mount. J. Math. **4** (1974), 69–79.
- [Go] Goebel K., Kuczumow T., *A contribution to the theory of nonexpansive mappings*, Bull. Calcutta Math. Soc. **70** (1978), 355–357.
- [Gö] Göhde D., *Zum Prinzip der kontraktiven Abbildung*, Math. Nachr. **30** (1965), 251–258.
- [Gu] Gulevich N.M., *Existence of fixed points of nonexpansive mappings satisfying the Rothe condition*, J. Soviet Math. **26** (1984), 1607–1611.
- [KR] Kirk W.A., Ray W.O., *Fixed-point theorems for mappings defined on unbounded sets in Banach spaces*, Studia Math. **64** (1979), 127–138.
- [KKM] Knaster B., Kuratowski C., Mazurkiewicz S., *Ein Beweis des Fixpunktsatzes für n -dimensionale Simplexe*, Fund. Math. **14** (1929), 132–137.
- [K] Krasnosel'skii M.A., *New existence theorems for solutions of nonlinear integral equations*, Dokl. Akad. Nauk SSSR **88** (1953), 949–952.
- [M] Martinez-Yanez C., *A remark on weakly inward contractions*, Nonlinear Anal. TMA **16** (1991), 847–848.
- [O] Opial Z., *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
- [P] Petryshyn W.V., *A new fixed point theorem and its application*, Bull. Amer. Math. Soc. **78** (1972), 225–229.
- [R] Ray W.O., *Zeros of accretive operators defined on unbounded sets*, Houston J. Math. **5** (1979), 133–139.
- [S] Schaefer H.H., *Neue Existenzsätze in der Theorie nichtlinearer Integralgleichungen*, Ber. Verh. Sächs. Akad. Wiss. Leipzig Math.-Natur. Kl. **101** (1955), no.7, 40pp.
- [Sh] Shinbrot M., *A fixed point theorem and some applications*, Arch. Rational Mech. Anal. **17** (1964), 255–271.
- [Z] Zhang S., *Star-shaped sets and fixed points of multivalued mappings*, Math. Japonica **36** (1991), 327–334.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151–742, KOREA

E-mail: shpark@math.snu.ac.kr

(Received May 14, 1995)