

Billy E. Rhoades

A fixed point theorem for a multivalued non-self mapping

Commentationes Mathematicae Universitatis Carolinae, Vol. 37 (1996), No. 2, 401--404

Persistent URL: <http://dml.cz/dmlcz/118843>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A fixed point theorem for a multivalued non-self mapping

B.E. RHOADES

Abstract. We prove a fixed point theorem for a multivalued non-self mapping in a metricaly convex complete metric space. This result generalizes Theorem 1 of Itoh [2].

Keywords: multivalued non-self mapping, metricaly convex metric space

Classification: 47H10, 54H25

Let $CB(X)$ denote the family of all nonempty closed bounded subsets of X , H the Hausdorff metric on X , and for any set A , $D(x, A)$ denotes the distance from x to A .

Lemma 1 ([3]). *If $A, B \in CB(X)$ and $x \in A$, then for any positive number ε there exists a $y \in B$ such that $d(x, y) \leq H(A, B) + \varepsilon$.*

Lemma 2 ([1]). *Let K be a nonempty closed subset of a complete metricaly convex metric space (X, d) . Then, for any $x \in K$, $y \notin K$, there exists a point $z \in \partial K$ such that $d(x, z) + d(z, y) = d(x, y)$.*

Let $F : X \rightarrow CB(X)$. We shall be interested in the following contractive definition. For each x, y in X ,

$$(1) \quad H(Fx, Fy) \leq \alpha d(x, y) + \beta \max\{D(x, Fx), D(y, Fy)\} + \gamma [D(x, Fy) + D(y, Fy)],$$

where $\alpha, \beta, \gamma \geq 0$ and such that

$$s := \left(\frac{1 + \alpha + \gamma}{1 - \beta - \gamma} \right) \left(\frac{\alpha + \beta + \gamma}{1 - \gamma} \right) < 1.$$

Theorem 1. *Let (X, d) be a complete metricaly convex metric space, K a nonempty closed subset of X . Let $F : X \rightarrow CB(X)$ satisfy (1). If for each $x \in \partial K$, $Fx \subset K$, then there exists a $z \in K$ such that $z \in Fz$.*

PROOF: We shall construct two sequences $\{x_n\}$ and $\{x'_n\}$ as follows. Let $x_0 \in K$. Let $x'_1 \in Tx_0$. If $x'_1 \in K$, set $x_1 = x'_1$. If $x'_1 \notin K$, choose x_1 so that $d(x_0, x_1) + d(x_1, x'_1) = d(x_0, x'_1)$. Pick $x'_2 \in Fx_1$ so that $d(x_1, x'_2) \leq H(Fx_0, Fx_1) + (1 - \beta - \gamma)\varepsilon$, with $\varepsilon = s$. If $x'_2 \in K$, set $x_2 = x'_2$. If $x'_2 \notin K$, since X is metricaly convex, choose x_2 so that $d(x_1, x_2) + d(x_2, x'_2) = d(x_1, x'_2)$. Continuing in this way

we obtain two sequences $\{x_n\}, \{x'_n\}$ satisfying the following properties: $x'_{n+1} \in Fx_n$ and such that $d(x'_n, x'_{n+1}) \leq H(Fx_{n-1}, Fx_n) + (1 - \beta - \gamma)\varepsilon^n$; $x_{n+1} = x'_{n+1}$ if $x'_{n+1} \in K$; and, if $x'_{n+1} \notin K$, x_{n+1} is chosen to satisfy $d(x_n, x_{n+1}) + d(x_{n+1}, x'_{n+1}) = d(x_n, x'_{n+1})$.

Define

$$P = \{x_n \in \{x_n\} : x_n = x'_n\}, \quad Q = \{x_n \in \{x_n\} : x_n \neq x'_n\}.$$

Three cases now arise.

Case 1. $x_n, x_{n+1} \in P$. From (1),

$$\begin{aligned} d(x_n, x_{n+1}) &\leq H(Fx_{n-1}, Fx_n) + (1 - \beta - \gamma)\varepsilon^n \\ &\leq \alpha d(x_{n-1}, x_n) + \beta \max\{D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n)\} \\ &\quad + \gamma[D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})] + (1 - \beta - \gamma)\varepsilon^n \\ &\leq \alpha d(x_{n-1}, x_n) + \beta \max\{d(x_{n-1}, x_n), d(x_n, x_{n+1})\} \\ &\quad + \gamma d(x_{n-1}, x_{n+1}) + (1 - \beta - \gamma)\varepsilon^n \\ (2) \quad &\leq \max \left\{ \frac{(\alpha + \beta + \gamma)d(x_{n-1}, x_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \gamma}, \right. \\ &\quad \left. \frac{(\alpha + \gamma)d(x_{n-1}, x_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \beta - \gamma} \right\} \\ &\leq \max \left\{ \frac{\alpha + \beta + \gamma}{1 - \gamma}, \frac{\alpha + \gamma}{1 - \beta - \gamma} \right\} d(x_{n-1}, x_n) + \varepsilon^n \\ &= kd(x_{n-1}, x_n) + \varepsilon^n, \end{aligned}$$

where

$$k := \frac{\alpha + \beta + \gamma}{1 - \gamma}.$$

Case 2. $x_n \in P, x_{n+1} \in Q$. From (1),

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_n, x'_{n+1}) \leq H(Fx_{n-1}, Fx_n) + (1 - \beta - \gamma)\varepsilon^n \\ &\leq \alpha d(x_{n-1}, x_n) + \beta \max\{D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n)\} \\ &\quad + \gamma[D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})] + (1 - \beta - \gamma)\varepsilon^n \\ &\leq \alpha d(x_{n-1}, x_n) + \beta \max\{d(x_{n-1}, x_n), d(x_n, x'_{n+1})\} \\ &\quad + \gamma d(x_{n-1}, x'_{n+1}) + (1 - \beta - \gamma)\varepsilon^n. \end{aligned}$$

If the maximum of the coefficient of β is $d(x_{n-1}, x_n)$, then we have

$$d(x_n, x'_{n+1}) \leq \frac{(\alpha + \beta + \gamma)d(x_{n-1}, x_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \gamma}.$$

If the maximum of the coefficient of β is $d(x_n, x'_{n+1})$, then we have

$$d(x_n, x'_{n+1}) \leq \frac{(\alpha + \gamma)d(x_{n-1}, x_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \beta - \gamma}.$$

Therefore, in all cases,

$$(3) \quad d(x_n, x_{n+1}) \leq kd(x_{n-1}, x_n) + \varepsilon^n.$$

Case 3. $x_n \in Q, x_{n+1} \in P$. Then, using (1) and Case 2,

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_n, x'_n) + d(x'_n, x_{n+1}) \\ &\leq d(x_n, x'_n) + H(Fx_{n-1}, Fx_n) + (1 - \beta - \gamma)\varepsilon^n \\ &\leq d(x_{n-1}, x'_n) + \alpha d(x_{n-1}, x_n) \\ &\quad + \beta \max\{D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n)\} \\ &\quad + \gamma[D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})] + (1 - \beta - \gamma)\varepsilon^n \\ &\leq d(x_{n-1}, x'_n) + \alpha d(x_{n-1}, x_n) + \beta \max\{d(x_{n-1}, x'_n), d(x_n, x_{n+1})\} \\ &\quad + \gamma[d(x_{n-1}, x_{n+1}) + d(x_n, x'_n)] + (1 - \beta - \gamma)\varepsilon^n. \end{aligned}$$

Note that

$$\begin{aligned} d(x_{n-1}, x_{n+1}) + d(x_n, x'_n) &\leq d(x_{n-1}, x_n) + d(x_n, x_{n+1}) + d(x_n, x'_n) \\ &= d(x_{n-1}, x'_n) + d(x_n, x_{n+1}). \end{aligned}$$

Therefore

$$\begin{aligned} d(x_n, x_{n+1}) &\leq \left\{ \frac{(1 + \alpha + \beta + \gamma)d(x_{n-1}, x'_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \gamma} \right. \\ &\quad \left. \frac{(1 + \alpha + \gamma)d(x_{n-1}, x'_n) + (1 - \beta - \gamma)\varepsilon^n}{1 - \beta - \gamma} \right\} \\ &\leq \max \left\{ \frac{1 + \alpha + \beta + \gamma}{1 - \gamma}, \frac{1 + \alpha + \gamma}{1 - \beta - \gamma} \right\} d(x_{n-1}, x'_n) + \varepsilon^n \\ &\leq \left(\frac{1 + \alpha + \gamma}{1 - \beta - \gamma} \right) d(x_{n-1}, x'_n) + \varepsilon^n \\ &\leq sd(x_{n-2}, x_{n-1}) + \frac{(1 + \alpha + \gamma)\varepsilon^{n-1}}{1 - \beta - \gamma} + \varepsilon^n. \end{aligned}$$

Using (2)–(4), it can be shown by induction that

$$d(x_{2n}, x_{2n+1}) \leq s^n \left(\delta + \frac{3n}{1 - \beta - \gamma} \right)$$

and that

$$d(x_{2n+1}, x_{2n+2}) \leq s^{(2n+1)/2} \left(\delta + \frac{3n+1}{1-\beta-\gamma} \right).$$

Then, for any $m > n$,

$$d(x_m, x_n) \leq \sum_{i=n}^{m-1} d(x_i, x_{i+1}) \leq \delta \sum_{i=n}^{m-1} s^{i/2} + \frac{1}{1-\beta-\gamma} \sum_{i=n}^{m-1} s^{i/2} (3i+1),$$

and $\{x_n\}$ is Cauchy, hence convergent. Call the limit z . From the way in which the $\{x_n\}$ were chosen, there exists an infinite subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} = x'_{n_k}$.

$$\begin{aligned} D(x_{n_k}, Fz) &\leq H(Fx_{n_i-1}, Fz) \\ &\leq \alpha d(x_{n_k-1}, z) + \beta \max\{D(x_{n_k-1}, Fx_{n_k-1}), D(z, Fz)\} \\ &\quad + \gamma [D(x_{n_k-1}, Fz) + D(z, Fx_{n_k-1})] \\ &\leq \alpha d(x_{n_k-1}, z) + \beta \max\{d(x_{n_k-1}, x_{n_k}), D(z, Fz)\} \\ &\quad + \gamma [D(x_{n_k-1}, Fz) + d(z, x_{n_k})]. \end{aligned}$$

Taking the limit as $k \rightarrow \infty$ yields $D(z, Fz) \leq (\beta + \gamma)D(z, Fz)$, which implies that $z \in Fz$. \square

REFERENCES

- [1] Blumenthal L.M., *Theory and Applications of Distance Geometry*, The Clarendon Press, Oxford, 1943.
- [2] Itoh Shigeru, *Multivalued generalized contractions and fixed point theorems*, Comment. Math. Univ. Carolinae **18** (1977), 247–258.
- [3] Nadler S.B., Jr., *Multi-valued contraction mappings*, Pacific J. Math. **30** (1969), 475–488.

INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405–5701, USA

(Received September 2, 1994, revised November 24, 1995)