

Isaac Gorelic

The G_δ -topology and incompactness of ω^α

Commentationes Mathematicae Universitatis Carolinae, Vol. 37 (1996), No. 3, 613--616

Persistent URL: <http://dml.cz/dmlcz/118867>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

The G_δ -topology and incompactness of ω^α

ISAAC GORELIC

Abstract. We establish a relation between covering properties (e.g. Lindelöf degree) of two standard topological spaces (Lemmas 4 and 5). Some cardinal inequalities follow as easy corollaries.

Keywords: Lindelöf degree, G_δ topology, cardinal functions

Classification: 54D20, 45B10

The present note is a contribution into the study of the Lindelöf degree in powers of topological spaces. It answers a question of W.A.R. Weiss.

In what follows, $\kappa^* \subset \beta\kappa$ is the space of all free ultrafilters over a discrete set of κ points, $\mu\kappa \subset \beta\kappa$ is the space of all *uniform* ultrafilters over κ , ω^α denotes the α -th power of the discrete set of integers, $(\mu\kappa, G_\delta)$ is $\mu\kappa$ with the finer G_δ topology. $L(X)$ denotes the Lindelöf degree of X , and $e(X) := \sup\{A \subset X : A \text{ is closed and discrete}\}$ — its *extent*.

Kenneth Kunen ([1]) proves $L(\mu(2^\kappa)^+, G_\delta) \geq \kappa^+$, for the same κ 's as in our Corollary 6. Our Corollary 6 gives here

$$L(\mu(2^\kappa)^+, G_\delta) \geq (2^\kappa)^+.$$

J. Mycielski proved ([2]), by inductive “stepping up”, that, for α less than the 1st weakly inaccessible cardinal,

$$e(\omega^\alpha) = \alpha.$$

Our Corollary 9 is a weaker statement for a larger class of cardinals. This Corollary was obtained first by Łoś [3] in 1959 using group-theoretic methods. See also Juhász [4].

Stevo Todorčević ([5]) proves, assuming the combinatorial statement \square_κ ,

$$L(\omega^\kappa) = \kappa.$$

1. If $\mathcal{A} = \{A_n : n < \omega\}$ is a countable disjoint partition of the cardinal κ , then

$$\begin{aligned} \mu\kappa &= \left(\bigcup_{n < \omega} S_n^{\mathcal{A}} \right) \dot{\cup} (\tilde{S}^{\mathcal{A}}), \text{ where} \\ S_n^{\mathcal{A}} &= \{u \in \mu\kappa : A_n \in u\} \text{ and} \\ \tilde{S}^{\mathcal{A}} &= \{u \in \mu\kappa : \{\bigcup_{n \geq i} A_n : i < \omega\} \subset u\}. \end{aligned}$$

Note that $\tilde{S}^{\mathcal{A}}$ is a G_δ set in $\mu\kappa$.

2. We say that a cover of κ^* or of $\mu\kappa$ is a *proper G_δ -cover* if every set in it is of the form $\tilde{S}^{\mathcal{A}}$ for some countable partition \mathcal{A} of κ .

3. **Lemma.** *If κ is a regular cardinal and $\mu\kappa$ has a proper G_δ -cover of size α , then ω^α has a subset of size κ without a CAP (complete accumulation point).*

PROOF: Suppose $\mu\kappa = \bigcup \{\tilde{S}^{\mathcal{A}^\gamma} : \gamma < \alpha\}$ for some collection $\mathcal{C} = \{\mathcal{A}^\gamma : \gamma < \alpha\}$ of countable partitions $\mathcal{A}^\gamma = \{A_n^\gamma : n < \omega\}$ of κ . For every point $p \in \kappa$ define its history in \mathcal{C} $\bar{p} : \alpha \rightarrow \omega$ by setting $\bar{p}(\gamma) := n$ such that $p \in A_n^\gamma$. Let $P = \{\bar{p} : p < \kappa\} \subset \omega^\alpha$.

Claim 1. $|P| = \kappa$, moreover, for every $p \in \kappa$, $K_p := \{q \in \kappa : \bar{q} = \bar{p}\}$ has size $|K_p| < \kappa$. Indeed, if not, then no $v \ni K_p$ is covered:

$$\forall \gamma < \alpha \quad v \notin \tilde{S}^{\mathcal{A}^\gamma},$$

because

$$v \in S_{\bar{p}(\gamma)}^{\mathcal{A}^\gamma}.$$

And $|P| = \kappa$ follows from the regularity of κ . □

Claim 2. P has no CAP in ω^α . If not, let $\varphi \in \omega^\alpha$ be a CAP of P . Then for every finite $F \subset \alpha$

$$|\{p < \kappa : \bar{p} \upharpoonright F = \varphi \upharpoonright F\}| = \kappa,$$

by Claim 1.

Therefore, the family $\mathcal{F} := \{A_{\varphi(\gamma)}^\gamma : \gamma < \alpha\}$ has the uniform finite intersection property (i.e. $\forall \mathcal{F}_0 \in [\mathcal{F}]^{< \aleph_0} \cap \mathcal{F}_0 \neq \emptyset$). [By $\bar{p} \upharpoonright F = \varphi \upharpoonright F \iff p \in \bigcap_{\gamma \in F} A_{\varphi(\gamma)}^\gamma$].

Pick a $u \in \mu\kappa$ extending \mathcal{F} .

Then $u \notin \bigcup_{\gamma < \alpha} \tilde{S}^{\mathcal{A}^\gamma} = \mu\kappa$. Contradiction. Hence $P \subset \omega^\alpha$ has no CAP in ω^α , so it is as required. □

4. Lemma. *If κ^* has a proper G_δ -cover of size α , then ω^α has a closed discrete subset of size κ .*

PROOF: Here we, similarly, study the family P of the histories of points $p \in \kappa$ in the family of partitions of κ defining our proper G_δ -cover, in this case of κ^* . Only finitely many points $p \in \kappa$ may have the same history, so $|P| = \kappa$, and, arguing as in Claim 2 of the previous lemma, $P \subset \omega^\alpha$ has no limit points in ω^α whatsoever. \square

5. Theorem. *If κ is a regular not Ulam measurable cardinal, then*

$$L(\omega^{L(\mu\kappa, G_\delta)}) \geq \kappa.$$

6. Corollary. *$L(\mu\kappa, G_\delta) \geq \kappa$, for the same κ 's.*

PROOF OF THEOREM 5 AND COROLLARY 6: Since every ultrafilter over κ is countably incomplete, there is a proper G_δ cover of $\mu\kappa$, and so $L(\mu\kappa, G_\delta) = \alpha \Rightarrow$ there is a proper G_δ -cover of size $\alpha \Rightarrow$ (By Lemma 3) ω^α has a subset of size κ without a CAP \Rightarrow

- (a) $L(\omega^\alpha) \geq \kappa$, and
- (b) $\alpha \geq \kappa$ (because $\alpha \geq L(\omega^\alpha)$).

\square

7. Corollary. *$L(\omega^{2^\kappa}) \geq \kappa$, for the same κ 's as in Theorem 5.*

PROOF: $(\mu\kappa, G_\delta)$ has a base of size $(2^\kappa)^\omega = 2^\kappa$. Hence $L(\mu\kappa, G_\delta) \leq 2^\kappa$ and so $L(\omega^{2^\kappa}) \geq L(\omega^{L(\mu\kappa, G_\delta)}) \geq \kappa$. \square

8. Theorem. *If κ is not Ulam measurable, then*

$$L(\kappa^*, G_\delta) \geq L(\omega^{L(\kappa^*, G_\delta)}) \geq e(\omega^{L(\kappa^*, G_\delta)}) \geq \kappa.$$

PROOF: Immediate from Lemma 4. \square

9. Corollary. *If $\kappa < 1^{st}$ measurable cardinal, then*

$$e(\omega^{2^\kappa}) \geq \kappa,$$

i.e. ω^{2^κ} has a closed discrete subspace of size κ .

PROOF: Same as of Corollary 7. \square

10. Corollary. *Let λ be a strong limit cardinal \leq the 1^{st} measurable cardinal. Then the set $\{e(\omega^\alpha) : \alpha < \lambda\}$ is cofinal in λ . Hence, if $cf(\lambda) > \omega$, the set $\{\alpha < \lambda : e(\omega^\alpha) = \alpha\}$ is closed and unbounded in λ .*

Remark. Murray Bell observed that the converses of Lemmas 3 and 4 are also true.

Acknowledgement. The author is very grateful to Professor William Weiss for many useful discussions.

REFERENCES

- [1] Kunen K., *Box products of ordered spaces*, Topology and its applications **20** (1985).
- [2] Mycielski J., *α -incompactness of N^α* , Bull. Acad. Pol., vol. XII, no. 8, 1964.
- [3] Łoś J., *Linear equations and pure subgroups*, Bull. Acad. Polon. Sci. **7** (1959).
- [4] Juhász I., *On closed discrete subspaces of product spaces*, Bull. Acad. Pol., vol. XVII, no. 4, 1969.
- [5] Todorčević S., *Incompactness of \mathbb{N}^θ* , Handwritten notes, 1990.

UNIVERSITY OF TORONTO, CANADA

(Received July 19, 1995)