

Mircea Balaj

Separation of $(n + 1)$ -families of sets in general position in \mathbf{R}^n

Commentationes Mathematicae Universitatis Carolinae, Vol. 38 (1997), No. 4, 743--748

Persistent URL: <http://dml.cz/dmlcz/118969>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1997

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Separation of $(n + 1)$ -families of sets in general position in \mathbf{R}^n

MIRCEA BALAJ

Abstract. In this paper the main result in [1], concerning $(n + 1)$ -families of sets in general position in \mathbf{R}^n , is generalized. Finally we prove the following theorem: If $\{A_1, A_2, \dots, A_{n+1}\}$ is a family of compact convexly connected sets in general position in \mathbf{R}^n , then for each proper subset I of $\{1, 2, \dots, n + 1\}$ the set of hyperplanes separating $\cup\{A_i : i \in I\}$ and $\cup\{A_j : j \in \bar{I}\}$ is homeomorphic to S_n^+ .

Keywords: family of sets in general position, convexly connected sets, Fan-Glicksberg-Kakutani fixed point theorem

Classification: Primary 52A37; Secondary 47H10

1. Introduction

In this paper we continue the investigation of a previous article [1], regarding the separability of the members of an $(n + 1)$ -family of sets in general position in \mathbf{R}^n . In the beginning we recall some definitions and notations.

A family \mathcal{A} of sets in \mathbf{R}^n is said to be in general position if any m -flat, $0 \leq m \leq n - 1$, intersects at most $m + 1$ members of \mathcal{A} . Let $m = \min\{n + 1, \text{card } \mathcal{A}\}$. It is easy to see that the family \mathcal{A} is in general position if and only if for every choice of sets $A_1, A_2, \dots, A_m \in \mathcal{A}$ and every choice of points $x_1 \in A_1, x_2 \in A_2, \dots, x_m \in A_m$, the set $\{x_1, x_2, \dots, x_m\}$ is affinely independent.

A set $A \subset \mathbf{R}^n$ is called (cf.[5] and [8, p. 174]) convexly connected if there is no hyperplane H such that $H \cap A = \emptyset$ and A contains points in both open halfspaces determined by H .

If A is a compact set and H a hyperplane in \mathbf{R}^n , then the distance between A and H is defined to be $d(A, H) = \min\{\|x - y\| : x \in A, y \in H\}$. If $H = \{x \in \mathbf{R}^n : \langle x, b \rangle = \lambda\}$ is a hyperplane, the corresponding closed halfspace $\{x \in \mathbf{R}^n : \langle x, b \rangle \leq \lambda\}$, $\{x \in \mathbf{R}^n : \langle x, b \rangle \geq \lambda\}$ are denoted respectively by H^{\leq} , H^{\geq} . A set A is said to be separated from a set B by the hyperplane H provided that A lies in one of the closed halfspaces H^{\leq} , H^{\geq} and B lies in the other. The set A is strictly separated from B by H provided that the separating hyperplane H is disjoint from both A and B . If \mathcal{A} is a family of sets containing at least two members, we say that a hyperplane H separates the members of \mathcal{A} if there exists a nontrivial partition $(\mathcal{B}, \mathcal{C})$ of \mathcal{A} such that $\cup \mathcal{B} \subset H^{\leq}$, $\cup \mathcal{C} \subset H^{\geq}$.

The unit sphere in \mathbf{R}^{n+1} and the set $\{x = (x_1, x_2, \dots, x_{n+1}) : \|x\| = 1, x_i \geq 0, 1 \leq i \leq n + 1\}$ are denoted by S_n, S_n^+ respectively. For every subset I of $\{1, 2, \dots, n + 1\}, \bar{I}$ denotes the complement of I in $\{1, 2, \dots, n + 1\}$.

In [1] among other results we have obtained the following

Theorem 1. *Let $\{A_1, A_2, \dots, A_{n+1}\}$ be a family of compact convexly connected sets in general position in \mathbf{R}^n . Then*

(i) *for each proper subset I of $\{1, 2, \dots, n + 1\}$, there exists exactly one hyperplane H such that*

$$(1) \quad H \text{ separates strictly the sets } \cup \{A_i : i \in I\}, \cup \{A_j : j \in \bar{I}\}$$

and

$$(2) \quad d(A_1, H) = d(A_2, H) = \dots = d(A_{n+1}, H);$$

(ii) *there exist exactly $2^n - 1$ hyperplanes satisfying (2).*

In this paper we obtain a generalization of the previous result. Also, we prove that for every nontrivial partition $(\mathcal{B}, \mathcal{C})$ of an $(n + 1)$ -family of compact convexly connected sets in general position in \mathbf{R}^n , the set of hyperplanes separating $\cup \mathcal{B}$ and $\cup \mathcal{C}$ is homeomorphic to S_n^+ .

2. Basic results

We start with the following result which generalizes Lemma 3 in [1].

Lemma 2. *Let $[x_1, x_2, \dots, x_{n+1}]$ be an n -simplex in \mathbf{R}^n . Then for each $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{n+1}) \in S_n^+$ and for each proper subset I of $\{1, 2, \dots, n + 1\}$ there exists exactly one hyperplane H such that*

$$(3) \quad H \text{ separates the sets } \{x_i : i \in I\} \text{ and } \{x_j : j \in \bar{I}\},$$

$$(4) \quad d(x_i, H) = k\alpha_i \text{ for some } k \text{ and all } i, \quad 1 \leq i \leq n + 1.$$

PROOF: The distance from an arbitrary point x_0 to a hyperplane

$$(5) \quad H = \{x \in \mathbf{R}^n : \langle x, b \rangle = \lambda\}$$

is given by the Ascoli's formula (see [6, p.21])

$$(6) \quad d(x_0, H) = \frac{|\langle x_0, b \rangle - \lambda|}{\|b\|}.$$

Since the pair $(b, \lambda) \in (\mathbf{R}^n \setminus \{0\}) \times \mathbf{R}$ for which the hyperplane H admits the representation (5) is unique up to a non-zero multiplicative constant, the conditions (3) and (4) are equivalent with

$$(7) \quad \langle x_i, b \rangle - \lambda = \beta_i, \quad 1 \leq i \leq n + 1$$

where

$$\beta_i = \begin{cases} \alpha_i, & \text{if } i \in I, \\ -\alpha_i, & \text{if } i \in \bar{I}. \end{cases}$$

Denoting by $(x_{i1}, x_{i2}, \dots, x_{in})$ the coordinates of x_i , $1 \leq i \leq n + 1$, and by (b_1, b_2, \dots, b_n) the coordinates of b , we are lead to an $(n + 1) \times (n + 1)$ linear system

$$(8) \quad x_{i1}b_1 + x_{i2}b_2 + \dots + x_{in}b_n - \lambda = \beta_i, \quad 1 \leq i \leq n + 1.$$

From the affine independence of the points x_1, x_2, \dots, x_{n+1} , it follows that the determinant D of order $n + 1$ having the general row $(x_{i1}, x_{i2}, \dots, x_{in}, 1)$ is different from zero. This proves that the system (8) possesses the unique solution

$$(9) \quad \begin{cases} b_j = \frac{D_j}{D}, & 1 \leq j \leq n \\ \lambda = -\frac{D_{n+1}}{D}, \end{cases}$$

where D_j is the determinant of order $n + 1$ having the general row $(x_{i1}, x_{i2}, \dots, x_{i,j-1}, \beta_i, x_{i,j+1}, \dots, x_{in}, 1)$ and D_{n+1} is the determinant of the same order, with the general row $(x_{i1}, \dots, x_{in}, \beta_i)$. Since at least two β_i are distinct, from (7) it can be easily deduced that $b \neq 0$. All these show that there exists a unique hyperplane H which satisfies the conditions (3) and (4). Note that $d(x_i, H) = \frac{\alpha_i}{\|b\|}$ and that the points x_i , $i \in I$, lie in the closed halfspace H^{\geq} , while the points x_j , $j \in \bar{I}$, lie in H^{\leq} . □

Let a point α_0 lie on the surface S_n^+ with all coordinates equal. The proof of the following lemma repeats the previous proof (taking $I = \{1, 2, \dots, n + 1\}$).

Lemma 3. *Let $\Delta = [x_1, x_2, \dots, x_{n+1}]$ be an n -simplex in \mathbf{R}^n . Then for every $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{n+1}) \in S_n^+ \setminus \{\alpha_0\}$ there exists exactly one hyperplane H such that*

- (i) *the simplex Δ is contained in one of the closed half-spaces determined by H , and*
- (ii) *$d(x_i, H) = k\alpha_i$ for some k and all i , $1 \leq i \leq n + 1$.*

The following generalization of Theorem 1 is our main result.

Theorem 4. *Let $\{A_1, A_2, \dots, A_{n+1}\}$ be a family of compact convexly connected sets in general position in \mathbf{R}^n . Then*

- (i) *for each $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{n+1}) \in S_n^+$ and for each proper subset I of $\{1, 2, \dots, n + 1\}$ there exists exactly one hyperplane H such that*

$$(10) \quad H \text{ separates the sets } \cup \{A_i : i \in I\} \text{ and } \cup \{A_j : j \in \bar{I}\}$$

and

$$(11) \quad d(A_i, H) = k\alpha_i \text{ for some } k \text{ and all } i, \quad 1 \leq i \leq n + 1;$$

(ii) if for each $\alpha \in S_n^+$, $N(\alpha)$ denotes the number of the hyperplanes H satisfying (11), then

$$N(\alpha) = \begin{cases} 2^n, & \text{if } \alpha \in S_n^+ \setminus \{\alpha_0\} \\ 2^n - 1, & \text{if } \alpha = \alpha_0. \end{cases}$$

PROOF: (i) A similar argument to that used in proving Corollary 7 in [1] permits us to suppose the compact sets A_i being convex. Let $A = A_1 \times A_2 \times \dots \times A_{n+1}$. The elements of A are denoted by \bar{x}, \bar{y}, \dots . Let I be an arbitrary fixed proper subset of $\{1, 2, \dots, n + 1\}$. By Lemma 2, for each $\bar{x} = (x_1, x_2, \dots, x_{n+1}) \in A$, ($x_i \in A_i$, $1 \leq i \leq n + 1$) there exists a unique hyperplane, denoted by $H(\bar{x})$, such that

$$(12) \quad d(x_i, H(\bar{x})) = k\alpha_i \text{ for some } k \text{ (dependent on } \bar{x}) \text{ and all } i, \quad 1 \leq i \leq n + 1,$$

$\{x_i : i \in I\} \subset H^{\geq}(\bar{x})$ and $\{x_j : j \in \bar{I}\} \subset H^{\leq}(\bar{x})$. The equation of $H(\bar{x})$ is $\langle x, b \rangle = \lambda$, where $b = (b_1, b_2, \dots, b_n) \in \mathbf{R}^n \setminus \{0\}$ and $\lambda \in \mathbf{R}$ are given by the formulas (9).

We define the map $f : A \rightarrow 2^A$ by $f(\bar{x}) = P_1(\bar{x}) \times P_2(\bar{x}) \times \dots \times P_{n+1}(\bar{x})$, $\bar{x} \in A$ and $P_i(\bar{x})$ defined by

$$(13) \quad \begin{cases} \text{(a)} & P_i(\bar{x}) = \{x \in A_i : \langle x, b \rangle = \min\{\langle y, b \rangle : y \in A_i\}\}, & i \in I, \\ \text{(b)} & P_i(\bar{x}) = \{x \in A_i : \langle x, b \rangle = \max\{\langle y, b \rangle : y \in A_i\}\}, & i \in \bar{I}. \end{cases}$$

Since the sets A_i are compact, the sets $P_i(\bar{x})$ are nonempty. If $y_i \in P_i(\bar{x})$, $1 \leq i \leq n + 1$, then $P_i(\bar{x})$ coincides with the intersection of the set A_i with the hyperplane through y_i parallel to $H(\bar{x})$. Thus each $P_i(\bar{x})$ is a compact convex set, and $f(\bar{x})$ is a compact convex set for each $\bar{x} \in A$. Using Lemma 4 in [1] it can be easily verified that f is upper semicontinuous.

By the Fan-Glicksberg-Kakutani fixed point theorem (see [2] and [4]), there is a point $\bar{z} = (z_1, z_2, \dots, z_{n+1}) \in A$ such that $\bar{z} \in f(\bar{z})$. Let $\langle x, b^0 \rangle - \lambda^0 = 0$ be the equation of the hyperplane $H(\bar{z})$, with $b^0 = (b_1^0, b_2^0, \dots, b_n^0)$ and λ^0 given by the formulas (9). For each $i \in I$, $z_i \in H^{\geq}(\bar{z})$ and by definition of f , $z_i \in P_i(\bar{z})$. Thus, we infer from (13a) that $A_i \subset H^{\geq}(\bar{z})$ for all $i \in I$.

Then, for each $i \in I$, we have

$$d(A_i, H(\bar{z})) = \min \left\{ \frac{|\langle x, b^0 \rangle - \lambda^0|}{\|b^0\|} : x \in A_i \right\} = \frac{\langle z_i, b^0 \rangle - \lambda^0}{\|b^0\|} = d(z_i, H(\bar{z})).$$

In a similar manner, we obtain that $A_j \subset H^{\leq}(\bar{z})$ and $d(A_j, H(\bar{z})) = d(z_j, H(\bar{z}))$, for all $j \in \bar{I}$. Therefore $H(\bar{z})$ separates the sets $\cup\{A_i : i \in I\}$, $\cup\{A_j : j \in \bar{I}\}$ and by (12) the sets $\{d(A_1, H(\bar{z})), d(A_2, H(\bar{z})), \dots, d(A_{n+1}, H(\bar{z}))\}$, $\{\alpha_1, \alpha_2, \dots, \alpha_{n+1}\}$ are proportional.

In the second part of the proof we verify the uniqueness of the hyperplane H which satisfies (10) and (11), for an arbitrary fixed set of indices I .

By the way of contradiction, suppose that there exist two distinct hyperplanes $H' = \{x \in \mathbf{R}^n : \langle x, b' \rangle - \lambda' = 0\}$, $H'' = \{x \in \mathbf{R}^n : \langle x, b'' \rangle - \lambda'' = 0\}$ satisfying (10) and (11). For each $i \in \{1, 2, \dots, n + 1\}$ let x'_i and x''_i be points in A_i such that $d(A_i, H') = d(x'_i, H')$, $d(A_i, H'') = d(x''_i, H'')$. Then, for a convenient choice of the pairs (b', λ') , (b'', λ'') we have

$$(14) \quad \begin{cases} (a) & \min\{\langle x, b' \rangle - \lambda' : x \in A_i\} = \langle x'_i, b' \rangle - \lambda' = \alpha_i & \text{if } i \in I, \\ (b) & \max\{\langle x, b' \rangle - \lambda' : x \in A_j\} = \langle x'_j, b' \rangle - \lambda' = -\alpha_j & \text{if } i \in \bar{I} \end{cases}$$

and

$$(15) \quad \begin{cases} (a) & \min\{\langle x, b'' \rangle - \lambda'' : x \in A_i\} = \langle x''_i, b'' \rangle - \lambda'' = \alpha_i & \text{if } i \in I, \\ (b) & \max\{\langle x, b'' \rangle - \lambda'' : x \in A_j\} = \langle x''_j, b'' \rangle - \lambda'' = -\alpha_j & \text{if } i \in \bar{I}. \end{cases}$$

Then, for each $i \in I$, by (14a) and (15a) it follows that $\langle x'_i, b' - b'' \rangle + \lambda'' - \lambda' \leq 0$ and $\langle x''_i, b' - b'' \rangle + \lambda'' - \lambda' \geq 0$. Obviously H' and H'' cannot be parallel, hence $b' \neq b''$. The convexity of A_i implies that the hyperplane $H = \{x \in \mathbf{R}^n : \langle x, b' - b'' \rangle + \lambda'' - \lambda' = 0\}$ intersects all sets A_i , $i \in I$. Using a similar argument we obtain that H intersects all sets A_j , $j \in \bar{I}$. Therefore H intersects each member of the family $\{A_1, A_2, \dots, A_{n+1}\}$ which is in general position. The contradiction obtained completes the proof.

(ii) From (i) we deduce that there exist exactly $2^n - 1$ hyperplanes which satisfy (11) and separate the members of the family $\{A_1, A_2, \dots, A_{n+1}\}$.

$N(\alpha_0) = 2^n - 1$ is the assertion (ii) in Theorem 1. If $\alpha \in S_n^+ \setminus \{\alpha_0\}$, arguing as above, Lemma 3 yields a unique hyperplane which leaves all sets A_i on the same side and which satisfies (11). □

Let $\{A_1, A_2, \dots, A_{n+1}\}$ be a family of compact convexly connected sets in general position in R^n . For each proper subset I of $\{1, 2, \dots, n + 1\}$ let $\mathcal{H}(I)$ denote the set of hyperplanes which separate the sets $\cup\{A_i : i \in I\}$ and $\cup\{A_j : j \in \bar{I}\}$. To each hyperplane $H \in \mathcal{H}(I)$ there corresponds a unique point $(b^H, \lambda^H) = (b_1^H, b_2^H, \dots, b_n^H, \lambda^H) \in S_n$ such that $H = \{x \in \mathbf{R}^n : \langle x, b^H \rangle = \lambda^H\}$ and $\cup\{A_i : i \in I\} \subset H^\geq$. This correspondence permits to identify $\mathcal{H}(I)$ with a subset of S_n , namely $\{(b^H, \lambda^H) : H \in \mathcal{H}(I)\}$.

The following known results are needed in the proof of Theorem 7.

Lemma 5 [7, Theorem 1]. *If M is a compact convex set in \mathbf{R}^n , then the function $h : \mathbf{R}^n \rightarrow \mathbf{R}$ defined by $h(b) = \max\{\langle x, b \rangle : x \in M\}$ is continuous.*

Lemma 6 [3, p.207, Lemma 3]. *Let X and Y be topological spaces, X compact and Y separated. If $f : X \rightarrow Y$ is a continuous bijection, then f is a homeomorphism.*

Theorem 7. Let $\{A_1, A_2, \dots, A_{n+1}\}$ be a family of compact convexly connected sets in general position in \mathbf{R}^n . Then for every proper subset I of $\{1, 2, \dots, n+1\}$ the sets $\mathcal{H}(I)$ and S_n^+ are homeomorphic.

PROOF: Let I be a proper subset of $\{1, 2, \dots, n+1\}$ arbitrarily fixed. Define $f : \mathcal{H}(I) \rightarrow S_n^+$ by $f(H) = \frac{1}{\|d_H\|} d_H$, where $d_H = (d(A_1, H), d(A_2, H), \dots, d(A_{n+1}, H))$. By Theorem 4, f is a bijection. By Lemma 5, each component of f is continuous, hence f is continuous too. Then, taking into account the quoted identification, $\mathcal{H}(I) = f^{-1}(S_n^+)$ is a closed subset of the compact set S_n . So $\mathcal{H}(I)$ is compact and the assertion of Theorem 7 follows now from Lemma 6. \square

Remark. Theorems 4 and 7 can be reformulated obtaining analogous informations about the hyperplanes which strictly separate the members of the family $\{A_1, A_2, \dots, A_{n+1}\}$. For instance we have:

Let $\{A_1, A_2, \dots, A_{n+1}\}$ be a family of compact convexly connected sets in general position in \mathbf{R}^n . Then for each proper subset I of $\{1, 2, \dots, n+1\}$ the set of hyperplanes strictly separating $\cup\{A_i : i \in I\}$ and $\cup\{A_j : j \in \bar{I}\}$ is homeomorphic to $\{(\alpha_1, \alpha_2, \dots, \alpha_{n+1}) \in S_n : \alpha_i > 0, 1 \leq i \leq n+1\}$.

REFERENCES

- [1] Balaj M., *(n+1)-families of sets in general position*, Beitrage zur Algebra und Geometrie **37** (1996), 67–74.
- [2] Fan K., *Fixed-point and minimax theorems in locally convex topological linear spaces*, Proc. Nat. Acad. Sci. U.S.A. **38** (1952), 121–126.
- [3] Gaal S.A., *Point Set Topology*, Academic Press, New York and London, 1964.
- [4] Glicksberg I.L., *A further generalization of the Kakutani fixed point theorem, with application to Nash equilibrium points*, Proc. Amer. Math. Soc. **3** (1952), 170–174.
- [5] Hanner O., Radström H., *A generalization of a theorem of Fenchel*, Proc. Amer. Math. Soc. **2** (1951), 589–593.
- [6] Singer I., *Best Approximation in Normed Linear Spaces by Elements of Linear Subspaces* (in Romanian), Edit. Academiei Române, București, 1967.
- [7] Valentine F.A., *The dual cone and Helly type theorems*, in: Convexity, V.L. Klee ed., Proc. Sympos. Pure Math. **7**, Amer. Math. Soc., 1963, pp. 473–493.
- [8] Valentine F.A., *Konvexe Mengen*, Manheim, 1968.

DEPARTMENT OF MATHEMATICS, ORADEA UNIVERSITY, STR. ARMATEI ROMÂNE NR. 5,
3700 ORADEA, ROMANIA

E-mail: balmir@lego.soroscj.ro

(Received December 4, 1996, revised June 17, 1997)