

Marek Wójtowicz

An elementary proof of a theorem on sublattices of finite codimension

Commentationes Mathematicae Universitatis Carolinae, Vol. 39 (1998), No. 1, 99--100

Persistent URL: <http://dml.cz/dmlcz/118989>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

An elementary proof of a theorem on sublattices of finite codimension

MAREK WÓJTOWICZ

Abstract. This paper presents an elementary proof and a generalization of a theorem due to Abramovich and Lipecki, concerning the nonexistence of closed linear sublattices of finite codimension in nonatomic locally solid linear lattices with the Lebesgue property.

Keywords: linear lattice, Lebesgue property, lattice homomorphism

Classification: 46A40, 47B65

In 1990 Y.A. Abramovich and Z. Lipecki proved, by means of Boolean algebra techniques and Liapunov's convexity theorem, the following result ([1, Theorem 2]; cf. [3, Example 27.8]):

Let $X = (X, \tau)$ be a Hausdorff locally solid linear lattice such that:

- (i) *X is nonatomic and Dedekind complete,*
- (ii) *X has the Lebesgue property.*

Then X contains no proper closed sublattices of finite codimension.

(By a sublattice of a linear lattice we always mean a *linear* sublattice. X has the Lebesgue property (or, τ is a Lebesgue topology on X) provided that for every MS-sequence (x_α) in X with $x_\alpha \downarrow 0$ we have $x_\alpha \rightarrow 0(\tau)$. For other undefined notions and basic results concerning linear lattices (= Riesz spaces) in this paper we refer the reader to the monographs [2], [3]).

Here we give an elementary and short proof of a more general result, namely, we show that the two assumptions in the above theorem, i.e. that τ is Hausdorff and X is Dedekind complete are superfluous. It should be noted that every infinite dimensional linear lattice possesses sublattices of arbitrary finite codimension ([1, Theorem 3]).

Theorem. *Let X be a nonatomic linear lattice, and let Y be a sublattice of X with $\dim X/Y < \infty$. Then Y is order dense in X .*

If, additionally, τ is a Lebesgue topology on X , then Y is τ -dense in X .

In particular, the topological dual X' of any nonatomic locally solid linear lattice (X, τ) with the Lebesgue property is nonatomic (equivalently, X has no nontrivial continuous Riesz homomorphisms $X \rightarrow \mathbf{R}$).

Order denseness of Y in X is understood in the sense of ([2, Definition 1.9]), i.e. that $Y_+ \setminus \{0\}$ is cofinal in $X_+ \setminus \{0\}$.

PROOF: Let Q denote the quotient map $X \rightarrow X/Y$. Since X is nonatomic, every principal ideal $A_e = \{x \in X : |x| \leq \lambda e \text{ for some } \lambda \geq 0\}$, $e \in X^+$, is of infinite dimension. If Y were not order dense in X , then $A_e \cap Y = \{0\}$ for some $e \in X^+$, and hence Q restricted to A_e would be a linear isomorphism; thus $\dim A_e \leq \dim Q(X) < \infty$, a contradiction. This proves the first part of the theorem, and since for Lebesgue topologies order denseness implies topological denseness, the second part also follows; the particular case is implied by ([2, Theorem 3.13]; [3, Theorem 18.3 (iii)]). \square

Examples. 1. Let K be a topological Hausdorff space. The lattice $C(K)$ is nonatomic whenever K has no isolated points, thus every such lattice has the property described in the first part of the Theorem.

2. Let S_p denote the nonatomic sublattice, consisting of all step functions, of the (nonatomic) lattice $L_p = L_p(0, 1)$, $0 < p < \infty$. It is easily seen that S_p endowed with the p -norm topology has the Lebesgue property, and hence S_p possesses the property described in the second part of the Theorem without being even σ -Dedekind complete (compare with (i) above).

3. This example seems to be known; we include it for completeness of the paper. Let $1 < p < \infty$. Since every continuous linear functional f on the lattice L_p is order continuous ([2, Theorems 9.1, 22.1 and 22.4]), any family of seminorms (q_f) of the form $q_f(x) = |f|(|x|)$, $x \in L_p$, determines a Lebesgue topology on L_p ([2, p. 40]). This topology is Hausdorff whenever the family (q_f) is total.

REFERENCES

- [1] Abramovich Y.A., Lipceki Z., *On ideals and sublattices in linear lattices and F -lattices*, Math. Proc. Cambridge Phil. Soc. **108** (1990), 79–87.
- [2] Aliprantis C.D., Burkinshaw O., *Locally Solid Riesz Spaces*, Academic Press, 1978.
- [3] Luxemburg W.A.J., Zaanen A.C., *Riesz Spaces I*, North-Holland, 1971.

INSTITUTE OF MATHEMATICS, PEDAGOGICAL UNIVERSITY, PL. SŁOWIAŃSKI 9,
65–069 ZIELONA GÓRA, POLAND

(Received July 1, 1997, revised November 4, 1997)