

Jiří Tůma

Representing lattices by homotopy groups of graphs

Commentationes Mathematicae Universitatis Carolinae, Vol. 40 (1999), No. 2, 215--221

Persistent URL: <http://dml.cz/dmlcz/119077>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1999

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Representing lattices by homotopy groups of graphs

JIRÍ TŮMA

Abstract. In this paper we represent every lattice by subgroups of free groups using the concept of the homotopy group of a graph.

Keywords: subgroup lattice, lattice embedding, free group, homotopy group

Classification: 06B15, 20E05, 20E15

In this paper we present a method how to represent a given lattice L as a sublattice of the subgroup lattice of a free group. The method is based on the idea of the homotopy group of a graph. Our construction is such that if the lattice L is finite then the free group and all its subgroups representing the elements of L are finitely generated. The first part of the proof is formulated in a more general way to enable further possible modifications of the proof replacing the free group by a finite group \mathbf{G} .

The first proof that every lattice can be embedded into the subgroup lattice of a group was given in [Wh]. The proof was later simplified by the author of the present note in [Tů2] using the solution of the word problem for HNN-extensions.

Let Ω be a set. By a *twist* on Ω we mean a bijection $t : A \rightarrow B$ between two subsets A, B of Ω . The set A is called the *domain* of t and denoted by $\text{Dom}(t)$, while B is called the *range* of t and denoted by $\text{Rng}(t)$. If $t : A \rightarrow B$ is a twist on Ω , then the inverse mapping $t^{-1} : B \rightarrow A$ is also a twist on Ω and called the *inverse* of t . The value of a twist t at a point $a \in \text{Dom}(t)$ will be written as at .

By a *twisting structure* on Ω we mean a set $\mathcal{T} = \{t_i : i \in I\}$ of twists on Ω such that with every $t \in \mathcal{T}$ the inverse t^{-1} of t is also contained in \mathcal{T} .

The Cayley graph $\mathbf{G}(\mathcal{T})$ of a twisting structure \mathcal{T} is defined as follows. The vertex set of $\mathbf{G}(\mathcal{T})$ is Ω . The edge set of $\mathbf{G}(\mathcal{T})$ is the set $E = \{(a, t) : a \in \text{Dom}(t), t \in \mathcal{T}\}$. If $e = (a, t) \in E$, then a is the initial vertex $\alpha(e)$ of e and at is the terminal vertex $\omega(e)$ of e . Since \mathcal{T} contains with every twist t also the inverse t^{-1} of t , with every edge $e = (a, t) \in E$ there is also the edge $(at, t^{-1}) \in E$. The edge (at, t^{-1}) is called the *inverse* of e and denoted by e^{-1} . It is obvious that $\alpha(e^{-1}) = \omega(e)$ and $\omega(e^{-1}) = \alpha(e)$, thus the graph $\mathbf{G}(\mathcal{T})$ is a symmetric graph possibly with loops and parallel edges. We further define the value $\nu(e)$ of an edge $e = (a, t)$ as the twist t . Thus the Cayley graph $\mathbf{G}(\mathcal{T}) = (\Omega, E, \alpha, \omega, \nu)$ of a twisting structure \mathcal{T} is a symmetric graph with edges valued by elements of \mathcal{T} . It

is the union of the graphs of all partial bijections $t \in \mathcal{T}$. Note also that for every $t \in \mathcal{T}$ and $a \in \Omega$ there is at most one edge of $\mathbf{G}(\mathcal{T})$ with initial vertex a and value $\nu(e) = t$.

By a *congruence* of a twisting structure \mathcal{T} on Ω we mean an equivalence relation π on Ω satisfying the following condition:

$$(*) \quad \text{whenever } (a, b) \in \pi, t \in \mathcal{T} \text{ and } a, b \in \text{Dom}(t), \text{ then also } (at, bt) \in \pi.$$

Thus a congruence of \mathcal{T} is a congruence of the partial algebra (Ω, \mathcal{T}) , where each $t \in \mathcal{T}$ is considered to be a partial unary operation on Ω . Obviously the set $\mathbf{C}(\mathcal{T})$ of all congruences of the twisting structure \mathcal{T} on Ω is closed under arbitrary intersections and contains the least and the greatest equivalence relations on Ω . Thus $\mathbf{C}(\mathcal{T})$ when ordered by inclusion is a complete lattice. It is called the *congruence lattice* of \mathcal{T} . The meet $\pi \wedge \rho$ of two congruences π, ρ of \mathcal{T} is their set-theoretical intersection $\pi \cap \rho$, while their join $\pi \vee \rho$ in $\mathbf{C}(\mathcal{T})$ is the least equivalence relation on Ω containing the set-theoretical union $\pi \cup \rho$ and satisfying the condition (*).

The following simple representation result was proved in [Tů1].

Theorem 1. *Every lattice L can be represented as a sublattice of $\mathbf{C}(\mathcal{T})$ for some twisting structure \mathcal{T} . □*

For the sake of completeness we present the construction. Given a lattice L , we may assume that it has a least element 0. For any two non-zero elements $a < b$ of L we define a twist $t_{a,b}$ on L with $\text{Dom}(t_{ab}) = \text{Rng}(t_{ab}) = \{0, a, b\}$ and such that

$$bt_{ab} = b, at_{ab} = 0, 0t_{ab} = a.$$

Moreover, if $a, b \in L$ are two non-comparable and non-zero elements, then we define a twist t_{ab} on L such that $\text{Dom}(t_{ab}) = \text{Rng}(t_{ab}) = \{a, b, a \vee b\}$ and

$$at_{ab} = a, bt_{ab} = a \vee b, (a \vee b)t_{ab} = b.$$

Let the twisting structure \mathcal{T} on L consist of all twists of the form t_{ab} , where $0 \neq a, b \in L$. Then the mapping assigning to every $x \in L$ the partition of L with one block the interval $[0, x]$ and the other blocks singletons is an embedding of L into $\mathbf{C}(\mathcal{T})$.

The main purpose of this note is to investigate a canonical mapping Φ from $\mathbf{C}(\mathcal{T})$ into the subgroup lattice of the free group $\mathbf{F}(\mathcal{T})$. Here $\mathbf{F}(\mathcal{T})$ denotes the free group generated by the set \mathcal{T} of free generators. Under sufficiently general conditions on \mathcal{T} we can prove that the canonical mapping Φ is an embedding of any member from a large class of sublattices of $\mathbf{C}(\mathcal{T})$. To this end we recall some basic ideas from combinatorial group theory related to homotopy groups of graphs. By a graph we mean a quadruple (V, E, α, ω) , where V, E are non-empty sets and $\alpha, \omega : E \rightarrow V$ are two incidence functions, $\alpha(e)$ is called the initial vertex of an edge $e \in E$ and $\omega(e)$ is the terminal vertex of e . By a path in the

graph (V, E, α, ω) we mean a sequence $p = e_1e_2 \cdots e_k$ of edges of E such that $\omega(e_i) = \alpha(e_{i+1})$ for every $i = 1, 2, \dots, k - 1$. A path $p = e_1e_2 \cdots e_k$ is called a loop if $\omega(e_k) = \alpha(e_1)$, and a loop $p = e_1e_2 \cdots e_k$ is called a loop at a vertex $v \in V$ if $\alpha(e_1) = v$.

We will generalize these concepts to the Cayley graph of a twisting structure \mathcal{T} . Let π be a congruence of \mathcal{T} . A sequence $p = e_1e_2 \cdots e_k$ of edges of the Cayley graph $\mathbf{C}(\mathcal{T})$ of \mathcal{T} is called a π -path if $(\omega(e_i), \alpha(e_{i+1})) \in \pi$ for every $i = 1, 2, \dots, k - 1$. A path $p = e_1e_2 \cdots e_k$ is called a π -loop if $(\alpha(e_1), \omega(e_k)) \in \pi$ and it is called a π -loop at a vertex $v \in \Omega$ if moreover $(v, \alpha(e_1)) \in \pi$. If $(\omega(e_i), \alpha(e_{i+1})) \in \tau$ for some relation τ on Ω , then we say that $(\omega(e_i), \alpha(e_{i+1}))$ is a τ -jump.

Suppose moreover that $\mathcal{T} \subset \mathbf{G}$, where \mathbf{G} is a group. Then we can assign to every π -path $p = e_1e_2 \cdots e_k$ its \mathbf{G} -value

$$\nu(p) = \nu(e_1)\nu(e_2) \cdots \nu(e_k) \in \mathbf{G}.$$

The set of \mathbf{G} -values of all π -loops at a vertex $v \in \Omega$ in the Cayley graph $\mathbf{C}(\mathcal{T})$ of a twisting structure \mathcal{T} is obviously a subgroup of \mathbf{G} . Indeed, $1 \in \mathbf{G}$ is the value of the empty π -path, if $g = \nu(p)$ for a π -loop $p = e_1e_2 \cdots e_k$ at v , then g^{-1} is the value of the inverse path $p^{-1} = e_k^{-1} \cdots e_1^{-1}$. And if $g = \nu(p)$, $h = \nu(q)$, then $gh = \nu(pq)$. Thus we can define a mapping

$$\Phi_{\mathbf{G}} : \mathbf{C}(\mathcal{T}) \rightarrow \text{Sub}(\mathbf{G})$$

from the congruence lattice of \mathcal{T} into the subgroup lattice of \mathbf{G} by

$$\Phi_{\mathbf{G}}(\pi) = \{\nu(p) : p \text{ is a } \pi\text{-loop at } v\}.$$

If $\pi \subseteq \rho$ are two congruences of \mathcal{T} , then obviously any π -loop at v is also a ρ -loop at v , thus we get the following simple lemma.

Lemma 2. *The mapping $\Phi_{\mathbf{G}}$ is order-preserving.* □

We say that a twisting structure \mathcal{T} is connected if its Cayley graph $\mathbf{G}(\mathcal{T})$ is connected. For connected twisting structures we have the following result.

Theorem 3. *If \mathcal{T} is a connected twisting structure, then the mapping $\Phi_{\mathbf{G}}$ is join-preserving.*

PROOF: First of all we describe the join $\pi \vee \rho$ of two congruences $\pi, \rho \in \mathbf{C}(\mathcal{T})$. Set $\sigma_0 = \pi \cup \rho$. If σ_{2i} is already defined for a natural number i , we define

$$\sigma_{2i+1} = \sigma_{2i} \cup \{(at, bt) : t \in \mathcal{T}, a, b \in \text{Dom}(t), (a, b) \in \sigma_{2i}\},$$

and

$$\sigma_{2i+2} \text{ is the transitive closure of } \sigma_{2i+1}.$$

Obviously, any congruence σ of \mathcal{T} containing both π and ρ must contain also σ_n for any natural number n . On the other hand,

$$\sigma = \bigcup_n \sigma_n$$

is an equivalence relation satisfying the condition (*), hence a congruence of \mathcal{T} . Thus $\sigma = \pi \vee \rho$ in $\mathbf{C}(\mathcal{T})$.

Since $\Phi_{\mathbf{G}}$ is order-preserving, we get that

$$\Phi_{\mathbf{G}}(\pi) \vee \Phi_{\mathbf{G}}(\rho) \subseteq \Phi_{\mathbf{G}}(\pi \vee \rho)$$

for any two congruences π, ρ of \mathcal{T} . To prove the opposite inclusion we have to show that the value $\nu(p)$ of any $(\pi \vee \rho)$ -path p at v is contained in the subgroup of \mathbf{G} generated by $\Phi_{\mathbf{G}}(\pi) \cup \Phi_{\mathbf{G}}(\rho)$.

So let $p = e_1 e_2 \cdots e_k$ be an arbitrary $(\pi \vee \rho)$ -loop at v . Thus $(\omega(e_i), \alpha(e_{i+1})) \in \pi \vee \rho$ for any $i = 1, 2, \dots, k - 1$ as well as $(v, \alpha(e_1)), (\omega(e_k), v) \in \pi \vee \rho$. Since $\pi \vee \rho = \bigcup_n \sigma_n$, there exists a natural number m such that

$$(v, \alpha(e_1)), (\omega(e_k), v) \in \sigma_m, (\omega(e_i), \alpha(e_{i+1})) \in \sigma_m$$

for every $i = 1, 2, \dots, k - 1$. Let us call such a $(\pi \vee \rho)$ -loop $p = e_1 e_2 \cdots e_k$ at v a σ_m -loop. We are going to prove that the value $\nu(p)$ of any σ_m -loop at v , $m \geq 1$, belongs to the subgroup of \mathbf{G} generated by the values of σ_{m-1} -loops at v .

If m is odd, let $i \leq k$ be such that $(\omega(e_i), \alpha(e_{i+1})) \in \sigma_m \setminus \sigma_{m-1}$. Thus there exist a twist $t \in \mathcal{T}$ and $a, b \in \sigma_{m-1}$ such that $at = \omega(e_i)$ and $bt = \alpha(e_{i+1})$. Consider the loop $p' = e_1 e_2 \cdots e_i (at, t^{-1})(b, t) e_{i+1} \cdots e_k$. Since $\omega(e_i) = \alpha(at, t^{-1})$, $(\omega(at, t^{-1}), \alpha(b, t)) = (a, b) \in \sigma_{m-1}$ and $\omega(b, t) = bt = \alpha(e_{i+1})$, p' is also a σ_m -loop at v and the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps in p' is one less than the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps in p . Similarly, if $(v, \alpha(e_1)) \in \sigma_m \setminus \sigma_{m-1}$, then again there are a twist $t \in \mathcal{T}$ and $(a, b) \in \sigma_m \setminus \sigma_{m-1}$ such that $at = v$ and $bt = \alpha(e_1)$. Again the loop $p' = (at, t^{-1})(b, t)p$ is a σ_m -loop at v (since $(v, t^{-1}) = (at, t^{-1})$) and the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps in p' is one less than the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps in p . The case $(\omega(e_k), v) \in \sigma_m \setminus \sigma_{m-1}$ is treated in exactly the same way. In all cases, $\nu(p') = \nu(p)$ and the number of $(\sigma_m \setminus \sigma_{m-1})$ -jumps in the path p' is one less than the number of $(\sigma_m \setminus \sigma_{m-1})$ -jumps in p . By a simple induction on the number of $(\sigma_m \setminus \sigma_{m-1})$ -jumps in p we prove that for every σ_m -loop p at v there exists a σ_{m-1} -loop p'' at v such that $\nu(p'') = \nu(p)$. Thus if m is odd, then the value $\nu(p)$ of any σ_m -loop p at v is equal to the value of a σ_{m-1} -loop at v .

If $m > 0$ is even and $p = e_1 e_2 \cdots e_k$ a σ_m -loop at v that is not a σ_{m-1} -loop, then either there exists some $i = 1, 2, \dots, k - 1$ such that $(\omega(e_i), \alpha(e_{i+1})) \in \sigma_m \setminus \sigma_{m-1}$ or $(v, \alpha(e_1)) \in \sigma_m \setminus \sigma_{m-1}$ or $(\omega(e_k), v) \in \sigma_m \setminus \sigma_{m-1}$. Let the first of the three possibilities occur. Since σ_m is the transitive closure of σ_{m-1} , there are elements $\omega(e_i) = a_1, a_2, \dots, a_l = \alpha(e_{i+1})$ such that $(a_j, a_{j+1}) \in \sigma_{m-1}$. Since the Cayley graph $\mathbf{G}(\mathcal{T})$ of \mathcal{T} is connected, there are paths q_i in $\mathbf{G}(\mathcal{T})$ of \mathcal{T} from v to a_i . Then

$p' = e_1 e_2 \cdots e_i q_1^{-1} q_1 q_2^{-1} \cdots q_{l-1} q_l^{-1} q_l e_{i+1} \cdots e_k$ is again a σ_m -loop at v in which the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps is one less than the number of $\sigma_m \setminus \sigma_{m-1}$ -jumps in p . Moreover, $\nu(p') = \nu(p)$. The other two cases are treated in exactly the same way. Thus also in this case the value of any σ_m -loop at v coincides with the value of a σ_{m-1} -loop at v .

Hence the value $\nu(p)$ of any $(\pi \vee \rho)$ -loop at v equals the value $\nu(p')$ of a σ_0 -loop p' at v . Recall that $\sigma_0 = \pi \cup \rho$. Let $p' = f_1 f_2 \cdots f_l$. For every $i = 1, 2, \dots, l$ let q_i be a path in $\mathbf{G}(\mathcal{T})$ from v to $\omega(f_i)$. Then

$$p'' = f_1 q_1^{-1} q_1 f_2 q_2^{-1} q_2 f_3 \cdots q_{l-1}^{-1} q_{l-1} f_l$$

is also a σ_0 -loop at v . Obviously, $\nu(p'') = \nu(p)$. Finally, observe that each $q_{i-1} f_i q_i^{-1}$, $i = 2, \dots, l-1$ is either a π -loop or a ρ -loop, since $(\omega(f_{i-1}), \alpha(f_i)) \in \sigma_0 = \pi \cup \rho$. Thus $\nu(q_{i-1} f_i q_i^{-1}) \in \Phi_{\mathbf{G}}(\pi) \cup \Phi_{\mathbf{G}}(\rho)$.

Similarly, we prove that also $\nu(f_1) q_1^{-1} \in \Phi_{\mathbf{G}}(\pi) \cup \Phi_{\mathbf{G}}(\rho)$ and $\nu(q_{l-1} f_l) \in \Phi_{\mathbf{G}}(\pi) \cup \Phi_{\mathbf{G}}(\rho)$. Thus $\nu(p'') \in \Phi_{\mathbf{G}}(\pi) \vee \Phi_{\mathbf{G}}(\rho)$. □

In the rest of the paper we restrict ourselves to the case that \mathbf{G} is the free group \mathbf{F} freely generated by \mathcal{T} . Let L be any non-empty collection of partitions on the set Ω closed under finite meets. We say that a twisting structure \mathcal{T} on Ω is *balanced* with respect to L if for every twist $t \in \mathcal{T}$, an element $a \in \Omega$ and any two $x, y \in \text{Dom}(t)$, whenever $(a, x) \in \pi \in L$ and $(a, y) \in \rho \in L$, then there exists some $z \in \text{Dom}(t)$ such that $(a, z) \in \pi \wedge \rho \in L$. The following lemma from [Tũ1] gives us a way to construct balanced sets.

Lemma 4. *Let X be a set and $\Omega = \mathbf{S}_X$, the group of all permutations of X of a finite type (i.e. generated by transpositions). For a set $Y \subset X$ let S_Y be the subgroup of \mathbf{S}_X consisting of all permutations p such that $p(x) = x$ for every $x \in X \setminus Y$. Let L be the set of partitions of Ω into left cosets of subgroups S_Y , $Y \subseteq X$. Then every left coset of every S_Y , $Y \subseteq X$, is balanced with respect to L .* □

By modifying Example 2.5. and Proposition 2.8. of [Tũ1] we get the following lemma.

Lemma 5. *For every lattice \mathbf{L} there exist a set Ω , a twisting structure \mathcal{T} on Ω with finite domains and a lattice embedding $\phi : \mathbf{L} \rightarrow \mathbf{C}(\mathcal{T})$ such that the twisting structure \mathcal{T} is balanced with respect to the lattice $L = \text{Im}(\phi)$.* □

No we are ready to prove the following counterpart to Theorem 3.

Theorem 6. *Let \mathcal{T} be a twisting structure on a set Ω and $L \subset \mathbf{C}(\mathcal{T})$ a sublattice of $\mathbf{C}(\mathcal{T})$. Suppose moreover that the domains of the elements of \mathcal{T} are finite and that \mathcal{T} is balanced with respect to L . Then the restriction of the canonical mapping $\Phi_{\mathbf{F}}$ to the lattice L is meet-preserving.*

PROOF: Let $\pi \in \mathbf{C}(\mathcal{T})$. First of all we prove that for any π -loop $p = e_1 e_2 \cdots e_k$ in $\mathbf{G}(\mathcal{T})$ at v such that $\nu(e_1) \nu(e_2) \cdots \nu(e_k)$ is not a reduced word in \mathbf{F} there

exists a subpath $p' = e_{i_1} \cdots e_{i_l}$ of p that is also a π -loop at v and the word $\nu(e_{i_1}) \cdots \nu(e_{i_l})$ is reduced. Indeed, if $\nu(e_1)\nu(e_2) \cdots \nu(e_k)$ is not reduced, then there is some $i = 1, 2, \dots, k - 1$ such that $\nu(e_i) = t = \nu(e_{i+1})^{-1}$. Thus there are some $a, b \in \text{Dom}(t)$ such that $e_i = (a, t)$ and $e_{i+1} = (bt, t^{-1})$. Moreover, since p is a π -path, we have $(\omega(e_i), \alpha(e_{i+1})) = (at, bt) \in \pi$. Since $\pi \in \mathbf{C}(\mathcal{T})$, we have also $(att^{-1}, btt^{-1}) = (a, b) \in \pi$. But we have also $(\omega(e_{i-1}), \alpha(e_i)) = (\omega(e_{i-1}), a) \in \pi$ and $(\omega(e_{i+1}), \alpha(e_{i+2})) = (b, \alpha(e_{i+2})) \in \pi$, we get $(\omega(e_{i-1}), \alpha(e_{i+2})) \in \pi$. Thus we can delete from p the edges e_i, e_{i+1} and the remaining path p' is still a π -loop at v . In this way we can subsequently delete from p pairs of subsequent edges with mutually inverse values to get a π -loop p' with required properties. Let us call such a π -path a *reduced* π -path.

Take now arbitrary congruences $\pi, \rho \in L$. Since $\Phi_{\mathbf{F}}$ is order-preserving, we have

$$\Phi_{\mathbf{F}}(\pi) \cap \Phi_{\mathbf{F}}(\rho) \supseteq \Phi_{\mathbf{F}}(\pi \cap \rho).$$

To prove the opposite inclusion take any reduced word $w = t_1 \cdots t_k \in \Phi_{\mathbf{F}}(\pi) \cap \Phi_{\mathbf{F}}(\rho)$. Then by the previous paragraph there is a reduced π -loop $p = e_1 e_2 \cdots e_k$ at v such that $\nu(p) = w$. Hence $\nu(e_i) = t_i$ for every $i = 1, 2, \dots, k$. Similarly, there is a reduced ρ -loop $q = f_1 f_2 \cdots f_l$ at v such that $\nu(f_i) = t_i$ for every $i = 1, 2, \dots, k$. Thus in particular, $(v, \alpha(e_1)) \in \pi$ and $(v, \alpha(f_1)) \in \rho$. Let us denote $\alpha(e_1) = a$ and $\alpha(f_1) = b$. Thus $a, b \in \text{Dom}(t_1)$. Since $\text{Dom}(t_1)$ is balanced with respect to L , there exists some $c \in \text{Dom}(t_1)$ such that $(v, c) \in \pi \cap \rho$. Hence also $(a, c) \in \pi$ and $(b, c) \in \rho$. Since both π and ρ are congruences of \mathcal{T} , we get also $(at_1, ct_1) \in \pi$ and $(bt_1, ct_1) \in \rho$. Moreover, $(at_1, \alpha(e_2)) = (\omega(e_1), \alpha(e_2)) \in \pi$ and $(bt_1, \alpha(f_2)) = (\omega(f_1), \alpha(f_2)) \in \rho$, we get $(ct_1, \alpha(e_2)) \in \pi$ and $(ct, \alpha(f_2)) \in \rho$. Denote by g_1 the edge (c, ct_1) . Thus $g_1 e_2 \cdots e_k$ is another π -loop at v and $g_1 f_2 \cdots f_k$ is another ρ -loop at v . Moreover, $(v, \alpha(g_1)) = (v, c) \in \pi \cap \rho$, and $\nu(g_1) = t_1$.

By repeating the same procedure with ct_1, e_2 and f_2 in place of v, e_1 and f_1 , we get another edge g_2 that can replace e_2 in p and f_2 in q and satisfies $\nu(g_2) = \nu(e_2) = \nu(f_2)$ and $(\omega(g_1), \alpha(g_2)) \in \pi \cap \rho$. After k steps we construct a $(\pi \cap \rho)$ -loop $r = g_1 g_2 \cdots g_k$ at v with $\nu(r) = w$. Hence $w \in \Phi_{\mathbf{F}}(\pi \cap \rho)$. \square

Finally, connectedness of \mathcal{T} also implies injectivity of $\Phi_{\mathbf{F}}$. Since the method of the proof will be also used in the proof of Theorem 8, we introduce some definitions here. If π is a congruence of \mathcal{T} , we define the quotient \mathcal{T}/π of \mathcal{T} as follows. The twisting structure \mathcal{T}/π will be defined on the set Ω/π of blocks of π . For any twist $t \in \mathcal{T}$ we define another twist t_π on Ω/π . The domain $\text{Dom}(t_\pi)$ consists of all blocks of π intersecting the domain $\text{Dom}(t)$. If $a \in \text{Dom}(t)$, then we define $[a]t_\pi = [at]$, where $[x]$ denotes the block of π containing x . The definition of t_π is correct since π is a congruence of \mathcal{T} . Hence t_π is also a bijection between two subsets of Ω/π and $\{t_\pi : t \in \mathcal{T}\}$ is a twisting structure on Ω/π . Thus \mathcal{T}/π is simply the quotient of the partial unary algebra (Ω, \mathcal{T}) by the congruence π .

It is also useful to mention that the graph of \mathcal{T}/π is naturally isomorphic to a quotient of the graph of \mathcal{T} . The vertices of $\mathbf{G}(\mathcal{T}/\pi)$ are blocks of the partition π on Ω . Whenever $a, b \in \text{Dom}(t)$ are such that $(a, b) \in \pi$, then the two edges (a, t)

and (b, t) of $\mathbf{G}(\mathcal{T})$ are identified into a single edge $([a], [a]t_\pi)$ of $\mathbf{G}(\mathcal{T}/\pi)$. If we assign to each edge $([a], [a]t_\pi)$ of $\mathbf{G}(\mathcal{T}/\pi)$ the value $t \in \mathbf{F}$, then we see that the values of π -loops at v in the graph $\mathbf{G}(\mathcal{T})$ are in one-to-one correspondence with the values of ordinary loops at $[v]$ in the graph of $\mathbf{G}(\mathcal{T}/\pi)$.

Lemma 7. *If \mathcal{T} is a connected twisting structure on Ω and $\pi < \rho$ two congruences of \mathcal{T} , then $\Phi_{\mathbf{F}}(\pi) \neq \Phi_{\mathbf{F}}(\rho)$.*

PROOF: Let $(a, b) \in \rho \setminus \pi$. Since \mathcal{T} is connected, there exist a reduced path $p = e_1 e_2 \cdots e_k$ in $\mathbf{G}(\mathcal{T})$ from v to a and a reduced path $q = f_1 f_2 \cdots f_l$ in $\mathbf{G}(\mathcal{T})$ from v to b . Then pq^{-1} is a ρ -loop in $\mathbf{G}(\mathcal{T})$ at v but it is not a π -loop at v . If pq^{-1} is not reduced, then there exists a twist $t \in \mathcal{T}$ such that $e_k = (at^{-1}, t)$ and $f_l = (bt^{-1}, t)$. Then also $(at^{-1}, bt^{-1}) \in \rho \setminus \pi$. So we can replace p by $p' = e_1 e_2 \cdots e_{k-1}$ and q by $q' = f_1 f_2 \cdots f_{l-1}$ to get a shorter ρ -loop $p'q'^{-1}$ at v that is not a π -loop at v . Hence we may assume that the ρ -loop $r = pq^{-1}$ at v is already reduced and it is not a π -loop. But then $\nu(r) \in \Phi_{\mathbf{F}}(\rho) \setminus \Phi_{\mathbf{F}}(\pi)$. \square

Putting together previous results we get the following theorem.

Theorem 8. *Every lattice L can be embedded into the subgroup lattice of a free group \mathbf{F} . If the lattice L is finite, then the group \mathbf{F} and all the subgroups of \mathbf{F} representing elements of L can be taken finitely generated.*

PROOF: It remains to prove the second assertion. However, if the lattice L is finite, then the twisting structure \mathcal{T} of Lemma 4 can be taken finite by [Tû1]. But then the group $\Phi_{\mathbf{F}}(\pi)$ is isomorphic to the homotopy group of the graph $\mathbf{G}(\mathcal{T}/\pi)$, by the remarks preceding Lemma 7. Since the graph $\mathbf{G}(\mathcal{T}/\pi)$ is finite, its homotopy group is a finitely free group. \square

REFERENCES

- [Tû1] Tûma J. On infinite partition representations and their finite quotients, Czech. Math. J. **45** (1995), 21–38.
- [Tû2] Tûma J., *A new proof of Whitman's embedding theorem*, J. Algebra **173** (1995), 459–462.
- [W] Whitman P., *Lattices, equivalence relations, and subgroups*, Bull. Amer. Math. Soc. **52** (1946), 507–522.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail: tuma@karlin.mff.cuni.cz

(Received January 23, 1998, revised January 3, 1999)