

Alessandro Fedeli; Attilio Le Donne

An independency result in connectification theory

Commentationes Mathematicae Universitatis Carolinae, Vol. 40 (1999), No. 2, 331--334

Persistent URL: <http://dml.cz/dmlcz/119089>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1999

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

An independency result in connectification theory

ALESSANDRO FEDELI, ATTILIO LE DONNE

Abstract. A space is called connectifiable if it can be densely embedded in a connected Hausdorff space.

Let ψ be the following statement: “a perfect T_3 -space X with no more than 2^c clopen subsets is connectifiable if and only if no proper nonempty clopen subset of X is feebly compact”.

In this note we show that neither ψ nor $\neg\psi$ is provable in ZFC.

Keywords: connectifiable, perfect, feebly compact

Classification: 54D25, 54C25, 03E35

The problem of finding those spaces which can be densely embedded in a connected Hausdorff space has been extensively studied in the last years and many results have been obtained (see, e.g., [1], [2], [6], [10] and [13]).

Despite all the efforts, a characterization of connectifiable spaces is still unknown.

In this note we present a characterization of connectifiable perfect T_3 -spaces with no more than 2^c clopen subsets, which can be neither proved nor disproved in ZFC.

We recall that a space X is called:

- (i) perfect if every closed subset of X is a G_δ -set;
- (ii) H-closed if every open cover of X has a finite subfamily whose union is dense, or equivalently, X is a closed subspace of every Hausdorff space in which it is contained;
- (iii) feebly compact if every countable open cover of X has a finite subfamily whose union is dense.

As usual, p will stand for the smallest cardinality of a maximal subfamily of $[\omega]^\omega$ with the strong finite intersection property (see, e.g., [4] and [12]).

Regarding connectifiability observe that

- (1) A connectifiable space contains no proper nonempty open H-closed subset ([13]).
- (2) Let X be a Hausdorff space with no more than 2^c clopen subsets. If every proper nonempty clopen subsets of X is not feebly compact, then X is connectifiable ([10]).

(3) There exists, in ZFC, a nonconnectifiable Hausdorff space of cardinality \mathfrak{c} with no proper nonempty H-closed subspace ([10]).

(4) It is consistent with ZFC that there is a nonconnectifiable normal Hausdorff space of cardinality \mathfrak{c} which has no proper nonempty H-closed subspace ([10]).

In our result we will make use of the following set-theoretic statements (which are consistent with ZFC):

(a) $p > \omega_1$;

(b) (Jensen's Combinatorial Principle \diamond) There are sets $A_\alpha \subset \alpha$ for $\alpha < \omega_1$ such that for every $A \subset \omega_1$ the set $\{\alpha < \omega_1 : A \cap \alpha = A_\alpha\}$ is stationary.

We refer the reader to [5] for topological terminology. For set-theoretic terminology see [8] and [4].

Theorem. *The following statement:*

“a perfect T_3 -space X with no more than $2^{\mathfrak{c}}$ clopen subsets X is connectifiable if and only if no proper nonempty clopen subset of X is feebly compact”

is independent of ZFC.

PROOF: First let us show that, under $p > \omega_1$, the above statement is true.

If X is not connectifiable then, by one of the above mentioned result, there is a proper nonempty feebly compact clopen subset A of X .

Now let us suppose that there is a proper nonempty feebly compact clopen subset A of X . Since X is T_3 and perfect, it follows that A is a countably compact perfect T_3 -space. So by a theorem of Weiss (here we use $p > \omega_1$) A is compact ([14], see also [12]).

Hence X is not connectifiable. Therefore the statement is consistent with ZFC.

Now let us prove the independency by showing that, under the Jensen's principle \diamond , there exists a connectifiable T_6 -space with exactly two proper nonempty clopen subsets, each of which is feebly compact.

Let S be the Ostaszewskii's space, this space is, under \diamond , an example of a noncompact countably compact perfectly normal space ([9]).

Let Z be the cone over S , i.e., let Z be the quotient of $S \times I$ obtained by identifying $S \times \{1\}$ with a point.

Now Z is noncompact ($S \times \{0\}$ is a noncompact space homeomorphic to a closed subspace of Z), countably compact and perfectly normal (Z is the continuous closed image of the countably compact perfectly normal space $S \times I$ under the natural mapping).

Now let $X = Z \oplus Z$, X is a perfectly normal space. The only proper nonempty clopen subsets of X are the two copies of Z , which are countably compact (= feebly compact) but not compact.

Since a Hausdorff space with open components is connectifiable if and only if it has no proper nonempty open H-closed subspace ([7]), it follows that X is connectifiable. \square

Example. It is worth noting that there is a ZFC example of a connectifiable perfect Hausdorff space, with no more than $2^{\mathfrak{c}}$ clopen subsets, which has proper nonempty feebly compact clopen subsets.

In fact let \mathcal{F} be the set of all free ultrafilters on ω and let Y be $\omega \cup \mathcal{F}$ endowed with the topology generated by the points of ω and all sets of the form $G \cup \{p\}$ where $G \in p \in \mathcal{F}$. Now fix $p \in \mathcal{F}$ and let X be the subspace $Y \setminus \{p\}$ of Y .

X is a Hausdorff space which is not H-closed (it is not closed in Y).

Now let us show that X is feebly compact. By a result in [3] it is enough to show that every locally finite system of pairwise disjoint nonempty open subsets of X is finite.

Suppose that $\mathcal{A} = \{A_n : n \in \omega\}$ is an infinite locally finite family of pairwise disjoint nonempty open subsets of X . Without loss of generality we may assume that, for every n , $A_n = \{\kappa_n\}$ for some $\kappa_n \in \omega$.

Let q be a free open ultrafilter on ω such that $q \neq p$ and $\{\kappa_n : n \in \omega\} \in q$. Since every neighbourhood of q in X meets infinitely many members of \mathcal{A} , we reach a contradiction. Therefore X is feebly compact.

Moreover X is perfect. In fact, every open subset A of X , is the union of the F_σ -set $\omega \cap A$ and the closed set $A \setminus \omega$ ($A \setminus \omega$ is a subset of the closed discrete subspace $X \setminus \omega$ of X).

Now let C be the cone over X and set $Z = C \oplus C$.

Z is a perfect Hausdorff space, and the only two proper nonempty clopen subsets of Z (namely the copies of X) are feebly compact.

Nonetheless Z has open components and no proper nonempty H-closed subspaces, therefore Z is connectifiable.

Remarks. (i) If L is the long line, then $X = L \oplus L$ is a ZFC example of a connectifiable hereditarily normal space of cardinality \mathfrak{c} which has proper nonempty feebly compact clopen subsets.

(ii) In [10] it is shown that, under $MA + \neg CH$, a disconnected perfectly normal space with no more than $2^{\mathfrak{c}}$ clopen subsets is connectifiable if and only if no nonempty clopen subset is relatively pseudocompact.

REFERENCES

- [1] Alas O.T., Tkačenko M.G., Tkachuk V.V., Wilson R.G., *Connectifying some spaces*, Topology Appl. **71** (1996), 203–215.
- [2] Alas O.T., Tkačenko M.G., Tkachuk V.V., Wilson R.G., *Connectedness and local connectedness of topological groups and extensions*, Comment. Math. Univ. Carolinae, to appear.
- [3] Bagley R.W., Connell E.H., McKnight J.D., Jr., *On properties characterizing pseudocompact spaces*, Proc. A.M.S. **9** (1958), 500–506.
- [4] van Douwen E.K., *The integers and topology*, Handbook of Set-theoretic Topology, (Kunen K. and Vaughan J.E., eds.) Elsevier Science Publishers B.V., North Holland, 1984, pp. 111–167.
- [5] Engelking R., *General Topology*, Sigma series in Pure Mathematics 6, Heldermann Verlag, Berlin, 1989.
- [6] Fedeli A., Le Donne A., *On locally connected connectifications*, Topology Appl. (1998).
- [7] Fedeli A., Le Donne A., *Connectifications and open components*, submitted.

- [8] Kunen K., *Set Theory*, North Holland, 1980.
- [9] Ostaszewski A.J., *On countably compact, perfectly normal spaces*, J. London Math. Soc. (2) **14** (1976), 505–516.
- [10] Porter J.R., Woods R.G., *Subspaces of connected spaces*, Topology Appl. **68** (1996), 113–131.
- [11] Swardson M.A., *Nearly realcompact spaces and T_2 -connectifiability*, Proceedings of the 8th Prague Topological Symposium, 1996, pp. 358–361.
- [12] Vaughan J.E., *Countably compact and sequentially compact spaces*, Handbook of Set-theoretic Topology, (Kunen K. and Vaughan J.E., eds.) Elsevier Science Publishers B.V., North Holland, 1984, pp. 569–602.
- [13] Watson S., Wilson R.G., *Embeddings in connected spaces*, Houston J. Math. **19** (1993), no. 3, 469–481.
- [14] Weiss W.A.R., *Countably compact spaces and Martin's axiom*, Canad. J. Math. **30** (1978), 243–249.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF L'AQUILA, 67100 L'AQUILA, ITALY

E-mail: fedeli@aquila.infn.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROME "LA SAPIENZA" 00100 ROME, ITALY

E-mail: ledonne@mat.uniroma1.it

(Received April 7, 1998)