

David Cruz-Uribe

A new proof of weighted weak-type inequalities for fractional integrals

Commentationes Mathematicae Universitatis Carolinae, Vol. 42 (2001), No. 3, 481--485

Persistent URL: <http://dml.cz/dmlcz/119262>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2001

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A new proof of weighted weak-type inequalities for fractional integrals

D. CRUZ-URIBE, SFO

Abstract. We give a new and simpler proof of a two-weight, weak (p, p) inequality for fractional integrals first proved by Cruz-Urbe and Pérez [4].

Keywords: weights, weak-type inequalities, fractional integrals

Classification: 42B20, 42B25

1. Introduction

For $0 < \alpha < n$, the fractional integral operator I_α is defined by

$$I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-\alpha}} dy.$$

In [4], Cruz-Urbe and Pérez proved a two-weight, weak-type norm inequality which answered a question posed by Sawyer and Wheeden [9].

Theorem. *Given a pair of weights (u, v) , $p, 1 < p < \infty$, and $\alpha, 0 < \alpha < n$, suppose that for some $r > 1$ and for all cubes Q ,*

$$(1.1) \quad |Q|^{\alpha/n} \left(\frac{1}{|Q|} \int_Q u^r dx \right)^{1/rp} \left(\frac{1}{|Q|} \int_Q v^{-p'/p} dx \right)^{1/p'} \leq C < \infty.$$

Then the fractional integral operator I_α satisfies the weak (p, p) inequality

$$(1.2) \quad u(\{x \in \mathbb{R}^n : |I_\alpha f(x)| > t\}) \leq \frac{C}{t^p} \int_{\mathbb{R}^n} |f|^{p'} v dx.$$

Their proof of Theorem 1.1 was fairly complex, and depended on a technical lemma resembling a good- λ inequality. The purpose of this paper is to give another, more elementary proof. It is based on three weighted norm inequalities

I want to thank the referee for detecting an error in the first version of this paper.

for the fractional integral operator and the closely related fractional maximal operator,

$$M_\alpha f(x) = \sup_{Q \ni x} \frac{|Q|^{\alpha/n}}{|Q|} \int_Q |f| dy.$$

These are stated in Section 2 and the proof of Theorem 1.1 is in Section 3.

Finally, we remark that we believe that condition (1.1) in Theorem 1.1 is not the best possible. In [4], Cruz-Uribe and Pérez proved an analogue of Theorem 1.1 for singular integrals, and in [3] they sharpened this result by replacing the local L^r norm on the left-hand side of (1.1) by the smaller Orlicz space norm $\|\cdot\|_{L(\log L)^{p-1+\delta}}$, $\delta > 0$. (A similar condition is sufficient for a strong (p, p) inequality for fractional integrals. See Pérez [7].) We conjecture that the corresponding weak (p, p) result holds for fractional integrals. For a partial result, see Cruz-Uribe and Fiorenza [2].

2. Preliminary results

The first result we need is due to Muckenhoupt and Wheeden [6, p. 262].

Theorem 2.1. *Given α , $0 < \alpha < n$, and a weight $w \in A_\infty$, there exists a constant C , depending only on α , n and the A_∞ constant of w , such that for all functions f ,*

$$\sup_{t>0} t w(\{x \in \mathbb{R}^n : |I_\alpha f(x)| > t\}) \leq C \sup_{t>0} t w(\{x \in \mathbb{R}^n : M_\alpha f(x) > t\}).$$

The second result is due to Sawyer [8, p. 285]; for a simple proof see [1].

Theorem 2.2. *Given α , $0 < \alpha < n$, and a weight w , there exists a constant C , depending only on α and n , such that for all functions f ,*

$$w(\{x \in \mathbb{R}^n : M_\alpha f(x) > t\}) \leq \frac{C}{t} \int_{\mathbb{R}^n} |f| M_\alpha w dx.$$

The third result is a special case of a theorem due to Pérez [7, p. 668].

Theorem 2.3. *Given α , $0 < \alpha < n$, $r > 1$, and a pair of weights (u, v) such that (1.1) holds, then there exists a constant s , $1 < s < \min(n/\alpha, r)$, and a constant C such that for all functions f ,*

$$(2.1) \quad \int_{\mathbb{R}^n} (M_{\alpha s} f)^{p'/s} v^{-p'/p} dx \leq C \int_{\mathbb{R}^n} |f|^{p'/s} u^{-p'/p} dx.$$

PROOF: Given s , $1 < s < \min(n/\alpha, r)$, condition (1.1) is equivalent to

$$|Q|^{s\alpha/n} \left(\frac{1}{|Q|} \int_Q [v^{-s/p}]^{p'/s} dx \right)^{s/p'} \left(\frac{1}{|Q|} \int_Q [u^{s/p}]^{rp/s} dx \right)^{s/rp} \leq C.$$

Pérez showed that this implies (2.1), provided that $(rp/s)' < p'/s$. This is true for $s = 1$, so by continuity it is true for $s > 1$ sufficiently small. \square

3. Proof of Theorem 1.1

The proof requires one lemma.

Lemma 3.1. *Given α , $0 < \alpha < n$, and s , $1 < s < n/\alpha$, then for all non-negative, locally integrable functions g , $M_\alpha(M(g^s)^{1/s})(x) \leq CM_{\alpha s}(g^s)(x)^{1/s}$, where M is the Hardy-Littlewood maximal operator.*

PROOF: Our proof is modeled on the proof of a similar result in García-Cuerva and Rubio de Francia [5, p.158]. It will suffice to show that there exists a constant C such that for any x and any cube Q containing x ,

$$\frac{|Q|^{\alpha/n}}{|Q|} \int_Q M(g^s)(y)^{1/s} dy \leq CM_{\alpha s}(g^s)(x)^{1/s}.$$

Let $g = g_1 + g_2$, where $g_1 = g\chi_{3Q}$. Then $M(g^s)(x)^{1/s} \leq M(g_1^s)(x)^{1/s} + M(g_2^s)(x)^{1/s}$. Since M is weak $(1, 1)$, by Kolmogorov's inequality (cf. [5, p. 485]),

$$\begin{aligned} \frac{|Q|^{\alpha/n}}{|Q|} \int_Q M(g_1^s)(y)^{1/s} dy &\leq C|Q|^{\alpha/n} \left(\frac{1}{|Q|} \int_{\mathbb{R}^n} g_1^s dy \right)^{1/s} \\ &\leq C|Q|^{\alpha/n} \left(\frac{1}{|3Q|} \int_{3Q} g^s dy \right)^{1/s} \\ &\leq CM_{\alpha s}(g^s)(x)^{1/s}. \end{aligned}$$

Further, $M(g_2^s)^{1/s} \in A_1$ with a constant independent of g , (see [5, p. 158]), so

$$\frac{|Q|^{\alpha/n}}{|Q|} \int_Q M(g_2^s)(y)^{1/s} dy \leq C|Q|^{\alpha/n} M(g_2^s)(x)^{1/s}.$$

There exists a cube P containing x such that

$$M(g_2^s)(x) \leq \frac{2}{|P|} \int_P g_2^s dy.$$

Since P must intersect $\mathbb{R}^n \setminus 3Q$, $Q \subset 3P$. Therefore,

$$|Q|^{\alpha/n} M(g_2^s)(x)^{1/s} \leq C|3P|^{\alpha/n} \left(\frac{1}{|3P|} \int_{3P} g^s dy \right)^{1/s} \leq CM_{\alpha s}(g^s)(x)^{1/s}.$$

□

We can now prove Theorem 1.1. Fix p , $1 < p < \infty$, and a function $f \in L^p(v)$; by a standard argument we may assume that f is non-negative, bounded and has

compact support. For each $t > 0$, let $E_t = \{x \in \mathbb{R}^n : I_\alpha f(x) > t\}$. By duality there exists a function $G_t \in L^{p'}$, $\|G_t\|_{p'} = 1$, such that

$$u(E_t)^{1/p} = \|u^{1/p} \chi_{E_t}\|_p = \int_{E_t} u^{1/p} G_t \, dx.$$

Fix $s > 1$ as in Theorem 2.3, and let $w_t = M(u^{s/p} G_t^s)^{1/s}$. Then $w_t \in A_1 \subset A_\infty$, and the A_∞ constant of w_t depends only on s . Hence, by Theorems 2.1 and 2.2, and by Lemma 3.1,

$$\begin{aligned} \sup_{t>0} t u(E_t)^{1/p} &= \sup_{t>0} t \int_{E_t} u^{1/p} G_t \, dx \\ &\leq \sup_{t>0} t w_t(E_t) \\ &\leq C \sup_{t>0} t w_t(\{x \in \mathbb{R}^n : M_\alpha f(x) > t\}) \\ &\leq C \sup_{t>0} \int_{\mathbb{R}^n} f M_\alpha(w_t) \, dx \\ &\leq C \sup_{t>0} \int_{\mathbb{R}^n} f M_{\alpha s}(u^{s/p} G_t^s)^{1/s} \, dx. \end{aligned}$$

Then, by Hölder’s inequality and Theorem 2.3,

$$\begin{aligned} \sup_{t>0} t u(E_t)^{1/p} &\leq C \sup_{t>0} \left(\int_{\mathbb{R}^n} f^p v \, dx \right)^{1/p} \left(\int_{\mathbb{R}^n} M_{\alpha s}(u^{s/p} G_t^s)^{p'/s} v^{-p'/p} \, dx \right)^{1/p'} \\ &\leq C \sup_{t>0} \left(\int_{\mathbb{R}^n} f^p v \, dx \right)^{1/p} \left(\int_{\mathbb{R}^n} (u^{s/p} G_t^s)^{p'/s} u^{-p'/p} \, dx \right)^{1/p'} \\ &= C \sup_{t>0} \left(\int_{\mathbb{R}^n} f^p v \, dx \right)^{1/p}. \end{aligned}$$

REFERENCES

- [1] Cruz-Uribe D., SFO, *New proofs of two-weight norm inequalities for the maximal operator*, Georgian Math. J. **7** (2000), 33–42.
- [2] Cruz-Uribe D., SFO, Fiorenza A., *The A_∞ property for Young functions and weighted norm inequalities*, Houston J. Math., to appear.
- [3] Cruz-Uribe D., SFO, Pérez C., *Sharp two-weight, weak-type norm inequalities for singular integral operators*, Math. Res. Lett. **6** (1999), 417–428.
- [4] Cruz-Uribe D., SFO, Pérez C., *Two-weight, weak-type norm inequalities for fractional integrals, Calderón-Zygmund operators and commutators*, Indiana Math. J. **49** (2000), 697–721.
- [5] García-Cuerva J., Rubio de Francia J.L., *Weighted Norm Inequalities and Related Topics*, North Holland Math. Studies 116, North Holland, Amsterdam, 1985.
- [6] Muckenhoupt B., Wheeden R., *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. **192** (1974), 261–274.

- [7] Pérez C., *Two weighted inequalities for potential and fractional type maximal operators*, Indiana Math. J. **43** (1994), 663–683.
- [8] Sawyer E.T., *Weighted norm inequalities for fractional maximal operators*, 1980 Seminar on Harmonic Analysis, CMS Conf. Proc. 1, pp.283–309, Amer. Math. Soc., Providence, 1981.
- [9] Sawyer E.T., Wheeden R., *Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces*, Amer. J. Math. **114** (1992), 813–874.

DEPARTMENT OF MATHEMATICS, TRINITY COLLEGE, HARTFORD, CT 06106-3100, USA

E-mail: david.cruzuribe@mail.trincoll.edu

(Received November 22, 2000, revised May 24, 20010)