

T. F. Zhuraev

On projectively quotient functors

Commentationes Mathematicae Universitatis Carolinae, Vol. 42 (2001), No. 3, 561--573

Persistent URL: <http://dml.cz/dmlcz/119271>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2001

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On projectively quotient functors

T.F. ZHURAEV

Abstract. We introduce notions of projectively quotient, open, and closed functors. We give sufficient conditions for a functor to be projectively quotient. In particular, any finitary normal functor is projectively quotient. We prove that the sufficient conditions obtained are necessary for an arbitrary subfunctor \mathcal{F} of the functor \mathcal{P} of probability measures. At the same time, any “good” functor is neither projectively open nor projectively closed.

Keywords: projectively closed functor, finitary functor, functor of probability measures

Classification: 54B30

Introduction

All spaces are assumed to be Tychonoff and all mappings are continuous.

Recall that a covariant functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ acting in the category of compact spaces is called *normal* if it has the following normality properties:

- *preserves the empty set and the singletons*, i.e., $\mathcal{F}(\emptyset) = \emptyset$ and $\mathcal{F}(\{1\}) = \{1\}$, where $\{k\}$ ($k \geq 0$) denotes the set $\{0, 1, \dots, k - 1\}$ of nonnegative integers smaller than k . In this notation, $0 = \{\emptyset\}$;
- *is monomorphic*, i.e., for any (topological) embedding $f: A \rightarrow X$, the mapping $\mathcal{F}(f): \mathcal{F}(A) \rightarrow \mathcal{F}(X)$ is also an embedding;
- *is epimorphic*, i.e., for any surjective mapping $f: X \rightarrow Y$, the mapping $\mathcal{F}(f): \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ is surjective;
- *preserves intersections*, i.e., for any family $\{A_\alpha : \alpha \in \mathcal{A}\}$ of closed subsets of a compact space X , the mapping $\mathcal{F}(i): \bigcap \{\mathcal{F}(A_\alpha) : \alpha \in \mathcal{A}\} \rightarrow \mathcal{F}(X)$ defined by $\mathcal{F}(i)(x) = \mathcal{F}(i_\alpha)(x)$, where $i_\alpha: A_\alpha \rightarrow X$ are the identity embeddings for all $\alpha \in \mathcal{A}$, is an embedding;
- *preserves preimages*, i.e., for any mapping $f: X \rightarrow Y$ and an arbitrary closed set $A \subset Y$, the mapping $\mathcal{F}(f \upharpoonright f^{-1}(A))(f^{-1}(A)) \rightarrow \mathcal{F}(A)$ is a homeomorphism;
- *preserves weight*, i.e., $w(\mathcal{F}(X)) = w(X)$ for any infinite compact space X ;
- *is continuous*, i.e., for any inverse spectrum $S = \{X_\alpha; \pi_\beta^\alpha : \alpha \in \mathcal{A}\}$ of compact spaces, the limit $f: \mathcal{F}(\lim S) \rightarrow \lim \mathcal{F}(S)$ of the mappings $\mathcal{F}(\pi_\alpha)$, where $\pi_\alpha: \lim S \rightarrow X_\alpha$ are the limiting projections of the spectrum S , is a homeomorphism.

In what follows, we assume that all functors under consideration are monomorphic and preserve intersections. We also assume that all functors preserve nonempty spaces. The latter assumption is not an essential limitation; the only functor it excludes from consideration is the empty functor, i.e., the functor \mathcal{F} that maps any space to the empty set.

Indeed, suppose that $\mathcal{F}(X) = \emptyset$ for some nonempty compact space X . Then $\mathcal{F}(\emptyset) = \mathcal{F}(1) = \emptyset$, because \mathcal{F} is monomorphic. Let Y be an arbitrary nonempty compact space. Consider the constant mapping $f: Y \rightarrow 1$. We have $\mathcal{F}(f)(\mathcal{F}(Y)) \subset \mathcal{F}(1) = \emptyset$; therefore, the space $\mathcal{F}(Y)$ is empty, because it is mapped to the empty set. Thus, we have proved that there exists a unique monomorphic functor that does not preserve nonempty spaces.

By exp , we denote the well-known functor of hyperspace of closed subsets. This functor maps every (nonempty) compact space X to the set $\text{exp}(X)$ of all its nonempty closed subsets endowed with the (finite) Vietoris topology (see [5]) and a continuous mapping $f: X \rightarrow Y$ to the mapping $\text{exp}(f): \text{exp}(X) \rightarrow \text{exp}(Y)$ defined by $\text{exp}(f)(A) = f(A)$.

In this paper, we introduce notions of projectively quotient, open, and closed functors. We give sufficient conditions for a functor to be projectively quotient (Theorem 1). In particular, any finitary normal functor is projectively quotient (Corollary 2). We prove that the sufficient conditions obtained are necessary for an arbitrary subfunctor \mathcal{F} of the functor \mathcal{P} of probability measures (Theorem 2). At the same time, any “good” functor is neither projectively open nor projectively closed (Theorems 3 and 4).

The main part

Let $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ be a functor. By $C(X, Y)$, we denote the space of continuous mappings from X to Y with the compact-open topology.

In particular, $C(\{k\}, Y)$ is naturally homeomorphic to the k th power Y^k of the space Y ; the homeomorphism takes each mapping $\xi: \{k\} \rightarrow Y$ to the point $(\xi(0), \dots, \xi(k - 1)) \in Y^k$.

For a functor \mathcal{F} , a compact space X , and a positive integer k , we define the mapping

$$\pi_{\mathcal{F}, X, k}: C(\{k\}, X) \times \mathcal{F}(\{k\}) \rightarrow \mathcal{F}(X)$$

by

$$\pi_{\mathcal{F}, X, k}(\xi, a) = \mathcal{F}(\xi)(a) \text{ for } \xi \in C(\{k\}, X) \text{ and } a \in \mathcal{F}(\{k\}).$$

When it is clear what functor \mathcal{F} and what space X are meant, we omit the subscripts \mathcal{F} and X and write $\pi_{X, k}$ or π_k instead of $\pi_{\mathcal{F}, X, k}$.

According to a theorem of Shchepin ([1], Theorem 3.1), the mapping

$$\mathcal{F}: C(Z, Y) \rightarrow C(\mathcal{F}(Z), \mathcal{F}(Y))$$

is continuous for any *continuous* functor \mathcal{F} and compact spaces Z and Y . This implies the following assertion.

Proposition 1 ([2]). *If \mathcal{F} is a continuous functor, X is a compact space, and k is a positive integer, then the mapping $\pi_{\mathcal{F},X,k}$ is continuous.*

Let \mathcal{F}_k be the subfunctor of a functor \mathcal{F} defined as follows. For a compact space X , $\mathcal{F}_k(X)$ is the image of the mapping $\pi_{\mathcal{F},X,k}$, and for a mapping $f: X \rightarrow Y$, $\mathcal{F}_k(f)$ is the restriction of $\mathcal{F}(f)$ to $\mathcal{F}_k(X)$. It is easy to verify that the diagram

$$\begin{array}{ccc}
 C(\{k\}, X) \times \mathcal{F}(\{k\}) & \xrightarrow{\bar{f} \times \text{id}} & C(\{k\}, Y) \times \mathcal{F}(\{k\}) \\
 \pi_{X,k} \downarrow & & \downarrow \pi_{Y,k} \\
 \mathcal{F}(X) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(Y),
 \end{array}$$

where $\bar{f}(\xi) = f \circ \xi$, is commutative; therefore, $\mathcal{F}(f)(\mathcal{F}_k(X)) \subset \mathcal{F}_k(Y)$, and \mathcal{F}_k is a functor.

A functor \mathcal{F} is called a *functor of degree n* if $\mathcal{F}_n(X) = \mathcal{F}(X)$ for any compact space X but $\mathcal{F}_{n-1}(X) \neq \mathcal{F}(X)$ for some X .

For a functor \mathcal{F} and an element $a \in \mathcal{F}(X)$, the *support* of a is defined as the intersection of all closed sets $A \subset X$ such that $a \in \mathcal{F}(A)$; it is denoted by $\text{supp}_{\mathcal{F}(X)}(a)$. When it is clear what functor and space are meant, we denote the support of a merely by $\text{supp}(a)$.

By definition,

$$(2) \quad f(\text{supp}(a)) \supset \text{supp}(\mathcal{F}(f)(a))$$

for a continuous mapping $f: X \rightarrow Y$ and $a \in \mathcal{F}(X)$. Clearly,

$$(3) \quad a \in \mathcal{F}(\text{supp}(a)).$$

If a functor \mathcal{F} *preserves preimages*, then \mathcal{F} preserves supports, i.e.,

$$(4) \quad f(\text{supp}(a)) = \text{supp}(\mathcal{F}(f)(a)).$$

Proposition 2. *For any functor \mathcal{F} and compact space X ,*

$$\mathcal{F}_k(X) = \{a \in \mathcal{F}(X) : |\text{supp}(a)| \leq k\}.$$

PROOF: The inclusion \subset follows from the definition of the set $\mathcal{F}_k(X)$ and condition (2). Let us verify the reverse inclusion. Suppose that $a \in \mathcal{F}(X)$ and $\text{supp}(a)$ consists of l different points x_0, x_1, \dots, x_{l-1} , where $l \leq k$. Consider the mapping $f: \text{supp}(a) \rightarrow \{k\}$ specified by $f(x_i) = i$. By (3), the element

$b = \mathcal{F}(f)(a) \in \mathcal{F}(\{k\})$ is defined. Let $\xi: \{k\} \rightarrow X$ be the mapping such that $\xi(i) = x_i$ for $i \leq l - 1$ and $\xi(j) = x_{l-1}$ for $l \leq j$. Then

$$\begin{aligned} \pi_k(\xi, b) &= \mathcal{F}(\xi)(b) \\ &= \mathcal{F}(\xi)(\mathcal{F}(f)(a)) = \mathcal{F}(\xi \circ f)(a) = \mathcal{F}(\text{id}_{\text{supp}(a)})(a) \\ &= \text{id}_{\mathcal{F}(\text{supp}(a))}(a) = a. \end{aligned}$$

Therefore, $a \in \mathcal{F}_k(X)$, which proves Proposition 2. □

The definition of support and property (3) imply the following assertion.

Proposition 3. *For a functor \mathcal{F} , a compact space X , and a closed subset A of X ,*

$$\mathcal{F}(A) = \{a \in \mathcal{F}(X) : \text{supp } a \subset A\}.$$

Chigogidze [3] extended an arbitrary intersection-preserving monomorphic functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ over the category Tych of Tychonoff spaces by setting

$$\mathcal{F}_\beta(X) = \{a \in \mathcal{F}(\beta X) : \text{supp}(a) \subset X\}$$

for any Tychonoff space X . If $f: X \rightarrow Y$ is a continuous mapping of Tychonoff spaces and $\beta f: \beta X \rightarrow \beta Y$ is the (unique) extension of f over their Stone-Ćech compactifications, then (2) implies that

$$\mathcal{F}(\beta f)(\mathcal{F}_\beta(X)) \subset \mathcal{F}_\beta(Y).$$

Therefore, we can define $\mathcal{F}_\beta(f) = \mathcal{F}(\beta f) \upharpoonright X$, which makes \mathcal{F}_β a functor.

Chigogidze proved [3] that, if a functor \mathcal{F} has some normality property, then \mathcal{F}_β also has this property (modified when necessary). The definition of the functor \mathcal{F}_β implies, in particular, that

$$(5) \quad f(\text{supp}_{\mathcal{F}_\beta(X)}(a)) = \text{supp}_{\mathcal{F}_\beta(Y)} \mathcal{F}_\beta(f)(a)$$

for any preimage-preserving functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$, continuous mapping $f: X \rightarrow Y$, and $a \in \mathcal{F}_\beta(X)$. In what follows, we denote both functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ and its extension $\mathcal{F}_\beta: \text{Tych} \rightarrow \text{Tych}$ over the category of Tychonoff spaces by the same symbol \mathcal{F} .

For a Tychonoff space X , a functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$, and a positive integer k , we put

$$\mathcal{F}_k(X) = \pi_{\mathcal{F}, \beta X, k}(C(\{k\}, X) \times \mathcal{F}(k))$$

and denote the restriction of $\pi_{\mathcal{F}, \beta X, k}$ to $C(\{k\}, X) \times \mathcal{F}(\{k\})$ by $\pi_{\mathcal{F}, X, k}$. If $f: X \rightarrow Y$ is a continuous mapping, then $\mathcal{F}(\beta f)(\mathcal{F}_k(X)) \subset \mathcal{F}_k(Y)$; this is implied by the

commutativity of diagram (1) for the mapping βf . Therefore, setting $\mathcal{F}_k(f) = \mathcal{F}(\beta f|_{\mathcal{F}(X)})$, we obtain a mapping

$$\mathcal{F}_k(f): \mathcal{F}_k(X) \rightarrow \mathcal{F}_k(Y).$$

Thus, we have defined a covariant functor $\mathcal{F}_k: \text{Tych} \rightarrow \text{Tych}$ that extends $\mathcal{F}_k: \text{Comp} \rightarrow \text{Comp}$. Proposition 2 implies the following assertion.

Proposition 4. *If $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ is a functor, then $\mathcal{F}_k: \text{Tych} \rightarrow \text{Tych}$ is a subfunctor of the functor \mathcal{F}_β , and*

$$(6) \quad \mathcal{F}_k(X) = \mathcal{F}_\beta(X) \cap \mathcal{F}_k(\beta X)$$

for any Tychonoff space X .

Proposition 5 ([1, Proposition 3.11]). *For any compact space X and functor \mathcal{F} , the mapping*

$$\text{supp}_{\mathcal{F}(X)} : \mathcal{F}(X) \rightarrow \text{exp } X$$

is lower semicontinuous.

Let $U \subset X$ be an open set. Put

$$\mathcal{F}_+(U) = \{a \in \mathcal{F}(X) : \text{supp}(a) \cap U \neq \emptyset\}.$$

Proposition 5 is equivalent to the assertion that the set $\mathcal{F}_+(U)$ is open for any open $U \subset X$.

Proposition 6. *For a compact space X , a functor \mathcal{F} , and a positive integer k , the set $\mathcal{F}_k(X)$ is closed in $\mathcal{F}(X)$.*

PROOF: Take $a \in \mathcal{F}(X) \setminus \mathcal{F}_k(X)$. According to Proposition 2, $|\text{supp}(a)| \geq k + 1$. Let x_0, \dots, x_k be pairwise different points from $\text{supp}(a)$, and let U_0, \dots, U_k be their pairwise disjoint neighborhoods. By Proposition 5, the set $\mathcal{F}_+(U_0) \cap \dots \cap \mathcal{F}_+(U_k)$ is then a neighborhood of a , and it is disjoint from $\mathcal{F}_k(X)$ by virtue of Proposition 2. \square

Remark 1. The definition of the set $\mathcal{F}_k(X)$ and Proposition 1 imply that $\mathcal{F}_k(X)$ is closed in $\mathcal{F}(X)$ for any continuous functor \mathcal{F} .

Propositions 4 and 6 imply the following assertion.

Proposition 7. *For a Tychonoff space X , a functor \mathcal{F} , and a positive integer k , the set $\mathcal{F}_k(X)$ is closed in $\mathcal{F}_\beta(X)$.*

Let us mention several simple but important facts.

Proposition 8. *If $f: X \rightarrow Y$ is a closed mapping and $Z \subset Y$, then the mapping*

$$f \upharpoonright f^{-1}(Z): f^{-1}(Z) \rightarrow Z$$

is also closed.

Proposition 9. *Let $f: X \rightarrow Y$ be a continuous surjective mapping, and let X_1, \dots, X_n be closed subsets of X such that*

1. $X = X_1 \cup \dots \cup X_n$;
2. all $f(X_i)$ are closed in Y ;
3. all $f \upharpoonright X_i: X_i \rightarrow f(X_i)$ are quotient mappings.

Then the mapping f is quotient.

Proposition 10. *If $f: X \rightarrow Y$ is a continuous mapping, X_0 is a subset of X : $f(X_0) = Y$, and $f \upharpoonright X_0$ is quotient, then f is also quotient.*

We say that a functor \mathcal{F} is *finitely open* if the set $\mathcal{F}_k(\{k+1\})$ is open in $\mathcal{F}(\{k+1\})$ for any positive integer k . For example, the *finitary* functors, i.e., the functors \mathcal{F} such that $\mathcal{F}(\{k\})$ are finite for all positive integers k , are finitely open. We say that a functor \mathcal{F} is *projectively quotient* if, for any Tychonoff space X and any positive integer k , the mapping

$$\pi_{\mathcal{F}, X, k}: C(\{k\}, X) \times \mathcal{F}(\{k\}) \rightarrow \mathcal{F}_k(X)$$

is quotient.

Theorem 1. *Any continuous finitely open functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ preserving the empty set and preimages is projectively quotient.*

PROOF: We prove that the mappings $\pi_{\mathcal{F}, X, k} = \pi_k$ are quotient by induction on k .

The mapping π_1 is bijective. Indeed, $\text{supp}(\mathcal{F}(\xi)(a)) = \xi(0)$ by (5). (We have $\text{supp}(a) \neq \emptyset$, because \mathcal{F} preserves the empty set.) This and the injectivity of the mapping $\mathcal{F}(\xi)$ gives the injectivity of π_1 . As to the inverse mapping, it takes $\mathcal{F}(\xi)(a)$ to the pair (ξ, a) . Therefore, the mapping $\pi_{\mathcal{F}, X, 1}$ is a homeomorphism as the restriction of the homeomorphism $\pi_{\mathcal{F}, \beta X, 1}$ to a subset.

Suppose that the mappings π_i are quotient for all positive integers $i \leq k$. Let us prove that π_{k+1} is quotient. We denote the identity embedding of the space $\{k\}$ into $\{k+1\}$ by e and put $\Phi_0 = \{a \in \mathcal{F}(\{k+1\}) : \text{supp}(a) \subset e(\{k\})\}$ and $\Phi_1 = \mathcal{F}(\{k+1\}) \setminus \mathcal{F}(\{k\})$. According to Proposition 3, the set Φ_0 coincides with $\mathcal{F}(e(\{k\}))$ and is therefore compact. The set Φ_1 is also compact, because \mathcal{F} is finitely open. Next, we put

$$Z_i^0 = C(\{k+1\}, X) \times \Phi_i \quad \text{for } i = 0, 1$$

and

$$Z_i^1 = C(\{k+1\}, \beta X) \times \Phi_i \quad \text{for } i = 0, 1$$

and denote the restriction of $\pi_{\beta X, k+1}$ to Z_i^j by f_i^j ($i, j \in 2$).

Let us prove that

$$(7) \quad f_0^0(Z_0^0) = \mathcal{F}_k(X).$$

For this purpose, it is sufficient to specify an epimorphism $g: Z_0^0 \rightarrow C(\{k\}, X) \times \mathcal{F}(\{k\})$ such that

$$(8) \quad f_0^0 = \pi_{X,k} \circ g.$$

Consider the mapping $h: C(\{k+1\}, X) \rightarrow C(\{k\}, X)$ defined by $h(\xi) = \xi \circ e$. Under the identification of $C(\{i\}, X)$ with the power space X^i , the mapping h corresponds to the projection $X^{k+1} \rightarrow X^k$ parallel to the last coordinate. Therefore, the mapping h is open. The mapping $g = h \times \text{id}_{\Phi_0}$ is also open. For $(\xi, a) \in C(\{k+1\}, X) \times \Phi_0$, we have

$$\begin{aligned} (\pi_{X,k} \circ g)(\xi, a) &= \pi_{X,k}(h(\xi), a) \\ &= \pi_{X,k}(\xi \circ e, a) = \mathcal{F}(\xi \circ e)(a) = \mathcal{F}(\xi)(\mathcal{F}(e)(a)) = \mathcal{F}(\xi)(a) \\ &= \pi_{X,k+1}(a) = f_0^0(a) \end{aligned}$$

(the equality $\mathcal{F}(\xi)(\mathcal{F}(e)(a)) = \mathcal{F}(\xi)(a)$ holds because $\text{supp}(a) \subset e(\{k\})$). This proves (8) and, thereby, (7). In addition, (8) implies that the mapping f_0^0 is quotient as the composition of the open mapping g and the quotient (by the induction hypothesis) mapping $\pi_{X,k}$.

Now, let us establish some properties of the mappings f_1^0 and f_1^1 . First,

$$(9) \quad Z_1^0 = (f_1^1)^{-1} f_1^1(Z_1^0).$$

Let us show this. Note that, according to (5),

$$(10) \quad \text{supp}(\pi_{k+1}(\xi, a)) = \xi(\text{supp}(a))$$

for any $\xi \in C(\{k+1\}, \beta X)$ and $a \in \mathcal{F}(\{k+1\})$. If $(\xi, a) \in Z_1^1$, then $\text{supp}(a) = \{k+1\}$, and hence

$$(11) \quad \text{supp } f_1^1(\xi, a) = \xi(\{k+1\}).$$

Therefore,

$$(12) \quad Z_1^0 = \{ \{ \xi, a \} \in Z_1^1 : \xi(\{k+1\}) \subset X \}.$$

Hence, if $f_1^1(\xi_0, a_0) = f_1^1(\xi_1, a_1)$ and $(\xi_0, a_0) \in Z_1^0$, then $(\xi_1, a_1) \in Z_1^0$. This proves (9).

Equality (9), Proposition 8, and the compactness of the set Z_1^1 , which follows from the assumption that \mathcal{F} is finitely open, imply that the mapping f_1^0 is closed and, consequently, quotient. Let us show that

$$(13) \quad f_1^0(Z_1^0) = f_1^1(Z_1^1) \cap \mathcal{F}_{k+1}(X).$$

It is sufficient to verify the \supset inclusion. Suppose that $f_1^1(\xi, a) \in f_1^1(Z_1^1) \cap \mathcal{F}_{k+1}(X)$. Then $X \supset \text{supp}(f_1^1(\xi, a)) = \xi(\{k + 1\})$ (the last equality is implied by (11)). Therefore, by (12), $(\xi, a) \in Z_1^0$, which proves (13).

Equality (13) and the compactness of $f_1^1(Z_1^1)$ imply the closedness of $f_1^0(Z_1^0)$ in $\mathcal{F}_{k+1}(X)$. In addition, the inclusion $f_1^1(Z_1^1) \supset \mathcal{F}_{k+1}(\beta X) \setminus \mathcal{F}_k(\beta X)$ and (13) give

$$(14) \quad f_1^0(Z_1^0) \supset \mathcal{F}_{k+1}(X) \setminus \mathcal{F}_k(X).$$

Put $Z = Z_0^0 \cup Z_1^0$ and $f = \pi_{X,k+1} \upharpoonright Z$. By (7) and (14), $\text{Im } f = \mathcal{F}_{k+1}(X)$. As mentioned, the mappings $f_1^0 = f|_{Z_1^0}$ are quotient, and the set $f_1^0(Z_1^0) = f(Z_1^0)$ is closed. Equality (7) and Propositions 4 and 7 imply that the set $f(Z_0^0)$ is also closed. According to Proposition 9, the mapping $f: Z \rightarrow \mathcal{F}_{k+1}(X)$ is quotient. This and Proposition 10 imply that the mapping $\pi_{X,k+1}$ is quotient too, which completes the proof of Theorem 1. \square

Corollary 1. *Any finitary continuous functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ preserving the empty set and preimages is projectively quotient.*

Corollary 2. *Any finitary normal functor, in particular, the hyperspace functor exp , is projectively quotient.*

In relation to Theorem 1 and its corollaries, several questions arise.

The first question is as follows:

Question 1. Is Theorem 1 valid without the assumption that the functor \mathcal{F} is finitely open?

This question is especially important because not all normal functors are finitely open. In particular, the functor \mathcal{P} of probability measures (which is the most interesting normal functor) is not finitely open. Theorem 2 proved below not only gives a negative answer to Question 1, but also characterizes the quotient normal subfunctors of the functor \mathcal{P} .

Arbitrary normal subfunctors of the functor \mathcal{P} are described in [5], [6].

Theorem 2. *A normal subfunctor \mathcal{F} of the functor \mathcal{P} is projectively quotient if and only if \mathcal{F} is finitely open.*

PROOF: The ‘if’ part follows from Theorem 1. Let us prove the ‘only if’ part. Suppose that the set $\mathcal{F}(\{k + 1\}) \setminus \mathcal{F}_k(\{k + 1\})$ is not closed for some positive

integer k . Take a sequence $\{\mu_n : n \in \omega\} \subset \mathcal{F}(\{k+1\}) \setminus \mathcal{F}_k(\{k+1\})$ of measures converging to a measure $\mu \in \mathcal{F}_k(\{k+1\})$. The support of the measure μ comprises no more than k points. By symmetry considerations, we can assume that these points are among the first k points of the set $\{k+1\}$, i.e., $\mu \in \mathcal{F}(\{k\}) \subset \mathcal{F}_k(\{k+1\})$. Each finitely supported probability measure is a convex combination of Dirac measures. Suppose that

$$\mu = m_0\delta(0) + \dots + m_{k-1}\delta(k-1)$$

and

$$\mu_n = m_0^n\delta(0) + \dots + m_{k-1}^n\delta(k-1) + m_k^n\delta(k).$$

Some of the numbers m_i may be zero, while all m_j^n are nonzero. Since the sequence $\{\mu_n\}$ converges to μ , the sequence $\{m_k^n\}$ converges to zero. Consider the mapping $f: \{k+1\} \rightarrow \{2\}$ that takes the set $\{k\}$ to 0 and the point $k \in \{k+1\}$ to 1. The image of the measure μ under this mapping is $\lambda = m_0\delta(f(0)) + \dots + m_{k-1}\delta(f(k-1)) = \delta(0)$, and the image of μ_n is $\lambda_n = e^n\delta(0) + m_k^n\delta(1)$, where $e^n = m_0^n + \dots + m_{k-1}^n$. Clearly, $\lambda_n \in \mathcal{F}(\{2\}) \setminus \mathcal{F}_1(\{2\})$ and $\lambda = \delta(0) \in \mathcal{F}_1(\{2\})$. The convergence of the sequence $\{\lambda_n\}$ to the measure λ implies that the functor \mathcal{F}_2 is not finitely open. Let us take the space ω of nonnegative integers as X and show that the mapping $\pi_2 = \pi_{\mathcal{F}, \omega, 2}: C(2, \omega) \times \mathcal{F}_2(\omega) \rightarrow \mathcal{F}_2(\omega)$ is not quotient.

Take the measure $\nu_n = e^n\delta(0) + m_k^n\delta(n+1)$ in $\mathcal{F}_2(\omega)$. Since the sequence $\{m_k^n\}_n$ converges to zero, the sequence $\bar{H} = \{\nu_n\}$ converges to the Dirac measure $\delta(0)$. Thus, it remains to prove that the set $\pi_2^{-1}(\bar{H})$ is not closed. Let us show that it is discrete. Identifying $C(2, \omega)$ with ω^2 , we easily see that $\pi_2^{-1}(\nu_n)$ consists of the two points

$$c_n^0 = ((0, n+1), e^n\delta(0) + m_k^n\delta(1))$$

and

$$c_n^1 = ((n+1, 0), m_k^n\delta(0) + e^n\delta(1)).$$

Clearly, the set $\pi_2^{-1}(\bar{H})$ is discrete in itself. Suppose that there exists a point $\eta \in \overline{\pi_2^{-1}(\bar{H})} \setminus \pi_2^{-1}(\bar{H})$. Let $\eta = ((a_0, a_1), \alpha)$. Then the definition of c_n^i for $i = 0, 1$ implies that $a_i = 0$ for some i . To be definite, we assume that $a_0 = 0$. Then $a_1 = n$ for some $n \geq 1$. Therefore, the set $\{(0, n)\} \times \mathcal{F}(\{2\})$ is a neighborhood of η containing only one point c_{n-1}^0 of the set $\pi_2^{-1}(\bar{H})$. This completes the proof of Theorem 2. □

The second question is also related to Theorem 1:

Question 2. Is Theorem 1 valid without the assumption that \mathcal{F} preserves the empty set?

A negative answer to this question is given by the *metrizable cone functor* Cone_m . Recall that $\text{Cone}_m(X)$ is the cone over the space X such that its vertex

v_x has a countable neighborhood base. For any metrizable space X , $\text{Cone}_m(X)$ is metrizable. Hence, for example, the mapping $\pi_{\mathbb{R},1} : \mathbb{R} \times [0, 1] \rightarrow \text{Cone}_m(\mathbb{R})$ is not quotient.

The third question is as follows:

Question 3. Is Theorem 1 valid without the assumption that \mathcal{F} preserves preimages?

I do not know the answer to this question. The best known non-preimage-preserving functor is the superextension functor λ , which has all the normality properties except this one.

Question 4. Is the functor λ projectively quotient?

Another group of problems is related to the potential possibility of obtaining stronger properties of the mapping $\pi_{\mathcal{F},X}$ under certain constraints on the functor \mathcal{F} . We say that a functor \mathcal{F} is *projectively open (closed)* if the mapping $\pi_{\mathcal{F},X,k}$ is open (closed) for any Tychonoff space X and a positive integer k . A functor \mathcal{F} is said to be *finitely nondegenerate* if the set $\mathcal{F}(\{k + 1\}) \setminus \mathcal{F}_k(\{k + 1\})$ is nonempty for some positive integer k .

Proposition 11. *If \mathcal{F} is a finitely nondegenerate functor preserving preimages, then*

$$\mathcal{F}(\{2\}) \setminus \mathcal{F}_1(\{2\}) \neq \emptyset.$$

PROOF: Let k be the positive integer mentioned in the definition of a finitely nondegenerate functor. Take some element $a \in \mathcal{F}(\{k + 1\}) \setminus \mathcal{F}_k(\{k + 1\})$. Consider the mapping f from $\{k + 1\}$ to $\{2\}$ that takes the set $\{k\}$ to the point $0 \in \{2\}$ and the point $k \in \{k + 1\}$ to the point $1 \in \{2\}$. Put $b = \mathcal{F}(f)(a)$. Since $\text{supp}(a) = \{k + 1\}$ and \mathcal{F} preserves supports, we have $\text{supp}(b) = f(\text{supp}(a)) = f(\{k + 1\}) = \{2\}$. This proves Proposition 11. \square

Proposition 12. *If \mathcal{F} is a finitely nondegenerate continuous functor preserving preimages and singletons, then $\mathcal{F}_1(X)$ is nowhere dense in $\mathcal{F}_2(X)$ for any nonempty first countable compact space X without isolated points.*

PROOF: Take $a \in \mathcal{F}_1(X)$. By the definition of $\mathcal{F}_1(X)$, there exist $\xi \in C(\{1\}, X) = X$ and $b \in \mathcal{F}_1(\{1\})$ such that $a = \pi_1(\xi, b) = \mathcal{F}(\xi)(b)$. Since \mathcal{F} is finitely nondegenerate, by Proposition 11, there exists $c \in \mathcal{F}(\{2\}) \setminus \mathcal{F}_1(\{2\})$. The assumptions made about the compact space X imply that there exist two disjoint sequences $\{x_n\}$ and $\{y_n\}$ converging to the point $x = \xi(0)$. Let $\xi_n \in C(\{2\}, X)$ be the mapping defined by $\xi_n(0) = x_n$ and $\xi_n(1) = y_n$. Put $a_n = \pi_2(\xi_n, c)$. Since $\text{supp}(c) = \{2\}$ and \mathcal{F} preserves supports, we have

$$\text{supp}(a_n) = \text{supp}(\mathcal{F}(\xi_n)(c)) = \xi_n(\text{supp}(c)) = \xi_n(\{2\}) = \{x_n, y_n\}.$$

Therefore, $a_n \in \mathcal{F}_2(X) \setminus \mathcal{F}_1(X)$. Let $\psi \in C(\{2\}, X)$ be the mapping defined by $\psi(i) = \xi(0) = x$ for $i = 0, 1$. Clearly, the sequence $\{\xi_n\}$ converges to ψ . Therefore, the sequence $\{(\xi_n, c)\}$ converges to (ψ, c) , and the sequence $\{\pi_2(\xi_n, c)\} = \{a_n\}$ converges to $\pi_2(\psi, c)$, because the mapping π_2 is continuous. Since $\text{supp}(c) = \{2\}$ and \mathcal{F} preserves supports, we have

$$\text{supp}(\pi_2(\psi, c)) = \psi(\text{supp}(c)) = \psi(\{2\}) = \{\psi(0), \psi(1)\} = \{x, x\} = \{x\};$$

this and (3) give $\pi_2(\psi, c) \in \mathcal{F}(\{x\})$. On the other hand, $\text{supp}(a) = \{x\}$, and hence $a \in \mathcal{F}\{x\}$. Therefore, $a = \pi_2(\psi, c)$, because \mathcal{F} preserves singletons. Thus, for an arbitrary point $a \in \mathcal{F}_1(X)$, we can find a sequence $\{a_n\} \subset \mathcal{F}_2(X) \setminus \mathcal{F}_1(X)$ converging to a . This proves Proposition 12. \square

Theorem 3. *No finite nondegenerate finitely open continuous functor \mathcal{F} preserving preimages and singletons is projectively open.*

PROOF: It is sufficient to show that the mapping $\pi_2 = \pi_{\mathcal{F}, I, 2}$, where I is a closed number interval, is not open. Since the functor \mathcal{F} is finitely open, the set $\mathcal{F}_1(\{2\})$ is open in $\mathcal{F}(\{2\})$. Therefore, the set $C(\{2\}, I) \times \mathcal{F}_1(\{2\})$ is open in $C(\{2\}, I) \times \mathcal{F}(\{2\})$. Thus, if the mapping π_2 were open, then the set $\pi_2(C(\{2\}, I) \times \mathcal{F}_1(\{2\}))$ would be open in $\mathcal{F}_2(I)$. But, since \mathcal{F} preserves supports, $\pi_2(C(\{2\}), I) \times \mathcal{F}_1(\{2\})$ is contained in (and coincides with) the set $\mathcal{F}_1(X)$, which is nowhere dense by Proposition 12. This completes the proof of Theorem 3. \square

Remark 2. In Theorem 3, the assumption that \mathcal{F} is finitely nondegenerate is essential. As an example, we can take the continuum exponent functor exp^c (of hyperspace of subcontinua). It is a finitary functor satisfying all the normality conditions, except it is not epimorphic. For any Tychonoff space X and positive integer k , we have $(\text{exp}^c)_k(X) = (\text{exp}^c)_1(X) = X$ and $C(\{k\}, X) \times \text{exp}^c(\{k\}) = X^k \times \{k\}$. The mapping $\pi_k = \pi_{\text{exp}^c, X, k}$ is open, because it is the sum of the mappings $\pi_k \upharpoonright X^k \times \{i\}$ of open subspaces $X^k \times \{i\}$, where each $\pi_k \upharpoonright X^k \times \{i\}$ coincides with the projection $X^k \rightarrow X$ onto the i th coordinate.

Theorem 4 proved below shows that no “good” functors \mathcal{F} can be projectively closed. As previously, we start with auxiliary statements. Recall that a functor $\mathcal{F}: \text{Comp} \rightarrow \text{Comp}$ is called a *functor with continuous supports* if, for any compact space X , the mapping

$$\text{supp}_{\mathcal{F}(X)}: \mathcal{F}(X) \rightarrow \text{exp } X$$

is continuous. Note that, for any Tychonoff space X , the mapping

$$\text{supp}_{\mathcal{F}_\beta(X)}: \mathcal{F}_\beta(X) \rightarrow \text{exp } X$$

is continuous as the restriction of the continuous mapping $\text{supp}_{\mathcal{F}(\beta X)}$ to $\mathcal{F}_\beta(X)$. In what follows, we denote the mapping $\text{supp}_{\mathcal{F}_\beta(X)}$ by $\text{supp}_{\mathcal{F}(X)}$.

Proposition 13. *For a functor \mathcal{F} with continuous supports, the mapping*

$$\text{supp}_{\mathcal{F}(X)}: \mathcal{F}(X) \rightarrow \exp X$$

is closed as a mapping onto its image.

Indeed, it suffices to show that $\text{supp}_{\mathcal{F}(X)}$ is the restriction of the closed mapping $\text{supp}_{\mathcal{F}(\beta X)}$ to its full preimage ([6]), i.e.,

$$\text{supp}_{\mathcal{F}(X)}^{-1}(K) = \text{supp}_{\mathcal{F}(\beta X)}^{-1}(K)$$

for any compact $K \subset X$. But this follows from the definition of the set

$$\mathcal{F}(X) = \mathcal{F}_\beta(X) \subset \mathcal{F}(\beta X).$$

Theorem 4. *No preimage-preserving continuous functor \mathcal{F} with continuous supports is projectively closed.*

PROOF: It is sufficient to show that the mapping $\pi_{\mathcal{F}, \mathbb{R}, 2}$ is not closed. First, we do this for the functor $\mathcal{F} = \exp$. Consider $Z_n = ((\frac{1}{n}, n), 0) \in \mathbb{R}^2 \times \exp(\{2\})$ and $Z = \{Z_n : n = 1, 2, \dots\}$.

Obviously, the set Z is closed in $\mathbb{R}^2 \times \exp(\{2\})$, while $\pi_{\exp, \mathbb{R}, 2}(Z) = \{\{\frac{1}{n}\} : n = 1, 2, \dots\}$ is a sequence converging to $\{0\} \in \exp_2(\mathbb{R})$; here, $\{0\}$ is the nonempty subset comprising one element $0 \in \mathbb{R}$. Thus, not only the mapping $\pi_{\exp, \mathbb{R}, 2}$, but also its restriction $\pi_{\{0\}}$ to the closed set $C(\{2\}, \mathbb{R}) \times \{0\}$ is not closed.

Now, suppose that $\pi_{\mathcal{F}, \mathbb{R}, 2}$ is closed for some functor \mathcal{F} . Take any $a \in \mathcal{F}(\{0\})$ (recall that all functors are assumed to preserve nonempty subsets). Then the mapping $\pi_a = \pi_{\mathcal{F}, \mathbb{R}, 2} \upharpoonright C(\{2\}, \mathbb{R}) \times \{a\}$ is closed as the restriction of the closed mapping $\pi_{\mathcal{F}, \mathbb{R}, 2}$ to a closed set. Let

$$h_a: C(\{2\}, \mathbb{R}) \times \{a\} \rightarrow C(\{2\}, \mathbb{R}) \times \{0\}$$

be the homeomorphism defined by

$$h_a(\xi, a) = (\xi, 0).$$

Then

$$(15) \quad \pi_{\{0\}} \circ h_a = \text{supp}_{\mathcal{F}, \mathbb{R}, 2} \circ \pi_a.$$

Indeed,

$$\begin{aligned} (\text{supp}_{\mathcal{F}, \mathbb{R}, 2} \circ \pi_a)(\xi, a) &= \text{supp}(\pi_a(\xi, a)) = \text{supp}(\mathcal{F}(\xi)(a)) = \xi(\text{supp}(a)) = \xi(0) \\ (\text{supp}(\mathcal{F}(\xi)(a)) = \xi(\text{supp}(a)) &\text{ because } \mathcal{F} \text{ preserves supports). On the other hand,} \end{aligned}$$

$$\pi_{\{0\}} \circ h_a(\xi, a) = \pi_{\{0\}}(\xi, 0) = \exp(\xi)(0) = \xi(0).$$

Thus, (15), the closedness of π_a , and Proposition 13 imply that the mapping $\pi_{\{0\}} \circ h_a$ is closed. Therefore, $\pi_{\{0\}}$ is also closed as a left divisor of a closed mapping (here, it is only essential that h_a is epimorphic). This contradiction completes the proof of Theorem 4. □

Acknowledgment. I would like to thank the referee for helpful remarks.

REFERENCES

- [1] Shchepin E.V., *Functors and uncountable powers of compact spaces*, Uspekhi Mat. Nauk **36** (1981), no. 3, 3–62.
- [2] Basmanov V.N., *Covariant functors, retracts and dimension*, Dokl. Akad. Nauk USSR **271** (1983), 1033–1036.
- [3] Chigogidze A.Ch., *Extension of normal functors*, Vestnik Mosk. Univ. Ser. I Mat. Mekh. **6** (1984), 40–42.
- [4] Fedorchuk V.V., *Probability measures in topology*, Uspekhi Mat. Nauk **46** (1991), no. 1, 41–80.
- [5] Fedorchuk V.V., Filippov V.V., *General Topology: Basic Constructions*, Moscow, Mosk. Gos. Univ., 1988.
- [6] Engelking R., *General Topology*, Warszawa, PWN, 1977.

DEPARTMENT OF GENERAL TOPOLOGY AND GEOMETRY, MECHANICS AND MATHEMATICS
FACULTY, MOSCOW STATE UNIVERSITY, VOROB'EVY GORY, MOSCOW, 119899 RUSSIA

(Received June 22, 2000, revised November 10, 2000)