

Igor V. Protasov

Remarks on extremally disconnected semitopological groups

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 2, 343--347

Persistent URL: <http://dml.cz/dmlcz/119324>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Remarks on extremally disconnected semitopological groups

I.V. PROTASOV

Abstract. Answering recent question of A.V. Arhangel'skii we construct in ZFC an extremally disconnected semitopological group with continuous inverse having no open Abelian subgroups.

Keywords: extremally disconnected and maximal spaces, semitopological group, Stone-Čech compactification

Classification: Primary 54H11; Secondary 54C05, 54G15

All topological spaces under consideration are supposed to be Hausdorff. A topological space X is called *extremally disconnected* if the closure of every open subset of X is open. A topological space X without isolated points is called *maximal* if X has an isolated point in any stronger topology. Every maximal space is extremally disconnected. A group G provided with a topology τ is called maximal if (G, τ) is maximal as a topological space.

A group G provided with a topology is called *left (right) topological* if all mappings $x \mapsto gx, g \in G$ ($x \mapsto xg, g \in G$) are continuous. A group G with a topology τ is called *semitopological* if (G, τ) is left and right topological.

In [1] A.V. Arhangel'skii established some properties of extremally disconnected semitopological groups and posed three problems.

Problem 1. *Is there in ZFC an example of a non-discrete extremally disconnected topological group?*

This is a reminiscence of old (and still unsolved) problem from [2]. It is worth of mentioning that for some types of extremally disconnected topological groups the answer to Problem 1 is negative. For example, if there exists a maximal topological group, then there exists a P -point in ω^* , the remainder of the Stone-Čech compactification of the discrete space ω ([4, Theorem 7.3]). For further results in this direction see [5, Theorem 5.1], [7, Theorem 2.5] and [9].

Problem 2. *Is there in ZFC an example of a non-discrete extremally disconnected semitopological group with continuous inverse?*

Several kinds of such examples follow from [5] and [8]. We describe three of them.

By [5, Theorem 1.2], every infinite group G of cardinality α admits a maximal left invariant topology τ of dispersion character α . Remind that the dispersion character of a topological space (X, τ) is the cardinal $\Delta(\tau) = \min\{|U| : U \in \tau, U \neq \emptyset\}$. If G is Abelian then (G, τ) is semitopological. If G is Boolean ($g^2 = e$ for every $g \in G$, e is the identity of G) then (G, τ) is a semitopological group with continuous inverse. Note that a maximal left topological group need not be regular. However, every countable group admits a maximal regular left invariant topology ([5, Corollary 2.7]). It is still unknown ([5, Problem 2.10]) whether there is in ZFC an example of a regular maximal left topological group of uncountable dispersion character.

By [5, Theorem 1.3 and 4.5], every infinite group G admits an extremally disconnected left invariant topology τ such that (G, τ) is zero-dimensional (i.e. every point of G has base of neighborhoods consisting of clopen subsets) and left totally bounded (i.e. for every neighborhood U of the identity, there exists a finite subset F with $G = FU$). By the above argument, there is a zero-dimensional example to Problem 2 of arbitrary dispersion character.

Let τ, τ' be left invariant topologies on a group G . We say that (G, τ') is an open refinement of (G, τ) if $\tau \subseteq \tau'$ and every nonempty open subset from (G, τ') contains a nonempty open subset from (G, τ) . By [8], every left topological group (G, τ) has an extremally disconnected open refinement. If (G, τ) is regular, then there exists a zero-dimensional extremally disconnected open refinement (G, τ') . Now suppose that, for every element $g \in G$, there exists a neighborhood U of identity in τ such $gx = xg$ for each $x \in U$. Then every open refinement (G, τ') of (G, τ) is a semitopological group. In addition, if the subset $\{g : g^2 = e\}$ is open in (G, τ) , then the mapping $x \mapsto x^{-1}$ is continuous in (G, τ') .

Problem 3. *Let G be an extremally disconnected semitopological group with continuous inverse. Does there exist an open and closed Abelian subgroup of G ?*

There are two reasons for considering this problem. By Malykhin's theorem [1, Theorem 2], every extremally disconnected topological group has a clopen Boolean subgroup. By [1, Theorem 3], for every non-discrete extremally disconnected semitopological group with continuous inverse, there exists a neighborhood U of the identity e such that $g^2 = e$ for every $g \in U$. In one special case this problem has been mentioned in [6]: does every maximal semitopological group with continuous inverse contain an open Boolean subgroup?

The following two theorems give us a negative answer to Problem 3.

Theorem 1. *For every infinite cardinal α , there exists a semitopological group (G, τ) with continuous inverse and following properties: $\Delta(\tau) = \alpha$, (G, τ) has no open Abelian subgroups, (G, τ) is extremally disconnected and zero-dimensional.*

Theorem 2. *For every infinite cardinal α , there exists a maximal semitopological group (G, τ) with continuous inverse such that $\Delta(\tau) = \alpha$ and (G, τ) has no open Abelian subgroups.*

To prove these theorems we need some definitions, constructions and results from [3], [5].

Given a discrete space X , we take the points of βX , the Stone-Ćech compactification of X , to be the ultrafilters on X , with the points of X identified with the principal ultrafilters. The topology of βX can be defined by stating that the sets of the form $\{p \in \beta X : A \in p\}$, where A is a subset of X , form a base for the open sets. We note that the sets of this form are clopen and that, for any $p \in \beta X$ and any $A \subseteq X$, $A \in p$ if and only if $p \in \overline{A}$, where \overline{A} is a closure of A in βX . If A is a subset of X we shall use A^* to denote $\overline{A} \setminus A$, in particular X^* is a set of all free ultrafilters on X . For every filter φ on X denote $\overline{\varphi} = \{p \in \beta X : \varphi \subseteq p\}$, $\varphi^* = \overline{\varphi} \cap G^*$.

Let G be a discrete group. There are two natural ways for extension of multiplication from G to βG . We follow [3, Chapter 4]. Given any $p, q \in \beta G$ and $A \subseteq G$, put

$$A \in pq \text{ if and only if } \{g \in G : g^{-1}A \in q\} \in p.$$

Take any member $P \in p$ and, for every $x \in P$, choose some element $Q_x \in q$. Then $\bigcup_{x \in P} xQ_x \in pq$ and the family of subsets of this form is a base of the ultrafilter pq . This multiplication on βG is associative, so βG is a semigroup and G^* is a subsemigroup of βG .

Every closed subsemigroup of βG has an idempotent p , $p^2 = p$ ([3, Theorem 2.5]). Given any idempotent $p \in G^*$, the family of subsets $\{P \cup \{e\} : P \in p\}$ is a filter of neighborhoods of e for the uniquely determined maximal left invariant topology on G ([5, §1]). A group G provided with this topology is denoted by $G(p)$. We need also another type of topologies determined by idempotents. Fix $p \in G^*$ with $p^2 = p$ and, for every subset $A \subseteq G$, put $\text{cl}(A, p) = \{x \in G : A \in xp\}$. Then the family $\{\text{cl}(A, p) : A \in p\}$ is a base of neighborhoods of e for the uniquely determined zero-dimensional extremally disconnected left invariant topology on G ([5, §1]). A group G provided with this topology is denoted by $G[p]$.

Let X be an infinite set of cardinality α . For every permutation f of X , put $\text{supp } f = \{x \in X : f(x) \neq x\}$. Consider the group $S(X)$ of all permutations of X with finite supports. For every nonempty subset $Y \subseteq X$, identify $S(Y)$ with the subgroup of all permutations $f \in S(X)$ such that $f(x) \in Y$, $x \in Y$ and $f(x) = x$, $x \in X \setminus Y$. The identity permutation is denoted by e .

Let $\mathbf{F} = \{Y \subseteq X : X \setminus Y \text{ is finite}\}$ be a filter of all cofinite subsets of X . Denote by φ_0 the filter on $S(X)$ with base $\{S(Y) : Y \in \mathbf{F}\}$. Note that φ_0^* is a subsemigroup of $\beta S(X)$ and

(*) for every $f \in S(X)$, there exists $F \in \varphi$ such that $fg = gf$ for every $g \in F$.

Put $S_2(X) = \{g \in S(X) : g^2 = e\}$ and denote by φ_2 the filter on $S(X)$ with the base $\{F \cap S_2(X) : F \in \varphi_0\}$. By (*), φ_2^* is a subsemigroup of $\beta S(X)$.

Call a subset $A \subseteq S(X)$ sparse if there exists $x \in X$ such that $|\{g(x) : g \in A\}| = \alpha$. A filter φ' on $S(X)$ is called sparse if every member of φ' is sparse.

Clearly, φ_2 is a sparse filter. By Zorn's Lemma, for every sparse filter φ' , there exists a sparse ultrafilter p with $\varphi' \subseteq p$. Hence, the subset SP of all sparse ultrafilters from φ_2^* is nonempty. Clearly, SP is closed in $\beta S(X)$. For every sparse subset $A \subseteq S(X)$ and every $f \in S(X)$, the subset fA is sparse. It follows that SP is a subsemigroup of $\beta S(X)$. We shall use the following claim

(**) for every sparse subset $A \subseteq S(X)$, there exist $h, g \in A$ such that $hg \neq gh$.

To prove (**), choose $x \in X$ such that the subset $\{f(x) : f \in A\}$ is infinite. Fix any $h \in A$ with $h(x) \neq x$ and pick $g \in A$ such that $g(x) \notin \text{supp } h$. Since $hg(x) = g(x)$ and $h(x) \neq x$, then $hg(x) \neq gh(x)$ so $hg \neq gh$.

PROOF OF THEOREM 2: Put $G = S(X)$ and choose any idempotent $p \in SP$. Consider the maximal left topological group $(G, \tau) = G(p)$. Since p is sparse then $\Delta(\tau) = \alpha$. Since $\varphi_2 \subseteq p$ then, by (*), (G, τ) is right topological with continuous inverse. By (**), (G, τ) has no open Abelian subgroups. \square

PROOF OF THEOREM 1: Put $G = S(X)$ and choose any idempotent $p \in SP$. Consider the extremally disconnected zero-dimensional left topological group $(G, \tau) = G[p]$. Clearly, $\Delta(\tau) = \alpha$. Denote by τ_e the filter of neighborhoods of e in τ . For every ultrafilter q on G , $\tau_e \subseteq q$ if and only if $qp = p$ ([5, §2]). By (*) and the definition of product of ultrafilters, $\varphi_2 \subseteq \tau_e$. By the above paragraph, (G, τ) is right topological with continuous inverse and (G, τ) has no open Abelian subgroups. \square

We conclude the paper with four remarks.

1. Using arguments from [5, §2], we can add the following statement to Theorem 2: there exists a countable zero-dimensional maximal semitopological group with continuous inverse and without open Abelian subgroups.

2. A topological space S is called strongly extremally disconnected if, for every open nonclosed subset U of S , there exists $x \in \text{cl } U \setminus U$ such that $\{x\} \cup U$ is a neighborhood of x . Let (G, τ) be a left topological group and let an ultrafilter q converge to the identity in τ . By [8, Theorem 4.12], the strongest left invariant topology τ_q on G in which q converges to e is strongly extremally disconnected. Put $G = S(X)$ and denote by τ the left invariant topology on G such that φ_2 is a filter of neighborhoods of e . Choose any ultrafilter $q \in SP$. Then (G, τ_q) is a particular example to Problem 3.

3. The group $S(X)$ has been used in [5, Example 6.2] to prove the following statement. Let B be a non-discrete extremally disconnected topological Abelian group. Then there exists an extremally disconnected topological group G with distinct left and right uniformities such that B is topologically isomorphic to some open subgroup of G .

4. A group G with a topology τ is called paratopological if the multiplication $(x, y) \mapsto xy$ is jointly continuous in G . By [8], every maximal paratopological group is a topological group. Let G be an extremally disconnected paratopological group. Is G a topological group?

REFERENCES

- [1] Arhangel'skii A.V., *On topological and algebraic properties of extremally disconnected semitopological groups*, Comment. Math. Univ. Carolinae **42.4** (2000), 803–810.
- [2] Arhangel'skii A.V., *Groups topologiques extremement discontinus*, C.R. Acad. Sci. Paris **265** (1967), 822–825.
- [3] Hindman N., Strauss D., *Algebra in the Stone-Ćech Compactification: Theory and Applications*, Walter de Gruyter, Berlin, 1998.
- [4] Protasov I.V., *Filters and topologies on semigroups* (in Russian), Mat. Stud. **3** (1994), 15–28.
- [5] Protasov I.V., *Maximal topologies on groups*, Siberian Math. J. **39** (1998), 1184–1194.
- [6] Protasov I.V., *On maximal topologies on groups*, Visn. Kyiv. Univ. Ser. Fiz-mat. nauk, no. 3 (1998), 251–253.
- [7] Protasov I.V., *Indecomposable topologies on groups*, Ukrainian Math. J. **50** (1998), 1879–1887.
- [8] Protasov I.V., *Extremal topologies on groups*, Mat. Stud. **15** (2001), 9–22.
- [9] Zelenyuk E.G., *Extremal ultrafilters and topologies on groups* (in Russian), Mat. Stud. **14** (2000), 121–140.

DEPARTMENT OF CYBERNETICS, KIEV UNIVERSITY, VOLODIMIRSKA 64, KIEV 01033, UKRAINE

(Received October 16, 2001, revised November 25, 2001)