

Frédéric Mynard

More on strongly sequential spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 3, 525--530

Persistent URL: <http://dml.cz/dmlcz/119342>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

More on strongly sequential spaces

FRÉDÉRIC MYNARD

Abstract. Strongly sequential spaces were introduced and studied to solve a problem of Tanaka concerning the product of sequential topologies. In this paper, further properties of strongly sequential spaces are investigated.

Keywords: sequential, strongly sequential, Fréchet, Tanaka topology

Classification: 54B10, 54D55, 54A20, 54B30

Strongly sequential spaces were introduced in [10] in order to solve a problem of Tanaka [12] of characterizing topologies whose product with every metrizable topology is sequential. In this paper, we identify a sequence $(x_n)_\omega$ with the corresponding filter (generated by $\{x_n : n \geq k\}_{k \in \omega}$) and a decreasing sequence of subsets with the filter it generates. In this way, the definition of the adherence of a filter ⁽¹⁾

$$\text{adh } \mathcal{H} = \bigcup_{\mathcal{F} \# \mathcal{H}} \lim \mathcal{F},$$

applies to sequences and decreasing sequences of subsets. Let cl_{Seq} denote the (idempotent) sequential closure ⁽²⁾ and let $\text{adh}_{\text{Seq}} \mathcal{H}$ be the union of limits of sequences $(x_n)_\omega$ that meshes with the filter \mathcal{H} .

A topology (more generally a convergence) is *strongly sequential* if

$$\text{adh } \mathcal{H} \subset \text{cl}_{\text{Seq}}(\text{adh}_{\text{Seq}} \mathcal{H}),$$

I am deeply indebted to professor S. Dolecki whose observations and suggestions are not only at the origin of this note but also have importantly improved its content. I would also like to thank professor Y. Tanaka for many valuable comments (in [13] and [14]) about preliminary versions of the present paper and about [10].

¹ Two families \mathcal{A} and \mathcal{B} of subsets *mesh*, in symbol $\mathcal{A} \# \mathcal{B}$, if $A \cap B \neq \emptyset$ for every $A \in \mathcal{A}$ and $B \in \mathcal{B}$.

² Let $\text{adh}_{\text{Seq}}^0 A = A$ and let $\text{adh}_{\text{Seq}}^1 A = \text{adh}_{\text{Seq}} A$ be the union of limits of sequences of A . If α is an ordinal number, let $\text{adh}_{\text{Seq}}^\alpha A = \text{adh}_{\text{Seq}}(\bigcup_{\beta < \alpha} \text{adh}_{\text{Seq}}^\beta A)$. For each subset A of X , there exists the least ordinal α for which $\text{adh}_{\text{Seq}}^{\alpha+1} A = \text{adh}_{\text{Seq}}^\alpha A$. This set is the *sequential closure* $\text{cl}_{\text{Seq}} A$ of A . The supremum of the above α 's for every subset A is the *sequential order* of the topology (or convergence). The topology is *sequential* if the closure cl coincide with the sequential closure.

for every countably based filter \mathcal{H} such that $\mathcal{H} = \mathcal{H}_{\text{ad}_T}$. The notation $\mathcal{H} = \mathcal{H}_{\text{ad}_T}$ means that \mathcal{H} has a filter-base consisting of sets that are unions of the closure of their points. Of course, this condition is always fulfilled in a T_1 (i.e., points are closed) convergence. In other words, a T_1 topology (convergence) is strongly sequential if whenever a decreasing sequence of subsets $(A_n)_\omega$ accumulates at x , the point x belongs to the sequential closure of the set of limit points of convergent sequences $(x_n)_n$ such that $x_n \in A_n$. One can also say that a T_1 topology (convergence) is strongly sequential if it is sequential and satisfies the following: if $(A_n)_{n \in \omega}$ is a decreasing sequence accumulating at x then $x \in \text{cl}\{y : y_n \rightarrow y; y_n \in A_n\}$. In the sequel, *regular* means regular and T_1 (contrary to [10]), so that the above characterization applies.

Strongly sequential spaces play a role with respect to sequential spaces similar to that played by strongly Fréchet spaces with respect to Fréchet spaces (see [10]) ⁽³⁾.

A topology (convergence) in which $\text{adh}\mathcal{H} \neq \emptyset$ implies that $\text{adh}_{\text{Seq}}\mathcal{H} \neq \emptyset$ for every countably based filter \mathcal{H} is called *Tanaka space* ⁽⁴⁾. Obviously, every strongly sequential space is a Tanaka space. Proposition 1 below shows that the converse is true among regular sequential spaces. On the other hand, Y. Tanaka [13] asked if a regular sequential inner-one A space is strongly sequential. Recall that a topology is *inner-one A* [9] if $\text{adh}(A_n)_\omega \neq \emptyset$ implies that there exists $x_n \in A_n$ such that $\{x_n : n \in \omega\}$ is not closed.

Proposition 1. *Let X be a regular sequential topology. The following are equivalent.*

1. X is strongly sequential;
2. X is a Tanaka topology;
3. X is inner-one A .

Y. Tanaka pointed out to me in [14] that he proved the equivalence between 2 and 3 in 1986.

PROOF: $1 \implies 2 \implies 3$ follows immediately from the definitions.

$3 \implies 1$. Let $(H_n)_\omega$ fulfill $x \in \bigcap_n \text{cl} H_n$. Then $x \in \bigcap_n \text{cl}(H_n \cap W)$ for every closed neighborhood W of x . As X is inner-one A , there exist sequences $(x_n^W)_\omega$ such that $x_n^W \in H_n \cap W$ and $\{x_n^W : n \in \omega\}$ is not closed, hence not sequentially closed, because of sequentiality. Modulo a rearrangement of the terms, $(x_n^W)_\omega$ admits a subsequence $(x_{n_k}^W)_k$ that converges to a point $x_W \in W \cap \text{adh}_{\text{Seq}}(H_n)$. By regularity, $x \in \text{cl}\{x_W : W = \text{cl} W \in \mathcal{N}(x)\}$. By sequentiality, this closure is equal to the sequential closure, so that $x \in \text{cl}_{\text{Seq}} \text{adh}_{\text{Seq}}(H_n)_\omega$. □

³ A topology is *Fréchet* if $\text{cl} = \text{adh}_{\text{Seq}}$ and *strongly Fréchet* if whenever a decreasing sequence $(A_n)_{n \in \omega}$ accumulates at x , there exists a sequence $x_n \in A_n$ that converges to x .

⁴ This property is called “property (C)” in [12].

In [12], Y. Tanaka proved, in the context of regular (T_1) topologies, that a topology whose product with every first-countable space is sequential is necessarily a Tanaka space. Under supplementary assumptions on X , he gave a characterization for the product $X \times Y$ of X with a first-countable space Y to be sequential. In view of [10, Theorem 5.1] (that gives a similar characterization in terms of strong sequentiality without these assumptions on X) and of Proposition 1, he could have dropped the supplementary assumptions on X in [12, Theorem 1.1]. By the way, these assumptions essentially reduce to the fact that X is Fréchet.

Proposition 2. *A regular Tanaka topology in which each point is G_δ , is strongly Fréchet.*

Notice that, although not stated independently, this result is shown along the lines of the proof of the main theorems of [12].

PROOF: Suppose that $x \in \text{cl} A$. Let $(B_n)_\omega$ be a sequence of open sets such that $\bigcap_n B_n = \{x\}$. By regularity, there is a sequence $(F_n)_\omega$ of closed neighborhoods of x such that $F_n \subset B_n$ for each n . It follows that $x \in \text{cl}(F_n \cap A)$, and, as X is a Tanaka topology, there exists a convergent sequence $x_n \in F_n \cap A$. On the other hand, $\lim(x_n)_n \subset \bigcap_n F_n \subset \bigcap_n B_n = \{x\}$, so that the topology is Fréchet. Moreover, a T_1 Tanaka Fréchet topology is strongly Fréchet (see [12] or [10]). \square

Now, if we drop the assumption of regularity, Tanaka and strongly sequential topologies no longer coincide (among sequential spaces).

Example 3. *[A sequential Tanaka topology which is not strongly sequential.]*

Consider the free bisequence

$$x_{n,k} \xrightarrow[k]{} x_n \xrightarrow[n]{} x_\infty,$$

with its usual topology ⁽⁵⁾. Denote $Y_n = \{x_{n,k} : k \in \omega\}$ and consider a family \mathcal{A} of subsets of $\{x_{n,k} : n, k \in \omega\}$ such that $\mathcal{A} \cup \{Y_n : n \in \omega\}$ is maximal almost disjoint (MAD) (see for example [6]). To the already convergent sequences of $Y = \mathcal{A} \cup \{x_{n,k} : n, k \in \omega\} \cup \{x_n : n \in \omega\} \cup \{x_\infty\}$, we add those generated by each $A \in \mathcal{A}$, each of which converges to the respective A seen as an element of Y . Endow Y with the finest topology for which the sequences above converge. This is obviously a sequential topology. It is moreover a Tanaka topology: since all the points but x_∞ are of countable character, it is enough to consider a decreasing sequence (H_p) that fulfills $x_\infty \in \bigcap_p \text{cl} H_p$ and such that every H_p is included in $\{x_{n,k} : n, k \in \omega\}$. If $w_p \in H_p$, then by maximality of $\mathcal{A} \cup \{Y_n : n \in \omega\}$, there

⁵ More precisely, $(x_n)_{n \in \omega}$ is a free sequence converging to x_∞ and for every n , $(x_{n,k})_{k \in \omega}$ is a free sequence converging to x_n . Sequences of the type $(x_{n,k})_{k \in \omega}$ are disjoint. All points $x_{n,k}$ are isolated, while a neighborhood basis of x_n is given by $\{\{x_n\} \cup \{x_{n,k} : k \geq p\} : p \in \omega\}$. Finally, a neighborhood basis for x_∞ is given by $\{\{x_\infty\} \cup \{x_n\} \cup \{x_{n,k} : k \geq m_n\} : p \in \omega, n \geq p, m_n \in \omega\}$.

is a subsequence of $(w_p)_\omega$ that converges to some $A \in \mathcal{A}$. On the other hand, Y is not strongly sequential because the filter \mathcal{H} generated by $(\bigcup_{n \geq m} Y_n)_{m \in \omega}$ verifies $x_\infty \in \text{adh } \mathcal{H}$ and $\text{adh}_{\text{Seq}} \mathcal{H} \subset \mathcal{A}$ which consists of isolated points, so that $x_\infty \notin \text{cl}_{\text{Seq}}(\text{adh}_{\text{Seq}} \mathcal{H})$.

The free bisequence

$$x_{n,k} \xrightarrow[k]{} x_n \xrightarrow[n]{} x_\infty,$$

with its usual topology is not a Tanaka space, hence not strongly sequential, contrary to my claim in [10, p.150]. Indeed, the filter \mathcal{H} generated by $\{x_{n,k} : n \geq m\}_{m \in \omega}$ fulfills $x_\infty \in \text{adh } \mathcal{H}$ but $\text{adh}_{\text{Seq}} \mathcal{H} = \emptyset$ because no sequence of the type $(x_{n_m, k_m})_m$ with $n_m \geq m$ converges. Other examples of non Fréchet strongly sequential topologies can however be provided. Indeed, in view of [10, Theorem 3.1] (that states that a convergence is strongly sequential if and only if its product with every metrizable topology is sequential) and of the classical theorem [8, Theorem 4.2] of Michael that states that a regular sequential topology is locally countably compact if and only if its product with every sequential topology is sequential, we get

Proposition 4. *A regular sequential, locally countably compact topology (convergence) is strongly sequential.*

In particular, each MAD compact topology ⁽⁶⁾ is a regular sequential locally countably compact, hence strongly sequential, topology of sequential order 2 [4, Theorem 3.5], hence not a Fréchet space.

On the other hand, a locally relatively countably compact ⁽⁷⁾ sequential topology need not be strongly sequential, as shows Example 3. Indeed, we only need to find a relatively countably compact neighborhood for x_∞ and $Y \setminus \mathcal{A}$ is such.

Proposition 4 can actually be strengthened. Recall that a topology is q if every point has a sequence (Q_n) of neighborhoods such that $x_n \in Q_n$ implies $\text{adh}(x_n) \neq \emptyset$. A slightly more general class of topologies is that of *bi-quasi- k* spaces. In such spaces, every adherent filter meshes with a countable family (Q_n) such that $x_n \in Q_n$ implies $\text{adh}(x_n) \neq \emptyset$. If this property holds only for countably based filters, the space is called *countably bi-quasi- k* .

Proposition 5. *Every regular sequential countably bi-quasi- k topology (in particular a regular sequential q -topology or a regular sequential bi- k topology) is strongly sequential.*

⁶ that is, the Alexandroff compactification of $N \cup \mathcal{A}$ where \mathcal{A} is a MAD family on a countable set N and where $N \cup \mathcal{A}$ is endowed with the topology in which the neighborhood filter of $A \in \mathcal{A}$ is generated by $\{W : \{A\} \in W, A \setminus W \text{ is finite.}\}$. This space has been called Alexandroff compactification of a Mrówka space, or a Franklin space, or an Isbell space or a ψ -space.

⁷ i.e., each point has a relatively countably compact neighborhood, that is, a neighborhood on which each countably based filter has non-empty adherence (in the whole set).

PROOF: Let \mathcal{H} be a countably based filter (of decreasing base $(H_n)_\omega$) such that $x \in \text{adh } \mathcal{H}$. As X is countably bi-quasi- k , there exists a sequence of sets (Q_n) such that $(Q_n) \# \mathcal{H}$ such that every sequence $x_n \in Q_n$ has non empty adherence. For every n choose $x_n \in H_n \cap Q_n$. The sequence (x_n) has non empty adherence, so that $\{x_n : n \in \omega\}$ is not closed, hence not sequentially closed, by sequentiality. Thus $(x_n)_\omega$ has a convergent subsequence so that $\text{adh}_{\text{Seq}} \mathcal{H} \neq \emptyset$. In view of Proposition 1, X is strongly sequential. \square

I thank the referee for having pointed out to me that Proposition 5 can also be deduced from Proposition 1 and [7, Lemma 9.1]. Proposition 5 answers positively a question of Y. Tanaka [13]: Are regular sequential countably bi- k spaces strongly sequential?

In view of Proposition 5, a non Fréchet strongly sequential topology need not be locally countably compact. Indeed, there exists a regular non Fréchet sequential q -topology which is not locally countably compact. For example, the product of a MAD-compact space and of a regular non locally countably compact first-countable space is a regular q -topology as a product of regular q -topologies which is sequential because the MAD-compact space is locally countably compact. Hence it is strongly sequential (of sequential order at least 2) and not locally countably compact (see also [3, Proposition 13]).

Strongly sequential spaces can be characterized in terms of their product properties: They are exactly the topologies whose product with every metrizable (or bisequential) topology is sequential (equivalently strongly sequential) [10, Theorem 3.1]. On the other hand, strong sequentiality appears in other results on product of sequential spaces, like [2, Theorem 12.1] and [11, Corollary 6.13]. This last example can be combined with Proposition 5 to the effect that

Theorem 6. *The product of a sequential regular bi-quasi- k topology with a strongly Fréchet topology is sequential.*

REFERENCES

- [1] Dolecki S., *Convergence-theoretic approach to quotient quest*, Topology Appl. **73** (1996), 1–21.
- [2] Dolecki S., Mynard F., *Convergence-theoretic mechanism behind product theorems*, Topology Appl. **104** (2000), 67–99.
- [3] Dolecki S., Nogura T., *Two-fold theorem on Fréchetness of products*, Czech. Math. J. **49** (124) (1999), 421–429.
- [4] Dolecki S., Nogura T., *Countably infinite products of sequential topologies*, to appear.
- [5] Engelking R., *Topology*, PWN, 1977.
- [6] Handbook of Set-theoretic Topology, edited by K. Kunen and J.E. Vaughan, North-Holland Publ., 1984.
- [7] Michael E., *A quintuple quotient quest*, Gen. Topology Appl. **2** (1972), 91–138.
- [8] Michael E., *Local compactness and cartesian product of quotient maps and k -spaces*, Ann. Inst. Fourier (Grenoble) **19** (1968), 281–286.

- [9] Michael E., Olson R.C., Siwiec D., *A-spaces and countably biquotient maps*, *Dissertationes Math.* **133** (1976).
- [10] Mynard F., *Strongly sequential spaces*, *Comment. Math. Univ. Carolinae* **41.1** (2000), 143–153.
- [11] Mynard F., *Coreflectively modified continuous duality applied to classical product theorems*, *Applied General Topology* **2** (2) (2002), 119–154.
- [12] Tanaka Y., *Products of sequential spaces*, *Proc. Amer. Math. Soc.* **54** (1976), 371–375.
- [13] Tanaka Y., *Private communication*, November 2000.
- [14] Tanaka Y., *Private communication*, June 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AUCKLAND, PRIVATE BAG 92019,
AUCKLAND, NEW ZEALAND

E-mail: mynard@olemiss.edu

(Received May 28, 2001, revised August 28, 2001)