

Anna Avallone; Giuseppina Barbieri  
Lyapunov measures on effect algebras

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 44 (2003), No. 3, 389--397

Persistent URL: <http://dml.cz/dmlcz/119396>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Lyapunov measures on effect algebras

ANNA AVALLONE, GIUSEPPINA BARBIERI

*Abstract.* We prove a Lyapunov type theorem for modular measures on lattice ordered effect algebras.

*Keywords:* Lyapunov measures, effect algebras, modular functions

*Classification:* 22B05, 06C15

### 1. Introduction

The celebrated Lyapunov's theorem says that the range of a non-atomic finite dimensional measure  $\mu$  on a  $\sigma$ -algebra is convex. In general, this is not true if  $\mu$  is infinite dimensional. On the other hand, Knowles showed that when  $\mu$  is properly non-injective with values in a locally convex linear space, then its range is still convex. In [11], De Lucia and Wright, after introducing a notion of a convex set, generalize Knowles' result to the case when  $\mu$  is group-valued.

In noncommutative measure theory it is known (see [5, Example 3.7]) that there are examples of nonatomic  $\mathbb{R}^n$ -valued measures on effect algebras which do not have a convex range. Nevertheless, in [5] it is proved (see 3.12) that a Lyapunov type theorem holds for  $\mathbb{R}^n$ -valued modular measures on lattice ordered effect algebras. Moreover, in [2], the result of [11] has been extended to modular functions on complemented lattices. Then a natural question arises:

Is it possible to extend the result of [11] to modular measures on effect algebras?

In this paper we give an affirmative answer to this question, introducing the notion of a pseudo non-injective measure (see Definition 4.1) in an effect algebra which is equivalent to the notion of properly non-injective measures in the Boolean case.

We recall that effect algebras have been introduced by D.J. Foulis and M.K. Bennett in 1994 (see [7]) for modelling unsharp measurement in a quantum mechanical system. They are a generalization of many structures which arise in quantum physics (see [6]) and in Mathematical Economics (see [14] and [9]), in particular of orthomodular lattices in noncommutative measure theory (e.g. see [12]) and MV-algebras in fuzzy measure theory.

## 2. Preliminaries

We will fix some notations. First we will give the definition of a D-poset. Examples of D-posets can be found in [10] and [13].

**Definition 2.1.** Let  $(L, \leq)$  be a partial ordered set (a poset for short). A partial binary operation  $\ominus$  on  $L$  such that  $b \ominus a$  is defined iff  $a \leq b$  is called a *difference* on  $(L, \leq)$  if the following conditions are satisfied for all  $a, b, c \in L$ :

- (1) if  $a \leq b$  then  $b \ominus a \leq b$  and  $b \ominus (b \ominus a) = a$ ,
- (2) if  $a \leq b \leq c$  then  $c \ominus b \leq c \ominus a$  and  $(c \ominus a) \ominus (c \ominus b) = b \ominus a$ .

**Definition 2.2.** Let  $(L, \leq, \ominus)$  be a poset with difference and let 0 and 1 be the least and greatest elements in  $L$ , respectively. The structure  $(L, \leq, \ominus)$  is called a *difference poset* (*D-poset* for short), or a *difference lattice* (*D-lattice* for short) if  $L$  is a lattice.

An alternative structure to a D-poset is that of an effect algebra introduced by Foulis and Bennett in [7]. These two structures, D-posets and effect algebras, are equivalent as shown in [13, Theorem 1.3.4].

We recall that a D-lattice is complete ( $\sigma$ -complete) if every set (countable set) has a supremum and an infimum.

If  $a \in L$ , we set  $a^\perp = 1 \ominus a$ .

We say that  $a$  and  $b$  are orthogonal if  $a \leq b^\perp$  and we write  $a \perp b$ . If  $a \perp b$ , we set  $a \oplus b = (a^\perp \ominus b)^\perp$ . If  $a_1, \dots, a_n \in L$  we define inductively  $a_1 \oplus \dots \oplus a_n = (a_1 \oplus \dots \oplus a_{n-1}) \oplus a_n$  if the right-hand side exists. The sum is independent of any permutation of the elements. We say that  $\{a_1, \dots, a_n\}$  is orthogonal if  $a_1 \oplus \dots \oplus a_n$  exists. We say that a family  $\{a_\alpha\}_{\alpha \in A}$  is *orthogonal* if every finite subfamily is orthogonal. If  $\{a_\alpha\}_{\alpha \in A}$  is orthogonal, we define  $\bigoplus_{\alpha \in A} a_\alpha := \sup\{\bigoplus_{\alpha \in F} a_\alpha : F \subset A \text{ finite}\}$  if the left-hand side exists.

If  $(G, +)$  is an abelian group, a function  $\mu : L \rightarrow G$  is called *modular* if, for every  $a, b \in L$ ,  $\mu(a \vee b) + \mu(a \wedge b) = \mu(a) + \mu(b)$ ;  $\mu$  is called a *measure* if, for every  $a, b \in L$ , with  $a \perp b$ ,  $\mu(a \oplus b) = \mu(a) + \mu(b)$ . It is easy to see that  $\mu$  is a measure iff for every  $a, b \in L$  with  $b \leq a$ ,  $\mu(a \ominus b) = \mu(a) - \mu(b)$ .

A measure  $\mu$  is said to be  $\sigma$ -*additive* if, for every orthogonal sequence in  $L$  such that  $a = \bigoplus_{n \in \mathbb{N}} a_n$  exists,  $\mu(a) = \sum_{n \in \mathbb{N}} \mu(a_n)$ . A measure  $\mu$  is said to be *completely additive* if for every orthogonal family  $\{a_\alpha\}_{\alpha \in A}$  in  $L$  such that  $a = \bigoplus_{\alpha \in A} a_\alpha$  exists, the family  $\{\mu(a_\alpha)\}_{\alpha \in A}$  is summable in  $G$  and  $\mu(a) = \sum_{\alpha \in A} \mu(a_\alpha)$ .

Recall that by 3.1 of [17] every modular function  $\mu : L \rightarrow G$  on any lattice generates a lattice uniformity,  $\mathcal{U}(\mu)$ , i.e. a uniformity which makes  $\wedge$  and  $\vee$  uniformly continuous.

We say that  $\mathcal{U}(\mu)$  is *exhaustive* if every monotone sequence  $\{a_n\}$  is a Cauchy sequence. We say that  $\mathcal{U}(\mu)$  is  $\sigma$ -*order* (*order*) *continuous* if every sequence (net)

$\{a_n\}$  which is order converging to  $a$  is converging to  $a$ . We say that a modular measure is exhaustive,  $\sigma$ -order (order) continuous iff  $\mathcal{U}(\mu)$  is so. By 2.2 of [4], a measure is  $\sigma$ -additive iff it is  $\sigma$ -order continuous.

*Throughout this article,  $(G, +)$  is an abelian topological Hausdorff group which has not  $\mathbb{Z}_2$  as a subgroup,  $L$  is a  $\sigma$ -complete  $D$ -lattice and  $\mu : L \rightarrow G$  is a  $\sigma$ -additive modular measure.*

### 3. Semi-convexity

We shall call  $x \in G$  *infinitely divisible* if for every  $n \in \mathbb{N}$  there exists  $y \in G$  such that  $2^n y = x$ . Since  $\mathbb{Z}_2$  is not a subgroup of  $G$  it is clear that when  $2^n y = x$ ,  $y$  is uniquely determined. In what follows we shall denote such a  $y$  by  $\frac{1}{2^n}x$ . If  $d = \frac{s}{2^n}$  is a dyadic rational number of the real interval  $[0, 1]$  and  $x \in G$  is infinitely divisible, we define  $dx$  to be  $sy$ , where  $y = \frac{1}{2^n}x$ . By [11] the definition of  $dx$  does not depend on the representation of  $d$ . Let  $D$  be the set of dyadic rationals in  $[0, 1]$ . For every infinite divisible  $x \in G$ , let  $g_x : D \rightarrow G$  be defined by  $g_x(d) = dx$  for  $d \in D$ . If  $t \in [0, 1]$  and  $\lim_{d \rightarrow t} g_x(d)$  exists in  $G$ , we define  $tx = \lim_{d \rightarrow t} g_x(d)$ . If  $M \subset G$ ,  $M$  is said to be *convex* if for every  $x, y \in M$  and  $t \in [0, 1]$ ,  $tx, (1-t)y$  exist and  $tx + (1-t)y \in M$ .

**Definition 3.1.** A measure  $\mu$  is said to be *semiconvex* if, for each  $b \in L$ , there exists  $c \in L$  such that  $c \leq b$  and  $\mu(b) = 2\mu(c)$ .

**Lemma 3.2.** *If  $\mu$  is semiconvex, then every element of  $\mu(L)$  is infinitely divisible.*

PROOF: For every  $a \in L$  and  $n \in \mathbb{N}$ , there exists  $b \leq a$  such that  $\mu(a) = 2^n \mu(b)$ . □

**Lemma 3.3.** *Suppose that  $\mu$  is semiconvex. Then for every  $a \in L$  and  $d \in D$ , there exists  $a_d \leq a$  such that  $\mu(a_d) = d\mu(a)$ . Moreover, if  $d_1 < d_2$ , then  $a_{d_1} \leq a_{d_2}$ .*

PROOF: Let  $a \in L$ .

(i) Claim 1: For every  $n \in \mathbb{N}$  there exists an orthogonal family  $\Pi_n = \{a_{n,1}, \dots, a_{n,2^n}\}$  in  $L$  such that  $\bigoplus_{j=1}^{2^n} a_{n,j} = a$  and, for every  $i \in \{1, \dots, 2^n\}$  we have:

- (a)  $2^n \mu(a_{n,i}) = \mu(a)$ ,
- (b)  $a_{n,2i-1} \oplus a_{n,2i} = a_{n-1,i}$ .

This is trivial for  $n = 1$ : Since  $\mu$  is semiconvex, we can choose  $a_{1,1} \leq a$  such that  $2\mu(a_{1,1}) = \mu(a)$ . Let  $a_{1,2} := a \ominus a_{1,1}$ . Then  $a_{1,1} \oplus a_{1,2} = a$  and  $2\mu(a_{1,2}) = 2\mu(a) - 2\mu(a_{1,1}) = \mu(a)$ .

By induction, suppose that Claim 1 holds for  $n \in \mathbb{N}$ . Since  $\mu$  is semiconvex, for every  $i \in \{1, \dots, 2^n\}$  we can find  $a_{n+1,2i-1}, a_{n+1,2i}$  in  $L$  such that  $a_{n+1,2i-1} \oplus a_{n+1,2i} = a_{n,i}$  and  $2\mu(a_{n+1,2i-1}) = 2\mu(a_{n+1,2i}) = \mu(a_{n,i})$ .

Set  $\Pi_{n+1} = \{a_{n+1,1}, a_{n+1,2}, \dots, a_{n+1,2^{n+1}}\}$ . Then  $\Pi_{n+1}$  is orthogonal since  $a = \bigoplus_{i=1}^{2^n} a_{n,i} = \bigoplus_{i=1}^{2^n} (a_{n+1,2i-1} \oplus a_{n+1,2i}) = \bigoplus_{i=1}^{2^{n+1}} a_{n+1,i}$  and for every  $i \in \{1, \dots, 2^{n+1}\}$  we have  $2^{n+1}\mu(a_{n+1,i}) = 2^n\mu(a_{n,i}) = \mu(a)$ .

(ii) Now we obtain a family  $\{b_{n,s} : n \in \mathbb{N}\}$  with  $s \in \{0, 1, \dots, 2^n\}$  such that:

- (1)  $b_{n,0} = 0$  and  $b_{n,2^n} = a$ ,
- (2)  $b_{n,i-1} \leq b_{n,i}$ ,
- (3)  $2^n\mu(b_{n,i}) = i\mu(a)$ ,
- (4) if  $\frac{r}{2^m} = \frac{s}{2^n}$ , then  $b_{m,r} = b_{n,s}$ .

It is sufficient to set  $b_{n,0} = 0$  and, for  $i \in \{1, \dots, 2^n\}$ ,  $b_{n,i} = \bigoplus_{j \leq i} a_{n,j}$ .

(iii) If  $d = \frac{r}{2^m}$ , set  $a_d = b_{m,r}$ . Then by (ii),  $a_d \leq a$  and  $2^m\mu(a_d) = r\mu(a)$ , from which  $\mu(a_d) = d\mu(a)$ . Moreover, by (ii), if  $d_1 < d_2$  then  $a_{d_1} \leq a_{d_2}$ . □

**Lemma 3.4.** *Suppose that  $\mu$  is semiconvex. Then for every  $a \in L$  and for every 0-neighborhood  $W$  in  $G$  there exists  $m \in \mathbb{N}$  such that for every  $p \in D$  with  $p \leq \frac{1}{2^m}$ ,  $p\mu(a) \in W$ .*

PROOF: Let  $a \in L$  and  $W$  be a 0-neighborhood in  $G$ . Since  $\mu$  is semiconvex, we can construct a decreasing sequence  $\{a_n\}$  in  $L$  such that  $a_n \leq a$  and  $2^n\mu(a_n) = \mu(a)$  for every  $n \in \mathbb{N}$ . Let  $b_1 := a \ominus a_1$  and for every  $n \geq 2$ , let  $b_n := a_{n-1} \ominus a_n$ . By 3.3 of [1],  $\{b_n\}$  is orthogonal and for every  $n \in \mathbb{N}$ ,  $2^n\mu(b_n) = 2^n\mu(a_{n-1}) - 2^n\mu(a_n) = 2\mu(a) - \mu(a) = \mu(a)$ . Suppose that for every  $m \in \mathbb{N}$  there exists  $c_m$  such that  $\mu(b_m \wedge c_m) \notin W$ . Since  $\{b_n\}$  is orthogonal,  $\{c_m \wedge b_m\}$  is orthogonal, too. Moreover, by 8.1.2 of [16],  $\mu$  is exhaustive. By 2.4 of [3],  $\mu$  is exhaustive if and only if  $\mu(a_n) \rightarrow 0$  for every orthogonal sequence  $\{a_n\}$  in  $L$ . Therefore, we obtain that  $\lim_m \mu(b_m \wedge c_m) = 0$ , a contradiction. Hence we can choose  $m \in \mathbb{N}$  such that  $\mu(b_m \wedge b) \in W$  for every  $b \in L$ . Set  $p = \frac{r}{2^m}$ , with  $p \leq \frac{1}{2^m}$ . By 3.3, we can find  $c \leq b_m$  such that  $\mu(c) = \frac{r}{2^{n-m}}\mu(b_m)$ . Then  $p\mu(a) = \frac{r}{2^n}\mu(a) = \frac{r}{2^{n-m}}\mu(b_m) = \mu(c) = \mu(c \wedge b_m) \in W$ . □

**Lemma 3.5.** *Suppose that  $\mu$  is semiconvex. Then for every  $a \in L$  and every  $t \in [0, 1]$  there exists  $a_t \leq a$  such that  $t\mu(a)$  is defined and  $t\mu(a) = \mu(a_t)$ . Moreover, the map  $t \mapsto a_t$  is increasing.*

PROOF: We repeat the same argument as in [2]. It follows from 3.3 that there exists a family of elements of  $L$   $\{a_d\}_{d \in D}$  such that  $\mu(a_d) = d\mu(a)$  for each  $d \in D$  and, also, for  $d_1 < d_2$ ,  $a_{d_1} \leq a_{d_2} \leq a$ . Let  $t \in [0, 1] \setminus D$ . We define  $\alpha_t, \beta_t$  by  $\alpha_t = \bigvee \{a_d : d \in D \text{ and } d < t\}$  and  $\beta_t = \bigwedge \{a_d : d \in D \text{ and } d > t\}$ . By using the  $\sigma$ -order continuity of  $\mu$  we find that  $\mu(\alpha_t) = \lim_d \nearrow_t \mu(a_d)$   $\mu(\beta_t) = \lim_d \searrow_t \mu(a_d)$ . Let  $V$  be any symmetric 0-neighbourhood in  $G$ . It follows from the construction and from 3.4 that we can find  $n \in \mathbb{N}$  and  $r \in \{0, 1, \dots, 2^n\}$  such that  $d = \frac{r}{2^n} < t < \frac{r+1}{2^n} = d'$ ,  $\mu(\beta_t) - \mu(\alpha_{d'}) \in V$ ,  $\mu(\alpha_t) - \mu(a_d) \in V$ , and  $\frac{1}{2^n}\mu(a) \in V$ . Then  $(\mu(\beta_t) - \mu(\alpha_t)) \in \mu(a_{d'}) - \mu(a_d) + 2V = \frac{1}{2^n}\mu(a) + 2V \subset 3V$ . Since the symmetric neighbourhoods form a base for 0-neighbourhoods, and since

the topology is Hausdorff,  $\mu(\beta_t) = \mu(\alpha_t)$ . Hence we can define  $a_t$ , for  $t \in [0, 1] \setminus D$  to be  $\alpha_t$ . Then it is clear that  $\mu(a_t) = t\mu(a)$  for each  $t \in [0, 1]$ .  $\square$

**Lemma 3.6.** *Let  $t \in [0, 1]$  and  $\nu_t : L \rightarrow G$  be defined as  $\nu_t(a) = t\mu(a)$ . Then  $\nu_t$  is a modular measure.*

PROOF: Let  $a, b \in L$ .

First suppose  $t = \frac{s}{2^n} \in D$ . By 3.3 we can find  $a_t, b_t \in L$  with  $a_t \leq a, b_t \leq b, 2^n\mu(a_t) = s\mu(a)$  and  $2^n\mu(b_t) = s\mu(b)$ . Then we have  $2^n\mu(a_t \vee b_t) + 2^n\mu(a_t \wedge b_t) = 2^n\mu(a_t) + 2^n\mu(b_t) = s\mu(a) + s\mu(b) = s\mu(a \wedge b) + s\mu(a \vee b)$ , from which  $\nu_t(a \vee b) + \nu_t(a \wedge b) = \nu_t(a) + \nu_t(b)$ .

Now let  $t \notin D$  and choose an increasing sequence  $\{d_n\}$  in  $D$  which converges to  $t$ . Then  $t\mu(a \vee b) + t\mu(a \wedge b) = \lim_n d_n\mu(a \vee b) + \lim_n d_n\mu(a \wedge b) = t\mu(a) + t\mu(b)$ , from which  $\nu_t(a \vee b) + \nu_t(a \wedge b) = \nu_t(a) + \nu_t(b)$ .

In a similar way we prove that  $\nu_t$  is a measure.  $\square$

#### 4. Lyapunov measures

In this section we set

$$I(\mu) = \{a \in L : \mu([0, a]) = \{0\}\}$$

and

$$N(\mu) = \{(a, b) \in L \times L : \mu \text{ is constant on } [a \wedge b, a \vee b]\}.$$

By 3.1 of [17] and 4.3 of [4]  $N(\mu)$  is a congruence relation and the quotient  $\hat{L} = L/N(\mu)$  is a D-lattice. Moreover, the function  $\hat{\mu} : \hat{L} \rightarrow G$  defined as  $\hat{\mu}(\hat{a}) = \mu(a)$  for  $a \in \hat{a} \in \hat{L}$  is trivially a modular measure.

We say that  $\mu$  is *closed* if  $\hat{L}$  is complete with respect to the uniformity  $\mathcal{U}(\hat{\mu})$  generated by  $\hat{\mu}$ .

**Definition 4.1.** We say that  $\mu$  is *pseudo non-injective* if for every  $a \in L \setminus I(\mu)$  there exist  $b, c \in L \setminus I(\mu), b \perp c, b \oplus c \leq a$  and  $\mu(b) = \mu(c)$ .

**Lemma 4.2.** (1)  $\mu$  is exhaustive.

- (2)  $\mu$  is closed iff  $\mu$  is order continuous and  $(\hat{L}, \leq)$  is complete.
- (3) If  $G$  is metrizable, then  $\mu$  is closed.
- (4) If  $\mu$  is order continuous, then  $\mu$  is completely additive.

PROOF: (1) By 8.1.2 of [16], every  $\sigma$ -order continuous lattice uniformity on a  $\sigma$ -complete lattice is exhaustive.

(2) By (1) and 1.2.6 of [16], the Hausdorff uniformity  $\mathcal{U}(\hat{\mu})$  generated by  $\hat{\mu}$  on  $\hat{L}$  is exhaustive. Then, by 6.3 of [16],  $(\hat{L}, \mathcal{U}(\hat{\mu}))$  is complete iff  $\mathcal{U}(\hat{\mu})$  is order continuous and  $(\hat{L}, \leq)$  is complete. Therefore, if  $\mu$  is closed, we have that  $(\hat{L}, \leq)$  is complete and  $\hat{\mu}$  is order continuous, too.

Conversely, if  $(\hat{L}, \leq)$  is complete and  $\mu$  is order continuous, then  $\hat{\mu}$  is order continuous by 7.1.9 of [16], and therefore  $\mu$  is closed.

(3) Since  $G$  is metrizable,  $\mathcal{U}(\mu)$  is metrizable and, by (1), it is exhaustive. By 8.1.4 of [16] (see also 3.5 and 3.6 of [17]), we get that  $(L, \leq)$  is complete and  $\mu$  is order continuous. By 7.1.9 of [16],  $(\bar{L}, \leq)$  is complete, too. Hence  $\mu$  is closed by (2).

(4) Let  $\{a_\alpha\}_{\alpha \in A}$  be an orthogonal family in  $L$  such that  $a = \sup\{\bigoplus_{\alpha \in F} a_\alpha : F \subset A \text{ finite}\}$  exists in  $L$ . For every finite  $F \subset A$ , let  $a_F = \bigoplus_{\alpha \in F} a_\alpha$ . Then  $\{a_F : F \subset A, F \text{ finite}\}$  is an increasing net in  $L$ , with  $a = \sup_F a_F$ . Since  $\mu$  is order continuous,  $\mu(a) = \lim_F \mu(a_F)$ . On the other hand  $\mu(a_F) = \sum_{\alpha \in F} \mu(a_\alpha)$ . Thus  $\mu(a) = \sum_{\alpha \in A} \mu(a_\alpha)$ .  $\square$

**Theorem 4.3.** *Let  $L$  be complete and let  $\mu$  be completely additive with  $I(\mu) = \{0\}$ . Then  $\mu$  is semiconvex if and only if  $\mu$  is pseudo non-injective.*

PROOF:  $\Rightarrow$ : Let  $a \in L \setminus I(\mu)$ .

First, suppose  $\mu(a) \neq 0$ . Then there exists  $b \leq a$  such that  $2\mu(b) = \mu(a)$ . Put  $c := a \ominus b$ . Then  $b \perp c$ ,  $b \oplus c = a$  and  $\mu(b) = \mu(c)$ , as  $2\mu(c) = 2\mu(a) - 2\mu(b) = \mu(a)$ . Moreover,  $b, c \notin I(\mu)$ , since  $\mu(b) = \mu(c) \neq 0$ .

Now let  $\mu(a) = 0$ . As  $a \notin I(\mu)$ , there exists  $d \leq a$  such that  $\mu(d) \neq 0$ . From above, there exist  $b, c \in L \setminus I(\mu)$ ,  $b \perp c$ ,  $b \oplus c \leq d$  and  $\mu(b) = \mu(c)$ . Obviously,  $b \oplus c \leq a$ .

$\Leftarrow$ : Let  $a \neq 0$ . We can suppose  $\mu(a) \neq 0$ .

(i) We will show that  $\exists h, 0 < h \leq a$  such that  $\mu(h) = \mu(a)$  and  $\mu(k) \neq 0$  for each  $0 < k \leq h$ .

We can suppose that there exists  $b \leq a$ ,  $b \neq 0$  and  $\mu(b) = 0$ , since otherwise (i) is satisfied with  $h = a$ .

Recall that in a complete D-lattice, if  $\{b_\gamma\}_{\gamma \in \Gamma}$  is an orthogonal family then, for every  $\bar{\gamma} \in \Gamma$ , the set  $\{\gamma \in \Gamma : b_\gamma = b_{\bar{\gamma}}\}$  is finite (see [DP] p.17). Then by Zorn's lemma we can find an orthogonal family  $\{a_\alpha\}_{\alpha \in A}$  with the following properties:

(1) For every  $\alpha \in A$ ,  $a_\alpha \neq 0$  and  $\mu(a_\alpha) = 0$ .

(2) For every finite  $F \subset A$ ,  $\bigoplus_{\alpha \in F} a_\alpha \leq a$ .

(3) If  $\{b_\gamma\}_{\gamma \in \Gamma}$  is an orthogonal family in  $L$  with (1) and (2), then for each  $\bar{\gamma} \in \Gamma$  the set  $\{\alpha \in A : a_\alpha = b_{\bar{\gamma}}\} \neq \emptyset$  and  $\{\gamma \in \Gamma : b_\gamma = b_{\bar{\gamma}}\} \subset \{\alpha \in A : a_\alpha = b_{\bar{\gamma}}\}$ .

Since  $L$  is complete,  $e = \bigoplus_{\alpha \in A} a_\alpha$  is well-defined. By (2) we get  $e \leq a$ . Since  $\mu$  is completely additive we have  $\mu(e) = \sum_{\alpha \in A} \mu(a_\alpha) = 0$ . Put  $h := a \ominus e$ . Then  $h \leq a$  and  $\mu(h) = \mu(a)$ .

We will show that, if  $0 < b \leq h$ ,  $\mu(b) \neq 0$ .

By way of contradiction, assume  $b \in L$ ,  $0 < b \leq h$  and  $\mu(b) = 0$ . Since  $b \leq h \leq e^\perp \leq (\bigoplus_{\alpha \in F} a_\alpha)^\perp$  for each finite  $F \subset A$ , we have, by 4.2 of [7] that every finite subfamily of  $\{a_\alpha\}_{\alpha \in A} \cup \{b\}$  is orthogonal. Moreover, if  $F \subset A$  is finite, we have  $b \bigoplus (\bigoplus_{\alpha \in F} a_\alpha) \leq h \oplus e = (a \ominus e) \oplus e = a$ . Then  $\{a_\alpha\}_{\alpha \in A} \cup \{b\}$  gives a contradiction with (3).

Let  $h$  be as in (i).

We claim that, if  $0 < k \leq h$ , then there exist  $c, d \in L$  such that  $0 < c < d \leq k$  and  $2\mu(c) = \mu(d)$ .

If  $0 < k \leq h$ ,  $\mu(k) \neq 0$  by (i) and, by pseudo non-injectivity, there exist  $b_1, b_2 \in L$ ,  $b_1 \perp b_2$ ,  $b_1 \oplus b_2 \leq k$ ,  $b_1 \neq 0$ ,  $b_2 \neq 0$  and  $\mu(b_1) = \mu(b_2)$ . Then for  $c := b_1$  and  $d := b_1 \oplus b_2$  we have  $0 < c < d \leq k$  as  $b_1$  and  $b_2$  are not zero and  $\mu(d) = \mu(b_1) + \mu(b_2) = 2\mu(c)$ .

(ii) Zorn's lemma ensures the existence of an orthogonal family  $\{d_\alpha\}_{\alpha \in A}$  with the following properties:

- (1) for every  $\alpha \in A$ ,  $d_\alpha \neq 0$  and there exists  $c_\alpha$  such that  $0 < c_\alpha < d_\alpha$  and  $2\mu(c_\alpha) = \mu(d_\alpha)$ ;
- (2) for every finite  $F \subset A$ ,  $\bigoplus_{\alpha \in F} d_\alpha \leq h$ ;
- (3) if  $\{c_\gamma : \gamma \in \Gamma\}$  is an orthogonal family in  $L$  with properties (1) and (2), then for every  $\bar{\gamma} \in \Gamma$  the set  $\{\alpha \in A : d_\alpha = c_{\bar{\gamma}}\} \neq \emptyset$  and  $\{\gamma \in \Gamma : c_\gamma = c_{\bar{\gamma}}\} \subset \{\alpha \in A : d_\alpha = c_{\bar{\gamma}}\}$ .

It is easy to see that the set  $\{c_\alpha : \alpha \in A\}$  is orthogonal. Put  $d = \bigoplus_{\alpha \in A} d_\alpha$  and  $c = \bigoplus_{\alpha \in A} c_\alpha$ . We get  $c \neq 0$ , since  $c_\alpha \neq 0$  for every  $\alpha \in A$ . By (2)  $d \leq h$ . Moreover, as  $\mu(d) = \sum_{\alpha \in A} \mu(d_\alpha) = 2 \sum_{\alpha \in A} \mu(c_\alpha) = 2\mu(c)$  and  $c \leq d$ , we obtain  $c < d$ .

(iii) We will show that  $d = h$ .

Suppose  $d < h$ . Then  $h \ominus d \neq 0$ . From above, there exist  $c_1, c_2 \in L$  with  $0 < c_1 < c_2 \leq h \ominus d$  and  $\mu(c_2) = 2\mu(c_1)$ .

We will check that  $\{d_\alpha\}_{\alpha \in A} \cup \{c_2\}$  has the same properties as  $\{d_\alpha\}_{\alpha \in A}$ .

Since  $c_2 \leq h \ominus d \leq d^\perp \leq (\bigoplus_{\alpha \in F} d_\alpha)^\perp$  for every finite  $F \subset A$ , from 4.2 of [7] it follows that every finite subfamily of  $\{d_\alpha\}_{\alpha \in A} \cup \{c_2\}$  is orthogonal and so, the family is orthogonal. Moreover, if  $F \subset A$  is finite, then  $c_2 \oplus (\bigoplus_{\alpha \in F} d_\alpha) \leq (h \ominus d) \oplus d = h$ . Obviously,  $c_2$  verifies (1). Then  $\{d_\alpha\}_{\alpha \in A} \cup \{c_2\}$  contradicts property (3). Hence  $d = h$ .

It follows that  $\mu(a) = \mu(h) = \mu(d) = 2\mu(c)$ . Therefore  $\mu$  is semiconvex. □

**Theorem 4.4.** *Let  $\mu$  be closed and pseudo non-injective. Then  $\mu(L)$  is convex.*

PROOF: It is clear that we can replace  $L$  by  $L/N(\mu)$  and  $\mu$  by  $\hat{\mu}$ . Then by 4.2 we can suppose  $L$  complete,  $\mu$  completely additive and  $I(\mu) = \{0\}$ . Hence by 4.3  $\mu$  is semiconvex.

Let  $b, c \in L$  and  $t \in [0, 1]$ .

First, suppose  $b \wedge c = 0$ .

By 3.3 there exist  $d, e \in L$  such that  $d \leq b$ ,  $e \leq c$ ,  $\mu(d) = t\mu(b)$  and  $\mu(e) = (1-t)\mu(c)$ . Since  $b \wedge c = 0$ , we have  $d \wedge e = 0$ . It follows that  $t\mu(b) + (1-t)\mu(c) = \mu(d) + \mu(e) = \mu(d \vee e) + \mu(d \wedge e) = \mu(d \vee e)$ .

Now let  $b, c \in L$ . Put  $b_1 := b \ominus (b \wedge c)$  and  $c_1 = c \ominus (b \wedge c)$ . By 1.8.5 of [13] we have  $b_1 \wedge c_1 = 0$ . Then, from above, there exist  $b_2, c_2 \in L$  with  $b_2 \leq b_1$ ,  $c_2 \leq c_1$  and  $t\mu(b_1) + (1-t)\mu(c_1) = \mu(b_2 \vee c_2)$ .



Since  $b = (b \wedge c) \oplus b_1$  and  $c = (b \wedge c) \oplus c_1$ , by 3.6 we obtain  $t\mu(b) = t\mu(b_1) + t\mu(b \wedge c)$  and  $(1-t)\mu(c) = (1-t)\mu(b \wedge c) + (1-t)\mu(c_1)$ . It follows that  $t\mu(b) + (1-t)\mu(c) = \hat{\mu}(b \wedge c) + t\mu(b_1) + (1-t)\mu(c_1) = \mu(b \wedge c) + \mu(b_2 \vee c_2)$ .

We claim that  $b \wedge c \perp b_2 \vee c_2$ . By 1.8.4 of [13] applied with  $c = a \wedge b$ , we obtain  $b_1 \vee c_1 = (b \ominus (b \wedge c)) \vee (c \ominus (b \wedge c)) = (b \vee c) \ominus (b \wedge c)$ , hence  $b_2 \vee c_2 \leq b_1 \vee c_1 \leq 1 \ominus (b \wedge c) = (b \wedge c)^\perp$ .

It follows that  $\mu(b \wedge c) + \mu(b_2 \vee c_2) = \mu((b \wedge c) \oplus (b_2 \vee c_2))$  and, therefore,  $t\mu(b) + (1-t)\mu(c) \in \mu(L)$ .  $\square$

**Corollary 4.5.** *Let  $\mu$  be closed. Then  $\mu$  is pseudo non-injective iff for every  $a \in L$ ,  $\mu([0, a])$  is convex.*

PROOF:  $\Leftarrow$ : From the assumptions we get that  $\mu$  is semiconvex. Hence,  $\hat{\mu}$  is semiconvex, too. Moreover, since  $\mu$  is closed, by 4.2 we have that  $L/N(\mu)$  is complete and  $\hat{\mu}$  is completely additive. Since  $I(\hat{\mu}) = \{\hat{0}\}$ , by 4.3 we have that  $\hat{\mu}$  is pseudo non-injective. We see that  $\mu$  is pseudo non-injective, too. Let  $a \in L \setminus I(\mu)$  and choose  $b \leq a$  such that  $\mu(b) \neq 0$ . Since  $\hat{\mu}$  is pseudo non-injective, there exist  $\hat{c}, \hat{d}; \hat{c}, \hat{d} \neq \hat{0}, \hat{c} \perp \hat{d}, \hat{c} \oplus \hat{d} \leq \hat{b}$  and  $\hat{\mu}(\hat{c}) = \hat{\mu}(\hat{d})$ . Then there exist  $c, d \in L \setminus I(\mu), c \perp d, c \oplus d \leq b \leq a$  and  $\mu(b) = \mu(c)$ .

$\Rightarrow$ : As in 4.4 we can suppose  $L = L/N(\mu)$ . Let  $a \in L$  and denote by  $\mu_a$  the restriction of  $\mu$  to  $[0, a]$ . Observe that  $[0, a]$  is a complete D-lattice and  $\mu_a$  is a  $\sigma$ -order continuous pseudo non-injective modular measure, since  $\mathcal{U}(\mu_a)$  coincides with the restriction of  $\mathcal{U}(\mu)$  to  $[0, a]$  and  $N(\mu_a) = N(\mu) \cap ([0, a] \times [0, a])$ . Hence by 4.4 we have that  $\mu([0, a])$  is convex.  $\square$

## REFERENCES

- [1] Avallone A., *Lattice uniformities on orthomodular structures*, Math. Slovaca **51** (2001), no. 4, 403–419.
- [2] Avallone A., *Liapunov modular functions*, submitted.
- [3] Avallone A., *Separating points of measures on effect algebras*, preprint.
- [4] Avallone A., Basile A., *On a Marinacci uniqueness theorem for measures*, J. Math. Anal. Appl., to appear.
- [5] Barbieri G., *Lyapunov's theorem for measures on D-posets*, Internat. J. Theoret. Phys., to appear.
- [6] Beltrametti E.G., Cassinelli G., *The Logic of Quantum Mechanics*, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [7] Bennett M.K., Foulis D.J., *Effect algebras and unsharp quantum logics. Special issue dedicated to Constantin Piron on the occasion of his sixtieth birthday*, Found. Phys. **24** (1994), no. 10, 1331–1352.
- [8] Bennett M.K., Foulis D.J., *Phi-symmetric effect algebras*, Found. Phys. **25** (1995), no. 12, 1699–172.
- [9] Butnariu D., Klement P., *Triangular Norm-Based Measures and Games with Fuzzy Coalitions*, Kluwer Acad. Publ., 1993.
- [10] Chovanec F., Kopka F., *D-posets*, Math. Slovaca **44** (1994), 21–34.
- [11] de Lucia P., Wright J.D.M., *Group valued measures with the Lyapunoff property*, Rend. Circ. Mat. Palermo (2) **40** (1991), no. 3, 442–452.

- [12] De Simone A., Navara M., Pták P., *On interval homogeneous orthomodular lattices*, Comment. Math. Univ. Carolinae **42** (2001), no. 1, 23–30.
- [13] Dvurečenskij A., Pulmannová S., *New Trends in Quantum Structures*, Mathematics and its Applications, 516, Kluwer Acad. Publ., Dordrecht; Ister Science, Bratislava, 2000.
- [14] Epstein L.G., Zhang J., *Subjective probabilities on subjectively unambiguous events*, Econometrica **69** ((2001)), no. 2, 265–306.
- [15] Knowles G., *Liapunov vector measures*, SIAM J. Control **13** (1975), 294–303.
- [16] Weber H., *Uniform lattices I, II. A generalization of topological Riesz spaces and topological Boolean rings; Order continuity and exhaustivity*, Ann. Mat. Pura Appl. (4) **160** (1991), 347–370 (1992); (4) **165** (1993), 133–158.
- [17] Weber H., *On modular functions*, Funct. Approx. Comment. Math. **24** (1996), 35–52.

UNIVERSITY OF BASILICATA, CONTRADA MACCHIA ROMANA, 85100 POTENZA, ITALY

*E-mail:* avallone@pzuniv.unibas.it

UNIVERSITY OF UDINE, VIA DELLE SCIENZE 208, 33100 UDINE, ITALY

*E-mail:* barbieri@dimi.uniud.it

(Received March 25, 2003, revised May 20, 2003)