

Jan M. Aarts; Robbert J. Fokkink
Mappings on the dyadic solenoid

Commentationes Mathematicae Universitatis Carolinae, Vol. 44 (2003), No. 4, 697--699

Persistent URL: <http://dml.cz/dmlcz/119424>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Mappings on the dyadic solenoid

JAN M. AARTS, ROBBERT J. FOKKINK

Abstract. Answering an open problem in [3] we show that for an even number k , there exist no k to 1 mappings on the dyadic solenoid.

Keywords: Pontryagin duality, k to 1 maps, solenoids

Classification: 22C05,54F15

Suppose that $P = (p_1, p_2, \dots)$ is a sequence of prime numbers. The P -adic solenoid S_P is the inverse limit sequence (S, f_n) where $S \approx \mathbb{R}/\mathbb{Z}$, the circle, and the bonding maps are homomorphisms $f_n(z) = p_n \cdot z \pmod 1$. The P -adic solenoid is a compact abelian group. In case P is a constant sequence of 2's, the inverse limit is called the *dyadic solenoid*, denoted by S_2 . We shall prove the following result.

Theorem 1. *Suppose that $f: S_2 \rightarrow S_2$ is a k to 1 map of the dyadic solenoid. Then k is odd.*

This answers a question in [3] and shows that the result in [7] is correct. The main ingredient in our proof is Scheffer's theorem [6] (see [5] for a recent application of Scheffer's theorem).

Theorem 2. *Suppose that G, H are compact and connected groups and that H is abelian. Suppose that $f: (G, e) \rightarrow (H, e)$ is a continuous map that preserves the unit element. Then f is homotopic to a unique homomorphism. The homotopy preserves the unit element.*

Solenoids have a local product structure of a Cantor set and an arc, [2]. The arc component Γ_e of the unit element e is a dense subgroup that is a $1 - 1$ homomorphic image of \mathbb{R} . The other arc components are translates of Γ .

Proposition 3. *Suppose that $f: S_2 \rightarrow S_2$ is a non-trivial homomorphism. Then f bijectively maps arc-components onto arc-components.*

PROOF: Since f is a homomorphism, it suffices to verify that the restriction to the unit component Γ_e is a bijection. Now Γ_e is an image of \mathbb{R} . A non-trivial homomorphism on \mathbb{R} is of the form $x \rightarrow rx$ for $r \neq 0$. In particular, it is a bijection. \square

Proposition 4. *Suppose that $f: S_2 \rightarrow S_2$ is not homotopic to a constant function. Then f maps arc-components onto arc-components.*

PROOF: By composing f with a translation, if necessary, we may assume that f preserves the unit element. By Scheffer’s theorem, f is homotopic to a non-trivial homomorphism h . The difference map $h - f: S_2 \rightarrow S_2$ has a compact image that is contained in Γ_e . So $f(x) = h(x) + t(x)$ for some $t(x)$ in a compact subset of Γ_e . Since h maps arc-components onto arc-components so does f . □

Under Pontryagin duality, the category of compact abelian groups is contravariantly equivalent to the category of discrete groups. The Pontryagin dual of the dyadic solenoid S_2 is isomorphic to the additive group $Q_2 = \{\frac{k}{2^n}: k \in \mathbb{Z}, n \geq 0\}$, see [4]. Each non-zero element of Q_2 has a unique representation $\frac{k}{2^n}$ for an odd number k and a non-negative integer n .

Lemma 5. *Suppose that $f: S_2 \rightarrow S_2$ is not homotopic to a constant map and that Γ is an arc-component. Then $f^{-1}(\Gamma)$ consists of an odd number of arc-components.*

PROOF: As S_2 is homogeneous, we may assume that Γ is the component of e . By the corollary, $f^{-1}(\Gamma)$ is a collection of arc-components that is necessarily the same for all mappings in the homotopy class of f . By Scheffer’s theorem, we may assume that f is a homomorphism and we see that the number of components in $f^{-1}(\Gamma)$ is the same for every possible choice of Γ . Consider the dual homomorphism $\hat{f}: Q_2 \rightarrow Q_2$. It is determined by the value $\hat{f}(1) = \frac{k}{2^n}$, for some odd number k . The image of \hat{f} is a subgroup of odd index k . By the contravariance of Pontryagin duality, the kernel of f is a subgroup of odd order k . The number of elements in the kernel is equal to the number of arc components by Proposition 3. □

Recall that $f: R \rightarrow R$ has a *proper* local maximum in c if there is an open interval I such that $c \in I$ and $f(x) < f(c)$ for all $c \neq x \in I$. A proper local minimum is defined likewise. A proper local extreme is either a maximum or a minimum. The value $f(c)$ is called a proper extreme value.

Proposition 6. *Suppose $f: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous map with finite fibers. Then the set of proper extreme values of f is countable.*

PROOF: It suffices to show that the set of proper local maxima is countable. As the fibers of f are finite, f has a proper local maximum in x whenever it has a local maximum in x . For each proper local maximum x , select an interval $I(x)$ with rational endpoints as in the definition of proper local maximum. Note that $I(x) \neq I(y)$ whenever $x \neq y$. The claim now follows as there are only countably many intervals with rational end points. □

Lemma 7. *Suppose that $f: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous surjection with finite fibers. Then the parity of $f^{-1}(z)$ is odd for each z that is not a proper extreme value.*

PROOF: Suppose z is not a proper extreme value of f . The graph of $y = f(x)$ intersects the horizontal line $y = z$ transversally, in finitely many points. As f is a surjection, we have $\lim_{x \rightarrow \infty} f(x) = \infty$ and $\lim_{x \rightarrow -\infty} f(x) = -\infty$, or the other way around. \square

PROOF OF THEOREM 1: Suppose that $f: S_2 \rightarrow S_2$ is a continuous k to 1 map. In particular, f is a surjection so it is not homotopic to a constant map. Without loss of generality we may assume that $f(e) = e$. By Lemma 5, $f^{-1}(\Gamma_e)$ consists of an odd number of arc-components. Each of these components is an image of the real line and is mapped surjectively onto Γ_e . As each of these maps can be lifted to \mathbb{R} , see [1], this results in an odd number of maps from the real line onto itself. By Lemma 7, outside a countable set of extreme values, each of these maps has fibers of odd parity. Now the sum of an odd number of odd numbers is odd, so k has to be odd. \square

REFERENCES

- [1] Aarts J.M., *The structure of orbits in dynamical systems*, Fund. Math. **129** (1988), 39–58.
- [2] Aarts J.M., Fokkink R.J., *The classification of solenoids*, Proc. Amer. Math. Soc. **111** (1991), 1161–1163.
- [3] Charatonik J.J., Covarrubias P.P., *On covering mappings on solenoids*, Proc. Amer. Math. Soc. **130** (2002), 2145–2154.
- [4] Hewitt E., Ross K.A., *Abstract Harmonic Analysis*, Vol. I, Die Grundlehren der mathematischen Wissenschaften, Bd. 115, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
- [5] Krupski P., *Means on solenoids*, Proc. Amer. Math. Soc. **131** (2003), 1925–1929.
- [6] Scheffer W.A., *Maps between topological groups that are homotopic to homomorphisms*, Proc. Amer. Math. Soc. **33** (1972), 562–567.
- [7] Zhou Youcheng, *Covering mappings on solenoids and their dynamical properties*, Chinese Sci. Bull. **45** (2000), 1066–1070.

TECHNISCHE UNIVERSITEIT DELFT, FACULTY OF INFORMATION TECHNOLOGY AND SYSTEMS,
P.O. BOX 5031, 2600 GA DELFT, NETHERLANDS

E-mail: j.m.aarts@its.tudelft.nl
r.j.fokkink@its.tudelft.nl

(Received January 30, 2003)