

Přemysl Jedlička

On left distributive left idempotent groupoids

Commentationes Mathematicae Universitatis Carolinae, Vol. 46 (2005), No. 1, 15--20

Persistent URL: <http://dml.cz/dmlcz/119504>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2005

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On left distributive left idempotent groupoids

PŘEMYSL JEDLIČKA

Abstract. We study the groupoids satisfying both the left distributivity and the left idempotency laws. We show that they possess a canonical congruence admitting an idempotent groupoid as factor. This congruence gives a construction of left idempotent left distributive groupoids from left distributive idempotent groupoids and right constant groupoids.

Keywords: groupoids, left distributivity, left idempotency

Classification: 20N02, 08A30

The left self-distributivity identity

$$(LD) \quad x \cdot (y \cdot z) = (x \cdot y) \cdot (x \cdot z)$$

is often studied together with the idempotency identity

$$(I) \quad x \cdot x = x$$

giving left distributive idempotent (LDI) groupoids. However, some structures, for instance the so-called LD-quasigroups [1] (left distributive left quasigroups equipped with another left distributive operation) satisfy, together with left distributivity, a weaker version of idempotency only, called left idempotency:

$$(LI) \quad (x \cdot x) \cdot y = x \cdot y.$$

The first results about left idempotent left distributive groupoids (LDLI) appeared in Kepka [4] where these groupoids were called pseudoidempotent left distributive groupoids. However, the first systematic study of these groupoids seems to have appeared as late as in [2].

In this paper, we study left distributive left idempotent (LDLI) groupoids and show that there exists a canonical congruence that, in fact, is the smallest idempotent congruence. Classes of that congruence are right constant groupoids, *i.e.*, groupoids satisfying the identity

$$(RC) \quad x \cdot z = y \cdot z.$$

This enables us to construct LDLI groupoids starting with an LDI groupoid and a family of right constant groupoids.

Kepka [3] found a decomposition similar to the current one for left symmetric left distributive (LSLD) groupoids. These groupoids form a subvariety of LDLI groupoids given by the identity

$$(LS) \quad x \cdot xy = y$$

and our decomposition is a generalization of the decomposition described for LSLD groupoids.

The smallest idempotent congruence

We begin with technical notes: if not specified differently, each groupoid mentioned here is equipped with the binary operation (\cdot) . The expression abc stands for $a \cdot (b \cdot c)$ and similarly a^k means $a \cdot a^{k-1}$.

Lemma 1. *Let G be an LI groupoid and let a be in G . Then we have, for all a, b in G ,*

$$a^k b = ab \quad \text{and} \quad (a^k)^l = a^{k+l-1}.$$

PROOF: First of all we prove $a^k b = ab$, for all a, b in G . It is evident for $k = 1$ and for $k > 1$ we have

$$a^k b = (a \cdot a^{k-1})b = (a^{k-1} \cdot a^{k-1})b = a^{k-1}b = ab.$$

Now we prove the other result by induction on l . Since it is true for $l = 1$, we continue with $l > 1$:

$$(a^k)^l = (a^k) \cdot (a^k)^{l-1} = a \cdot a^{k+l-2} = a^{k+l-1},$$

and that is what we wanted to prove. \square

Definition 2 ([5]). Let G be an LI groupoid. We define ip_G to be the smallest equivalence relation on G satisfying $(a, a^2) \in \text{ip}_G$.

Lemma 3. *Let G be an LI groupoid. Then, for all a, b in G , the following conditions are equivalent:*

- (i) $(a, b) \in \text{ip}_G$;
- (ii) there exist positive integers k, l satisfying $a^k = b^l$.

PROOF: (i) \Rightarrow (ii): The relation $(a, b) \in \text{ip}_G$ means that there exists a sequence $a = a_0, a_1, \dots, a_n = b$, such that we have $a_i = a_{i-1}^2$ or $a_i^2 = a_{i-1}$, for each $1 \leq i \leq n$. Using induction on n , we show that there exist positive integers k, l satisfying $a^k = b^l$. The claim is evident for $n = 0$. Let us suppose $n \geq 1$. The

induction hypothesis tells us that there exist k', l' satisfying $a^{k'} = a'_{n-1}$. We have two possibilities now:

- for $b^2 = a_{n-1}$ we have $b^{l'+1} = (b^2)^{l'} = a'_{n-1} = a^{k'}$;
- for $b = a^2_{n-1}$, we have $b^{l'} = (a^2_{n-1})^{l'} = (a'_{n-1})^2 = (a^{k'})^2 = a^{k'+1}$.

(ii) \Rightarrow (i): Evident. □

Example 4. The relation ip_G is not a congruence in general, for instance

$$\begin{array}{c|ccc} \cdot & 0 & 1 & 2 \\ \hline 0 & 1 & 0 & 2 \\ 1 & 1 & 0 & 2 \\ 2 & 1 & 2 & 2 \end{array}$$

is a simple LI groupoid with ip_G non-trivial. However, the relation ip_G is a congruence on any LDLI groupoid:

Proposition 5. *For each LDLI groupoid G , the relation ip_G is a congruence and, for any a, b, c in G with $(a, b) \in \text{ip}_G$, we have $ac = bc$.*

PROOF: Consider $(a, b) \in \text{ip}_G$ in G . Then there exist k, l satisfying $a^k = b^l$. Now, for all c in G , we have

$$\begin{aligned} a \cdot c &= a^k \cdot c = b^l \cdot c = b \cdot c, \\ (c \cdot a)^k &= c \cdot a^k = c \cdot b^l = (c \cdot b)^l. \end{aligned}$$

This implies that ip_G is a congruence. □

Note 6. Kepka and Nĕmec [5] proved Proposition 5 for a left cancellative LDLI groupoid. They also proved that, in the case of left cancellative LD groupoids, the LI identity is equivalent to the identity

$$xx \cdot x = xx.$$

This result is not true for non-cancellative ones, as we can see on the following example, which is LD, satisfies the cited identity but it is not LI ($(1 \cdot 1) \cdot 0 \neq 1 \cdot 0$):

$$\begin{array}{c|ccc} \cdot & 0 & 1 & 2 \\ \hline 0 & 1 & 2 & 2 \\ 1 & 1 & 2 & 2 \\ 2 & 2 & 2 & 2 \end{array}$$

It is easy to see that, for any LDLI groupoid G , the factor G/ip_G is LDLI and that the equivalence classes are right constant groupoids. Moreover, two ip_G congruent elements satisfy $a^k = b^l$ for some k and l .

Decomposition of LDLI groupoids

The result of Proposition 5 leads us to introduce the following definition:

Definition 7. A set A is a *connected monounary algebra* if it is equipped with a unary operation α satisfying, for all a, b in A , the relation $\alpha^k(a) = \alpha^l(b)$ for some k, l .

Every right constant groupoid G is equipped with a natural operation $o_G : a \mapsto a^2$ that describes the multiplication on G entirely. On the other hand, we can build, on every monounary algebra, a structure of left idempotent right constant groupoid. We say that a right constant groupoid is *connected* if its corresponding monounary algebra is connected. If G is an LDLI groupoid, all congruence classes of ip_G are connected right constant groupoids, according to Proposition 5. This permits us to find a decomposition of the groupoid G .

Proposition 8. (i) Let H be an LDI groupoid and let A_a , with $a \in H$, be a pairwise disjoint sets. Let $f_{a,b}$ be a mapping from A_b to A_{ab} , for every a, b in H . Let us define the groupoid $B(H, f)$ to be the set $\bigcup_{a \in H} A_a$ with the operation $*$ defined by $x * y = f_{a,b}(y)$, for x in A_a and y in A_b . Then the groupoid $B(H, f)$ is LI. Moreover, the mappings $f_{a,b}$ satisfy the identity

$$(1d) \quad f_{a,bc} \circ f_{b,c} = f_{ab,ac} \circ f_{a,c}$$

for all a, b and c in H if and only if the groupoid $B(H, f)$ is LD.

(ii) Let G be an LDLI groupoid. Then G is equal to $B(G/\text{ip}_G, f)$, where $f_{\bar{a},\bar{b}}(c) = ac$ and \bar{a} stands for the class of ip_G containing a .

PROOF: (i) Let us take arbitrary a, b, c from H , x from A_a , y from A_b and z from A_c . The element $x * x = f_{a,a}(x)$ belongs also to A_a because H is idempotent. Hence we have $(x * x) * y = f_{a,b}(y) = x * y$. For the left distributivity, since we have

$$\begin{aligned} x * (y * z) &= x * f_{b,c}(z) = f_{a,bc}(f_{b,c}(z)) = f_{ab,ac}(f_{a,c}(z)), \\ (x * y) * (x * z) &= f_{a,b}(y) * f_{a,c}(z) = f_{ab,ac}(f_{a,c}(z)), \end{aligned}$$

the groupoid $B(H, f)$ is LD if and only if Condition (1d) is satisfied.

(ii) We remark first that the definition of $f_{\bar{a},\bar{b}}$ depends neither on the choice of a , by Proposition 5, nor on the choice of b . The construction yields an LI groupoid and we want to show that the groupoid $B(G/\text{ip}_G, f)$ is equal to (G, \cdot) . Let us choose arbitrarily a, b in G , c in \bar{a} and d in \bar{b} . Then we have

$$c * d = f_{\bar{a},\bar{b}}(d) = a \cdot d = c \cdot d,$$

which completes the proof. □

Note 9. For all a in G , we have the equality $f_{\bar{a},\bar{a}} = o_G$ on the equivalence class \bar{a} . And when considering any a, b in G , the mapping $f_{\bar{a},\bar{b}}$ has to be a homomorphism:

$$f_{\bar{a},\bar{b}}(o_G(d)) = f_{\bar{a},\bar{b}}(f_{\bar{b},\bar{b}}(d)) = f_{\overline{ab},\overline{ab}}(f_{\bar{a},\bar{b}}(d)) = o_G(f_{\bar{a},\bar{b}}(d))$$

holds for any d in \bar{b} .

In the sequel, each element of the groupoid $B(H, f)$ is denoted by the pair (a, x) with a in H and x in A_a .

Example 10. Let H be an LDI groupoid and let A be a connected right constant groupoid. Let us take, for each a in H , a disjoint copy of A , denoted A_a . We define the mapping $f_{a,b}$ by $d \mapsto o_{H_b}(d)$, d in A_b . Then the groupoid $B(H, f)$ is isomorphic to the product $H \times A$.

We apply the congruence ip_G to get a classification of all nonidempotent simple LDLI groupoids. Although this classification follows directly from the results about simple LD groupoids presented in [5], we show it here because it uses a different approach.

Definition 11 ([5]). The groupoid $\text{Cyc}_r(n)$, with $n \geq 1$, is the set $\{0, 1, \dots, n-1\}$ with the operation $i \cdot j = j - 1$, for $j > 0$, and $i \cdot 0 = n - 1$.

The groupoid $\text{Path}_r(n)$, with $n \geq 1$, is the set $\{0, 1, \dots, n-1\}$ with the operation $i \cdot j = j - 1$, for $j > 0$, and $i \cdot 0 = 0$.

Proposition 12 (Stanovský [6]). *The only simple right constant groupoids are, up to isomorphism, the two-element idempotent right constant groupoid, $\text{Path}_r(2)$ and $\text{Cyc}_r(p)$, for p prime.*

Proposition 13. *The only simple nonidempotent LDLI groupoids are, up to isomorphism, $\text{Path}_r(2)$, and $\text{Cyc}_r(p)$, for p prime.*

PROOF: The congruence ip_G on an LDLI groupoid G is not trivial, unless G is idempotent or G is a connected right constant groupoid. The only nonidempotent simple right constant groupoids are, according to Proposition 12, the groupoids $\text{Path}_r(2)$, and $\text{Cyc}_r(p)$. \square

Acknowledgment. The author wishes to thank Patrick Dehornoy, Aleš Drápal, David Stanovský, the referee and the participants of the KAFKA seminar for their useful comments.

REFERENCES

- [1] Dehornoy P., *Braids and Self Distributivity*, Progress in Mathematics, vol. 192, Birkhäuser, 2000.
- [2] Jedlička P., *Propriété de treillis pour les groupes de Coxeter et les systèmes LDI*, Ph.D. Thesis (in French and Czech), Caen, 2004, 260 pp.

- [3] Kepka T., *Non-idempotent left symmetric left distributive groupoids*, Comment. Math. Univ. Carolinae **35** (1994), no. 1, 181–186.
- [4] Kepka T., *Notes on left distributive groupoids*, Acta Univ. Carolinae Math. Phys. **22.2** (1981), 23–37.
- [5] Kepka T., Němec P., *Selfdistributive groupoids, Part A1: Non-idempotent left distributive groupoids*, Acta Univ. Carolinae Math. Phys. **44.1** (2003), 3–94.
- [6] Stanovský D., *Homomorphic images of subdirectly irreducible algebras*, Contest Thesis, 2001, <http://www.karlin.mff.cuni.cz/~stanovsk/math/publ.htm>.

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF ALGEBRA,
SOKOLOVSKÁ 83, 186 75 PRAGUE 8, CZECH REPUBLIC

E-mail: jedlicka@karlin.mff.cuni.cz

(Received March 2, 2004, revised May 11, 2004)