

János Gerlits; István Juhász; Zoltán Szentmiklóssy  
Two improvements on Tkačenko's addition theorem

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 46 (2005), No. 4, 705--710

Persistent URL: <http://dml.cz/dmlcz/119560>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2005

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Two improvements on Tkačenko’s addition theorem

J. GERLITS, I. JUHÁSZ, Z. SZENTMIKLÓSSY

*Abstract.* We prove that (A) if a countably compact space is the union of countably many  $D$  subspaces then it is compact; (B) if a compact  $T_2$  space is the union of fewer than  $N(\mathbb{R}) = \text{cov}(\mathcal{M})$  left-separated subspaces then it is scattered. Both (A) and (B) improve results of Tkačenko from 1979; (A) also answers a question that was raised by Arhangel’skiĭ and improves a result of Gruenhage.

*Keywords:*  $D$ -space, left separated, compact, countably compact, scattered space, Novák number

*Classification:* 54D55, 54G12, 54A25

### 1. Introduction

We start by recalling a few well-known definitions and introducing some related notation. A space  $X$  is said to be a  $D$ -space if for any neighbourhood assignment  $\phi$  defined on  $X$  there is a closed discrete set  $D \subset X$  such that  $\phi[D] = \bigcup\{\phi(x) : x \in D\} = X$ . For any space  $X$  we set

$$D(X) = \min\{|\mathcal{A}| : X = \bigcup \mathcal{A} \text{ and } A \text{ is a } D\text{-space for each } A \in \mathcal{A}\}.$$

The space  $X$  is called left-separated if there is a well-ordering  $\prec$  on  $X$  such that all initial segments w.r.t.  $\prec$  are closed in  $X$ . Again, we set, for any space  $X$ ,

$$\text{ls}(X) = \min\{|\mathcal{A}| : X = \bigcup \mathcal{A} \text{ and } A \text{ is left-separated for each } A \in \mathcal{A}\}.$$

(Note that both  $D(X)$  and  $\text{ls}(X)$  can be finite.)

It was shown in [7] that left-separated spaces are  $D$ -spaces, hence we have  $D(X) \leq \text{ls}(X)$  for any  $X$ .

In [6], M. Tkačenko proved the following remarkable result: *If  $X$  is a countably compact  $T_3$ -space with  $\text{ls}(X) \leq \omega$  then*

- (i)  $X$  is compact,
- (ii)  $X$  is scattered,
- (iii)  $X$  is sequential.

It is easy to see that if in a scattered compact  $T_2$ -space any countably compact subspace is compact then it is sequential, hence (iii) immediately follows from (i) and (ii), although this is not how (iii) was proved in [6].

The aim of this note is to improve (i) and (ii) as follows.

- (A) *Any countably compact space  $X$  with  $D(X) \leq \omega$  is compact.*
- (B) *If  $X$  is compact  $T_2$  with  $\text{ls}(X) < N(\mathbb{R})$  then  $X$  is scattered.*

Here  $N(\mathbb{R})$  denotes the Novák number of the real line  $\mathbb{R}$ , i.e. the covering number  $\text{cov}(\mathcal{M})$  of the ideal  $\mathcal{M}$  of all meager subsets of  $\mathbb{R}$ .

If  $X$  is any crowded (i.e. dense-in-itself) space and  $Y \subset X$  then we denote by  $N(Y, X)$  the relative Novák number of  $Y$  in  $X$ , that is the smallest number of nowhere dense subsets of  $X$  needed to cover  $Y$ . In particular,  $N(X) = N(X, X)$  is the Novák number of  $X$ .

We should also mention that a weaker version of statement (A), in which  $D(X) < \omega$  is assumed instead of  $D(X) \leq \omega$ , has been established in [3].

## 2. The results

Similarly as in [6], we can actually prove the following higher-cardinal generalization of statement (A) from the introduction.

**Theorem 2.1.** *Let  $\kappa$  be any infinite cardinal and  $X$  be initially  $\kappa$ -compact with  $D(X) \leq \kappa$ . Then  $X$  is actually compact.*

The proof of Theorem 2.1 is based on the following lemma that may have some independent interest in itself.

**Lemma 2.2.** *Let  $X$  be any space and  $Y \subset X$  its  $D$  subspace. If  $\rho$  is a regular cardinal such that  $X$  has no closed discrete subset of size  $\rho$  (i.e.  $\hat{e}(X) \leq \rho$ ), moreover  $\mathcal{U} = \{U_\alpha : \alpha \in \rho\}$  is a strictly increasing open cover of  $X$  then there is a closed set  $Z \subset X$  such that  $Z \cap Y = \emptyset$  and  $Z \not\subset U_\alpha$  for all  $\alpha \in \rho$ .*

PROOF: If there is an  $\alpha \in \rho$  with  $Y \subset U_\alpha$  then  $Z = X - U_\alpha$  is clearly as required. So assume from here on that  $Y \not\subset U_\alpha$  for all  $\alpha \in \rho$ .

For every point  $y \in Y$  let  $\alpha(y)$  be the *minimal* ordinal  $\alpha$  such that  $y \in U_\alpha$  and then consider the neighbourhood assignment  $\phi$  on  $Y$  defined by

$$\phi(y) = U_{\alpha(y)}.$$

Since  $Y$  is a  $D$ -space there is a set  $E \subset Y$ , closed and discrete in  $Y$ , such that  $Y \subset \phi[E]$ . We claim that  $Z = E'$ , the derived set of  $E$ , is now as required.

Indeed,  $Z$  is closed in  $X$  and  $Z \cap Y = \emptyset$  as  $E$  has no limit point within  $Y$ . It remains to show that  $Z \not\subset U_\alpha$  for all  $\alpha \in \rho$ . Assume, indirectly, that  $Z \subset U_\alpha$  for some  $\alpha \in \rho$ . Note first that for any point  $y \in Y \cap U_\alpha$  we have  $\alpha(y) \leq \alpha$ , consequently  $\phi[E \cap U_\alpha] \subset U_\alpha$ . On the other hand,  $Z = E' \subset U_\alpha$  implies that  $E - U_\alpha$  is closed discrete in  $X$ , hence  $|E - U_\alpha| < \rho$  by our assumption. But then

$$\beta = \sup\{\alpha(y) : y \in E - U_\alpha\} < \rho$$

because  $\rho$  is regular, consequently we have

$$Y \subset \phi[E] = \phi[E \cap U_\alpha] \cup \phi[E - U_\alpha] \subset U_\alpha \cup U_\beta = U_{\max\{\alpha, \beta\}},$$

contradicting that no member of  $\mathcal{U}$  covers  $Y$ . □

Now, we can turn to the proof of our theorem.

PROOF OF THEOREM 2.1: It suffices to prove that for no regular cardinal  $\rho$  is there a strictly increasing open cover of  $X$  of the form  $\mathcal{U} = \{U_\alpha : \alpha \in \rho\}$ . For  $\rho \leq \kappa$  this is clear, for  $X$  is initially  $\kappa$ -compact. So assume now that  $\rho > \kappa$ , and assume indirectly that  $\mathcal{U} = \{U_\alpha : \alpha \in \rho\}$  is a strictly increasing open cover of  $X$ . Note also that  $X$  has no closed discrete subset of size  $\rho > \kappa$  because  $X$  is initially  $\kappa$ -compact.

By  $D(X) \leq \kappa$  we have  $X = \bigcup\{Y_\nu : \nu \in \kappa\}$ , where  $Y_\nu$  is a  $D$  subspace of  $X$  for each  $\nu \in \kappa$ . Using Lemma 2.2 then we may define by a straightforward transfinite recursion on  $\nu \in \kappa$  closed sets  $Z_\nu \subset X$  such that for each  $\nu \in \kappa$  we have  $Z_\nu \cap Y_\nu = \emptyset$ ,  $Z_\nu \not\subset U_\alpha$  for all  $\alpha \in \rho$ , moreover  $\nu_1 < \nu_2$  implies  $Z_{\nu_1} \supset Z_{\nu_2}$ . In this we make use of the fact that if  $\nu < \kappa$  and  $\{Z_\eta : \eta \in \nu\}$  is a decreasing sequence of closed sets in  $X$  such that  $\bigcap\{Z_\eta : \eta \in \nu\} \subset U$  for some open  $U \subset X$  then there is an  $\eta \in \nu$  with  $Z_\eta \subset U$  as well, using again the initial  $\kappa$ -compactness of  $X$ .

But then, applying once more that  $X$  is initially  $\kappa$ -compact, we conclude that

$$\bigcap\{Z_\nu : \nu \in \kappa\} \neq \emptyset,$$

contradicting that  $X = \bigcup\{Y_\nu : \nu \in \kappa\}$ . □

It should be noted that in the above result no separation axiom is needed. This is in contrast with Tkačenko's result from [6].

Let us now turn to our second statement (B). Again, we need to first give a preparatory result. For this we recall the cardinal function  $\delta(X)$  that was introduced in [8]:

$$\delta(X) = \sup\{d(S) : S \text{ is dense in } X\}.$$

Let us note here that if  $X$  is a compact  $T_2$ -space then  $\delta(X) = \pi(X)$ , as was shown in [4].

**Lemma 2.3.** *Assume that  $X$  is an arbitrary crowded topological space and  $Y \subset X$  is its left-separated subspace. Then we have*

$$N(Y, X) \leq \delta(X),$$

consequently

$$N(X) \leq \text{ls}(X) \cdot \delta(X).$$

PROOF: We shall prove  $N(Y, X) \leq \delta(X)$  by transfinite induction on the order type of the well-ordering that left-separates  $Y$ . So assume that  $\prec$  is a left-separating well-ordering of  $Y$  such that if  $Z$  is any proper initial segment of  $Y$ , w.r.t.  $\prec$ , then  $N(Z, X) \leq \delta(X)$ .

Let  $G$  be the union of all those open sets  $U$  in  $X$  for which  $Y$  (or more precisely:  $U \cap Y$ ) is dense in  $U$ . Clearly, then  $Y \setminus G$  is nowhere dense in  $X$  and  $Y \cap G$  is dense in  $G$ . The latter then implies

$$d(Y \cap G) \leq \delta(G) \leq \delta(X).$$

On the other hand, since  $\prec$  left-separates  $Y \cap G$ , any dense subset of  $Y \cap G$  must be cofinal in  $Y \cap G$  w.r.t.  $\prec$ , hence we clearly have

$$\text{cf}(Y \cap G, \prec) \leq d(Y \cap G) \leq \delta(X).$$

But any proper  $\prec$ -initial segment of  $Y \cap G$  may be covered by  $\delta(X)$  many nowhere dense sets, by the inductive hypothesis, hence we have

$$N(Y, X) \leq 1 + \delta(X) \cdot \delta(X) = \delta(X),$$

because  $d(X)$  and so  $\delta(X)$  is always infinite by definition. The second part now follows immediately.  $\square$

Note that again absolutely no separation axiom was needed in the above result. However, in the proof of the following theorem the assumption of Hausdorffness is essential.

**Theorem 2.4.** *Let  $X$  be a compact  $T_2$ -space satisfying  $\text{ls}(X) < N(\mathbb{R})$ . Then  $X$  must be scattered.*

PROOF: We actually prove the contrapositive form of this statement. So assume that  $X$  is not scattered, then it is well-known that some closed subspace  $F \subset X$  admits an irreducible continuous closed map  $f : F \rightarrow \mathbb{C}$  onto the Cantor set  $\mathbb{C}$ .

It is also well-known and easy to check that then we have  $\delta(F) = \delta(\mathbb{C}) = \omega$ , moreover  $N(F) = N(\mathbb{C}) = N(\mathbb{R}) > \omega$ . But then from Lemma 2.3 we conclude that

$$\text{ls}(X) \geq \text{ls}(F) = \text{ls}(F) \cdot \omega \geq N(F) = N(\mathbb{R}).$$

□

We would like to mention that 2.3 and 2.4 were motivated by the treatment of Tkačenko's results given in [5]. We also point out that Theorems 2.1 and 2.4 yield a slight strengthening of Tkačenko's theorem in that the  $T_3$  separation axiom may be replaced by  $T_2$  in it. This is new even in the case of left-separated spaces (i.e. the assumption  $\text{ls}(X) = 1$ ) that preceded Tkačenko's result in [2].

**Corollary 2.5.** *Let  $X$  be a countably compact  $T_2$  space that satisfies  $\text{ls}(X) \leq \omega$ . Then  $X$  is compact, scattered, and sequential.*

We finish by formulating a couple of natural problems concerning our results.

**Problem 2.6.** *Is the upper bound  $N(\mathbb{R})$  in Theorem 2.4 sharp? Can it actually be replaced by the cardinality of the continuum (in ZFC, of course)?*

Note that as metric or compact spaces are all  $D$ -spaces, in Theorem 2.4 one clearly cannot replace  $\text{ls}(X)$  with  $D(X)$ . Also, a compact ( $D$ -)space may fail to be sequential. Being left-separated, however, is clearly a hereditary property, hence left-separated spaces are actually hereditary  $D$ -spaces. Thus the following problems may be raised.

**Problem 2.7.** *Is a compact  $T_2$  hereditary  $D$ -space sequential? Does it contain a point of countable character?*

Concerning this problem we note that it follows easily from Theorem 2.1 that a compact  $T_2$ -space  $X$  satisfying  $D(Y) \leq \omega$  for all  $Y \subset X$  has countable tightness.

## REFERENCES

- [1] Arhangel'skii A.V., *D-spaces and covering properties*, Topology Appl. **146–147** (2005), 437–449.
- [2] Gerlits J., Juhász I., *On left-separated compact spaces*, Comment. Math. Univ. Carolinae **19** (1978), 53–62.
- [3] Gruenhage G., *A note on D-spaces*, preprint.
- [4] Juhász I., Shelah S.,  $\pi(X) = \delta(X)$  for compact  $X$ , Topology Appl. **32** (1989), 289–294.
- [5] Simon P., *Left-separated spaces: a comment to a paper of M.G. Tkačenko*, Comment. Math. Univ. Carolinae **20** (1979), 597–603.
- [6] Tkačenko M.G., *O bikompaktah predstavimyh v vide obedidenia schetnogo chisla levych podprostranstv I, II*, Comment. Math. Univ. Carolinae **20** (1979), 361–379 and 381–395.

- [7] van Douwen E., Pfeffer W.F., *Some properties of the Sorgenfrey-line and related spaces*, Pacific J. Math. **81** (1979), 371–377.
- [8] Weston J.H., Shilleto J., *Cardinalities of dense sets*, Gen. Topology Appl. **6** (1976), 227–240.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, P.O.BOX 127, 1364 BUDAPEST, HUNGARY  
*E-mail*: gerlits@renyi.hu

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, P.O.BOX 127, 1364 BUDAPEST, HUNGARY  
*E-mail*: juhasz@renyi.hu

EÖTVÖS LORÁNT UNIVERSITY, DEPARTMENT OF ANALYSIS, 1117 BUDAPEST, PÁZMÁNY PÉTER  
SÉTÁNY 1/A, HUNGARY  
*E-mail*: zoli@renyi.hu

(Received April 7, 2005, revised April 7, 2005)