

Swapan Kumar Ghosh

Intersections of minimal prime ideals in the rings of continuous functions

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 47 (2006), No. 4, 623--632

Persistent URL: <http://dml.cz/dmlcz/119623>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Intersections of minimal prime ideals in the rings of continuous functions

SWAPAN KUMAR GHOSH

*Abstract.* A space  $X$  is called  $\mu$ -compact by M. Mandelker if the intersection of all free maximal ideals of  $C(X)$  coincides with the ring  $C_K(X)$  of all functions in  $C(X)$  with compact support. In this paper we introduce  $\phi$ -compact and  $\phi'$ -compact spaces and we show that a space is  $\mu$ -compact if and only if it is both  $\phi$ -compact and  $\phi'$ -compact. We also establish that every space  $X$  admits a  $\phi$ -compactification and a  $\phi'$ -compactification. Examples and counterexamples are given.

*Keywords:* minimal prime ideal,  $P$ -space,  $F$ -space,  $\mu$ -compact space,  $\phi$ -compact space,  $\phi'$ -compact space, round subset, almost round subset, nearly round subset

*Classification:* Primary 54C40; Secondary 46E25

### 1. Introduction

By a space we always mean a completely regular Hausdorff space. It is well-known that if  $X$  is realcompact, then the intersection of all free maximal ideals of  $C(X)$  coincides with the ring  $C_K(X)$  of all functions in  $C(X)$  with compact support ([1, 8.19]). A space with the latter property is called  $\mu$ -compact by M. Mandelker in 1971 ([5]). A subset  $A$  of  $\beta X$  is called round by M. Mandelker in 1969 if for any zero set  $Z$  of  $X$ ,  $\text{cl}_{\beta X} Z$  is a neighbourhood of  $A$  whenever  $\text{cl}_{\beta X} Z \supseteq A$  ([4, 4]). In 1973, D.G. Johnson and M. Mandelker have shown that for any space  $X$ , there is a smallest  $\mu$ -compact space  $\mu X$  lying between  $X$  and  $\beta X$  ([3, 4.1]). They have also proved that  $\mu X$  is the smallest subspace of  $\beta X$  containing  $X$  for which  $\beta X - \mu X$  is round ([3, 4.3]). We define  $\phi$ -compact spaces in terms of intersections of minimal prime ideals of  $C(X)$ . The class of all  $\phi$ -compact spaces extends the class of all  $\mu$ -compact spaces. We prove that for any space  $X$ , there is a smallest  $\phi$ -compact space  $\phi X$  lying between  $X$  and  $\beta X$ . Mandelker's definition of round subsets of  $\beta X$  characterizes  $P$ -spaces. In fact,  $X$  is a  $P$ -space if and only if every subset of  $\beta X$  is round ([4, 5.6]). The question is what type of subsets of  $\beta X$  characterize  $F$ -spaces? We define almost round subsets of  $\beta X$ . It turns out that a space  $X$  is an  $F$ -space if and only if every subset of  $\beta X$  is almost round. We also establish that  $\phi X$  is the smallest subspace of  $\beta X$  containing  $X$  for which  $\beta X - \phi X$  is almost round. Our motivation to define  $\phi'$ -compact spaces is the theorem in which we show that a space is  $\mu$ -compact if

and only if it is both  $\phi$ -compact and  $\phi'$ -compact. We prove that for any space  $X$ , there is a smallest  $\phi'$ -compact space  $\phi'X$  lying between  $X$  and  $\beta X$ . We define nearly round subsets of  $\beta X$  and similar results as for round and almost round subsets are established. Finally we show that an  $F$ -space  $X$  is a  $P$ -space if and only if every subset of  $\beta X$  is nearly round.

## 2. Maximal, prime and minimal prime ideals

As usual,  $\beta X$  is the Stone-Ćech compactification of  $X$ . There is a one-one correspondence between the points of  $\beta X$  and the maximal ideals of  $C(X)$ , described in the following theorem ([1, 7.3]).

**Theorem 2.1** ([Gelfand-Kolmogoroff]). *The maximal ideals of  $C(X)$  are given by  $M^p = \{f \in C(X) : p \in \text{cl}_{\beta X} Z(f)\}$  ( $p \in \beta X$ ), here  $Z(f) = \{x \in X : f(x) = 0\}$  is the zero-set of  $f$ .*

Also the set  $O^p = \{f \in C(X) : \text{cl}_{\beta X} Z(f) \text{ is a neighbourhood of } p\}$  is an ideal of  $C(X)$ , for each  $p \in \beta X$ .

An ideal  $I$  of  $C(X)$  is called a  $z$ -ideal if  $Z(f) = Z(g)$  and  $f \in I$  implies  $g \in I$ . It is clear that for each  $p \in \beta X$ ,  $M^p$  and  $O^p$  are  $z$ -ideals of  $C(X)$ .

We now write down the following important theorem given in [1, 7.15].

**Theorem 2.2.** *Every prime ideal  $P$  of  $C(X)$  contains  $O^p$  for a unique  $p$  and  $M^p$  is the unique maximal ideal that contains  $P$ .*

It is well-known that  $X$  is an  $F$ -space if and only if  $O^p$  is prime for each  $p \in \beta X$  ([1, 14.25]), and  $X$  is a  $P$ -space if and only if  $O^p = M^p$  for each  $p \in \beta X$  ([1, 14.29]). Clearly every  $P$ -space is an  $F$ -space, the converse is not true. The space  $\beta\mathbb{R} \setminus \mathbb{R}$  is a compact  $F$ -space ([1, 14.27]). It fails to be a  $P$ -space since every compact  $P$ -space is finite ([1, 4k, 2]).

Every  $z$ -ideal in  $C(X)$  is an intersection of prime ideals ([1, 2.8]). Since  $O^p$  is a  $z$ -ideal we have the following theorem.

**Theorem 2.3.** *The ideal  $O^p$  is the intersection of all minimal prime ideals containing it.*

Let  $\mathcal{P}_{\min}(X)$  denote the class of all minimal prime ideals of  $C(X)$ . We define the relation ' $\sim$ ' on  $\mathcal{P}_{\min}(X)$  by  $P \sim Q$  if and only if  $P, Q$  are contained in a same maximal ideal. Obviously ' $\sim$ ' is an equivalence relation on  $\mathcal{P}_{\min}(X)$ . All the minimal prime ideals of  $C(X)$  contained in  $M^p$  (i.e. containing  $O^p$ ) for some  $p \in \beta X$  form an equivalence class which will be denoted by  $E_p$ . We state the following important characterization of minimal prime ideals of  $C(X)$  which is an immediate consequence of [2, Lemma 1.1].

**Theorem 2.4.** *Let  $P$  be a prime ideal of  $C(X)$ . Then  $P$  is minimal if and only if for any  $f \in P$ , there exists  $g \in C(X) - P$  such that  $fg = 0$ .*

**Notations 2.5.** Let  $X \subseteq Y \subseteq \beta X$  and  $p \in \beta X$ . The ideal  $\{f \in C(X) : \text{cl}_{\beta X} Z(f) \text{ is a neighbourhood of } p\}$  of  $C(X)$  will be denoted by  $O_X^p$  and the ideal  $\{f \in C(Y) : \text{cl}_{\beta X} Z(f) \text{ is a neighbourhood of } p\}$  of  $C(Y)$  will be denoted by  $O_Y^p$ .

We note that every minimal prime ideal in  $C(X)$  is a  $z$ -ideal ([1, 14.7]). Now we prove the following theorem.

**Theorem 2.6.** *Let  $X \subseteq Y \subseteq \beta X$  and  $p \in \beta X$ . If  $P_Y$  is a minimal prime ideal of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  and if  $f \in P_Y$  then there exists a minimal prime ideal  $P_X$  of  $C(X)$  with  $P_X \supseteq O_X^p$  such that  $f|_X \in P_X$ . Also if  $P_X$  is a minimal prime ideal of  $C(X)$  with  $P_X \supseteq O_X^p$  and if  $f \in P_X$  with  $f^Y \in C(Y)$  then there exists a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  such that  $f^Y \in P_Y$ , here  $f^Y$  is the continuous extension of  $f$  over  $Y$ .*

**PROOF:** Let  $f \in P_Y$  where  $P_Y$  is a minimal prime ideal of  $C(Y)$  with  $P_Y \supseteq O_Y^p$ . Then there exists  $g \in C(Y)$  such that  $fg = 0$  and  $g \notin P_Y$  (Theorem 2.4). Clearly,  $g \notin O_Y^p$ . Let  $g' = g|_X$ . Then  $Z(g') \subseteq Z(g)$  and hence  $g' \notin O_X^p$ . Let  $f' = f|_X$ . Clearly,  $f'g' = 0$ . Now  $g' \notin O_X^p$  implies that there exists a minimal prime ideal  $P_X$  of  $C(X)$  with  $P_X \supseteq O_X^p$  such that  $g' \notin P_X$ . Thus  $f' = f|_X \in P_X$ .

Conversely let,  $f \in P_X$  with  $f^Y \in C(Y)$  where  $P_X$  is a minimal prime ideal of  $C(X)$  such that  $P_X \supseteq O_X^p$ . Now there exists  $g \in C(X)$  with  $fg = 0$  such that  $g \notin P_X$  (Theorem 2.4). Let  $h = g \wedge 1$ . Since  $g \notin P_X$  and  $P_X$  is a  $z$ -ideal,  $h \notin P_X$ . Clearly  $fh = 0$ . Let  $h^Y$  be the continuous extension of  $h$  over  $Y$ . Then,  $f^Y h^Y = 0$ . We claim that there exists a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  such that  $h^Y \notin P_Y$ . If not, then  $h^Y \in O_Y^p$  and so there is a neighbourhood  $V$  of  $p$  in  $\beta X (= \beta Y)$  such that  $Z(h^Y) \supseteq V \cap Y$  ([1, 7.12(a)]). Thus,  $Z(h) = X \cap Z(h^Y) \supseteq V \cap Y \cap X = V \cap X$  and so,  $h \in O_X^p$  ([1, 7.12(a)]). Hence  $g \in O_X^p$  since  $O_X^p$  is a  $z$ -ideal. This shows that  $g \in P_X$ , a contradiction. So,  $h^Y \notin P_Y$  for some minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  and thus  $f^Y \in P_Y$ . □

### 3. $\phi$ -compact spaces and almost round subsets

Recall the equivalence relation introduced in Section 2. Let us now give the following definition.

**Definition 3.1.** Let  $A \subseteq \beta X$ . A family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$  is said to be adequate for  $A$  if  $\mathcal{F} \cap E_p \neq \emptyset \forall p \in A$ . A space  $X$  is defined to be  $\phi$ -compact if  $\bigcap \mathcal{F} \subseteq C_K(X)$  for every family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$ .

**Examples 3.2.** (a) Every  $F$ -space is  $\phi$ -compact. In fact, if  $X$  is an  $F$ -space then  $E_p = \{O^p\} \forall p \in \beta X$ . So if  $\mathcal{F}$  is a family of minimal prime ideals of  $C(X)$ ,

adequate for  $\beta X - X$  then  $O^p \in \mathcal{F} \forall p \in \beta X - X$ . Clearly,  $\bigcap \mathcal{F} \subseteq \bigcap_{p \in \beta X - X} O^p = C_K(X)$  and thus  $X$  is  $\phi$ -compact.

(b) Every  $\mu$ -compact space is  $\phi$ -compact (hence every realcompact space is  $\phi$ -compact). In fact, if  $\mathcal{F}$  is any family of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$  then  $\bigcap \mathcal{F} \subseteq \bigcap_{p \in \beta X - X} M^p$ . Now if  $X$  is  $\mu$ -compact then  $\bigcap_{p \in \beta X - X} M^p = C_K(X)$  and thus  $\bigcap \mathcal{F} \subseteq C_K(X)$ . So  $X$  becomes  $\phi$ -compact.

(c) The Tychonoff plank  $T$  is not  $\phi$ -compact. We know that there is only one free maximal ideal, say  $M^t$  in  $C(T)$ . Also  $O^t$  is not prime ([1, 8J, 6]). Thus if  $P$  is any minimal prime ideal of  $C(T)$  with  $P \subseteq M^t$  then  $O^t \not\subseteq P$  and hence  $T$  cannot be  $\phi$ -compact.

Our next theorem shows that every space  $X$  admits a  $\phi$ -compactification.

**Theorem 3.3.** *For every space  $X$ , there is a smallest  $\phi$ -compact space  $\phi X$  lying between  $X$  and  $\beta X$ . So  $X$  is  $\phi$ -compact if and only if  $X = \phi X$ .*

PROOF: Let  $\Phi$  denote the set of all  $\phi$ -compact spaces lying between  $X$  and  $\beta X$ . Clearly  $\Phi \neq \emptyset$  since  $\beta X \in \Phi$ . Let  $\phi X = \bigcap \Phi$ . To complete the theorem we shall show that  $\phi X$  is  $\phi$ -compact. Consider any family  $\mathcal{F}$  of minimal prime ideals of  $C(\phi X)$ , adequate for  $\beta(\phi X) - \phi X (= \beta X - \phi X)$  and suppose  $f \in \bigcap \mathcal{F}$ . Let  $Y \in \Phi$  and  $p \in \beta X - Y$ . Then  $p \in \beta X - \phi X$ . Since  $\mathcal{F}$  is adequate for  $\beta X - \phi X$ , there is a minimal prime ideal  $P_{\phi X}$  of  $C(\phi X)$  in  $\mathcal{F}$  with  $P_{\phi X} \supseteq O^p_{\phi X}$ . So  $f \in P_{\phi X}$ . Clearly  $f \in C^*(\phi X)$  and let  $f^Y$  be the continuous extension of  $f$  over  $Y$ . By Theorem 2.6, there is a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O^p_Y$  such that  $f^Y \in P_Y$ . Thus  $\mathcal{F}' = \{P_Y : P_Y \text{ is a minimal prime ideal of } C(Y) \text{ with } f^Y \in P_Y\}$  is adequate for  $\beta Y - Y$  and  $f^Y \in \bigcap \mathcal{F}'$ . Since  $Y$  is  $\phi$ -compact,  $f^Y \in C_K(Y)$ . So,  $\text{cl}_Y(Y - Z(f^Y))$  is compact and hence so is  $\bigcap_{Y \in \Phi} \text{cl}_Y(Y - Z(f^Y))$ . Clearly,  $\text{cl}_{\phi X}(\phi X - Z(f)) \subseteq \bigcap_{Y \in \Phi} \text{cl}_Y(Y - Z(f^Y))$ . Let  $p \in \bigcap_{Y \in \Phi} \text{cl}_Y(Y - Z(f^Y))$ . Then  $p \in Y \forall Y \in \Phi$  and so  $p \in \phi X$ . Take any neighbourhood  $U$  of  $p$  in  $\phi X$ . Then there is a neighbourhood  $V$  of  $p$  in  $Y$  (where  $Y \in \Phi$ ) such that  $V \cap \phi X = U$ . Also,  $V \cap (Y - Z(f^Y)) \neq \emptyset$ . Thus,  $V \cap (Y - Z(f^Y))$  is a non-void open set in  $Y$ . Since  $\phi X$  is dense in  $Y$ ,  $\phi X \cap V \cap (Y - Z(f^Y)) \neq \emptyset$  i.e.  $U \cap (\phi X - Z(f)) \neq \emptyset$ . So  $p \in \text{cl}_{\phi X}(\phi X - Z(f))$ . Thus,  $\text{cl}_{\phi X}(\phi X - Z(f)) = \bigcap_{Y \in \Phi} \text{cl}_Y(Y - Z(f^Y))$ . Hence  $f \in C_K(\phi X)$  and  $\phi X$  becomes  $\phi$ -compact. □

We now define almost round subsets as follows.

**Definition 3.4.** A subset  $A$  of  $\beta X$  is said to be almost round if  $\bigcap \mathcal{F} \subseteq \bigcap_{p \in A} O^p$  for every family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $A$ .

Obviously  $X$  is  $\phi$ -compact if and only if  $\beta X - X$  is almost round. We also note that the union of any collection of almost round subsets of  $\beta X$  is almost round.

We now prove the following two lemmas.

**Lemma 3.5.** *Let  $X \subseteq Y \subseteq vX$ . Then  $f \in O_X^p$  if and only if  $f^Y \in O_Y^p$  where  $f^Y$  is the continuous extension of  $f$  over  $Y$ .*

PROOF: The lemma follows from the fact that  $\text{cl}_{\beta X} Z(f) = \text{cl}_{\beta X} Z(f^Y)$ . □

**Lemma 3.6.** *Let  $X \subseteq Y \subseteq vX$ . Then  $Y$  is  $\phi$ -compact if and only if  $\beta X - Y$  is almost round (with respect to  $X$ ).*

PROOF: Let  $Y$  be  $\phi$ -compact and let  $\mathcal{F}$  be a family of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - Y$ . Suppose  $f \in \bigcap \mathcal{F}$  and  $f^Y$  is the continuous extension of  $f$  over  $Y$ . If  $p \in \beta X - Y$  then there is a minimal prime ideal  $P_X \in \mathcal{F}$  with  $P_X \supseteq O_X^p$ ,  $\mathcal{F}$  being adequate for  $\beta X - Y$ . So by Theorem 2.6, there is a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  such that  $f^Y \in P^Y$ . Thus  $\mathcal{F}' = \{P_Y : P_Y$  is a minimal prime ideal of  $C(Y)$  with  $f^Y \in P_Y\}$  is adequate for  $\beta X - Y$  and  $f^Y \in \bigcap \mathcal{F}'$ . Since  $Y$  is  $\phi$ -compact,  $f^Y \in C_K(Y)$ . Thus  $f^Y \in O_Y^p \forall p \in \beta X - Y$ . So by Lemma 3.5,  $f \in O_X^p \forall p \in \beta X - Y$ . Consequently,  $\bigcap \mathcal{F} \subseteq \bigcap_{p \in \beta X - Y} O_X^p$  and so  $\beta X - Y$  is almost round.

Conversely let  $\beta X - Y$  be almost round. Suppose  $\mathcal{F}'$  is any family of minimal prime ideals of  $C(Y)$ , adequate for  $\beta Y - Y (= \beta X - Y)$  and suppose  $f \in \bigcap \mathcal{F}'$ . Let  $f_1 = f|_X$  and  $p \in \beta X - Y$ . Since  $\mathcal{F}'$  is adequate for  $\beta X - Y$ , there is a minimal prime ideal  $P_Y \in \mathcal{F}'$  such that  $P_Y \supseteq O_Y^p$ . Also  $f \in P_Y$ . By Theorem 2.6, there is a minimal prime ideal  $P_X$  of  $C(X)$  with  $P_X \supseteq O_X^p$  such that  $f_1 \in P_X$ . Thus  $\mathcal{F} = \{P_X : P_X$  is a minimal prime ideal of  $C(X)$  with  $f_1 \in P_X\}$  becomes adequate for  $\beta X - Y$  and  $f_1 \in \bigcap \mathcal{F}$ . Since  $\beta X - Y$  is almost round,  $f_1 \in O_X^p \forall p \in \beta X - Y$  and so by Lemma 3.5,  $f \in O_Y^p \forall p \in \beta X - Y$ . So  $\bigcap \mathcal{F}' \subseteq \bigcap_{p \in \beta X - Y} O_Y^p = C_K(Y)$  and hence  $Y$  is  $\phi$ -compact. □

**Corollary 3.7.** *For any space  $X$ ,  $\beta X - \phi X$  is almost round.*

We now use Lemma 3.6 to prove the following theorem.

**Theorem 3.8.** *For any space  $X$ ,  $\phi X$  is the smallest subspace of  $\beta X$  containing  $X$  for which  $\beta X - \phi X$  is almost round.*

PROOF: Let  $X \subseteq Y \subseteq \beta X$  such that  $\beta X - Y$  is almost round. Then  $(\beta X - \phi X) \cup (\beta X - Y) = \beta X - (\phi X \cap Y)$  is almost round. Clearly  $X \subseteq \phi X \cap Y \subseteq vX$  and so Lemma 3.6 implies that  $\phi X \cap Y$  is  $\phi$ -compact. Since  $\phi X$  is the smallest  $\phi$ -compact space between  $X$  and  $\beta X$ ,  $\phi X \subseteq \phi X \cap Y$ . So  $\phi X \subseteq Y$  and the theorem follows. □

Almost round subsets characterize  $F$ -spaces in the following way.

**Theorem 3.9.**  *$X$  is an  $F$ -space if and only if every subset of  $\beta X$  is almost round.*

PROOF: The necessity follows from the fact that for an  $F$ -space  $X$ ,  $E_p = \{O^p\} \forall p \in \beta X$ .

To prove the sufficiency let  $p \in \beta X$ . Since  $\{p\}$  is almost round,  $O^p = P$  for any minimal prime ideal  $P$  with  $P \supseteq O^p$ . Thus  $O^p$  is prime and so  $X$  is an  $F$ -space.

Let  $X$  be a  $\phi$ -compact space. If  $\tau : X \rightarrow Y$  is a homeomorphism then  $\tau$  has an extension to a homeomorphism  $\tau_1 : \beta X \rightarrow \beta Y$  such that  $\tau|_{\beta X - X} : \beta X - X \rightarrow \beta Y - Y$  is also a homeomorphism. Also the map  $\psi : C(Y) \rightarrow C(X)$  defined by  $f \rightarrow f \circ \tau$  is an isomorphism. If  $\mathcal{F} = \{P_Y^\alpha : \alpha \in \Lambda\}$  is a family of minimal prime ideals of  $C(Y)$ , adequate for  $\beta Y - Y$  then clearly  $\mathcal{F}_X = \{\psi(P_Y^\alpha) : \alpha \in \Lambda\}$  becomes a family of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$ . It is now easy to see that  $Y$  is  $\phi$ -compact. Hence we have the following theorem. □

**Theorem 3.10.**  *$\phi$ -compactness is a topological property.*

**Example 3.11.** Let  $Y = \beta N - \{p\}$  where  $p \in \beta N - N$ . Then  $Y$  is an  $F$ -space and hence  $\phi$ -compact. The lone free maximal ideal of  $C(Y)$  is  $M_Y^p = \{f \in C(Y) : p \in \text{cl}_{\beta Y} Z(f)\}$ . Clearly  $p \in \text{cl}_{\beta N}(Y - N)$ . Define  $f : N \rightarrow R$  by  $f(n) = \frac{1}{n}$  and suppose  $h = f^\beta|_Y$ . Then  $h \in C(Y)$  and  $Z(h) = Y - N$ . Thus  $h \in M_Y^p$ . Now  $\text{cl}_Y(Y - Z(h)) = \text{cl}_Y N = Y$  which is not compact and so  $h \notin C_K(Y)$ . Hence  $Y$  is not  $\mu$ -compact.

**4.  $\phi'$ -compact spaces and nearly round subsets**

Recall the definition of a family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$  (Definition 3.1). Let us now give the following definition.

**Definition 4.1.** A space  $X$  is said to be  $\phi'$ -compact if for any  $f \in \bigcap_{p \in \beta X - X} M^p$ , there is a family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$  such that  $f \in \bigcap \mathcal{F}$ .

**Example 4.2.** Every  $\mu$ -compact space is  $\phi'$ -compact (hence every realcompact space is  $\phi'$ -compact). In fact, if  $X$  is  $\mu$ -compact and if  $f \in \bigcap_{p \in \beta X - X} M^p$  then  $f \in C_K(X)$  and so  $f$  is in every free minimal prime ideal of  $C(X)$ . So if  $\mathcal{F}$  is the collection of all free minimal prime ideals in  $C(X)$  then  $f \in \bigcap \mathcal{F}$ . Clearly  $\mathcal{F}$  is adequate for  $\beta X - X$ .

The following theorem relates  $\mu$ -compact spaces,  $\phi$ -compact spaces and  $\phi'$ -compact spaces.

**Theorem 4.3.** *A space is  $\mu$ -compact if and only if it is both  $\phi$ -compact and  $\phi'$ -compact.*

PROOF: Necessity follows from 3.2(b) and 4.2.

For sufficiency we assume that  $X$  is both  $\phi$ -compact and  $\phi'$ -compact. Let  $f \in \bigcap_{p \in \beta X - X} M^p$ . Since  $X$  is  $\phi'$ -compact, there is a family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$  such that  $f \in \bigcap \mathcal{F}$ . Now  $\phi$ -compactness of  $X$  implies  $\bigcap \mathcal{F} \subseteq C_K(X)$ . Thus  $f \in C_K(X)$  and so  $X$  is  $\mu$ -compact. □

**Example 4.4.** Recall the space  $Y = \beta N - \{p\}$  where  $p \in \beta N - N$  given in 3.11. The space is  $\phi$ -compact but not  $\mu$ -compact. Hence the space is also not  $\phi'$ -compact by the previous theorem.

**Notations 4.5.** Let  $X \subseteq Y \subseteq \beta X$  and  $p \in \beta X$ . The maximal ideal  $\{f \in C(X) : p \in \text{cl}_{\beta X} Z(f)\}$  of  $C(X)$  will be denoted by  $M_X^p$  and the maximal ideal  $\{f \in C(Y) : p \in \text{cl}_{\beta Y} Z(f)\}$  of  $C(Y)$  will be denoted by  $M_Y^p$ .

In our next theorem we shall show that every space  $X$  admits a  $\phi'$ -compactification.

**Theorem 4.6.** *For any space  $X$ , there is a smallest  $\phi'$ -compact space  $\phi'X$  lying between  $X$  and  $\beta X$ . Thus  $X$  is  $\phi'$ -compact if and only if  $X = \phi'X$ .*

PROOF: Let  $\Phi'$  be the family of all  $\phi'$ -compact spaces lying between  $X$  and  $\beta X$ . Then  $\Phi' \neq \emptyset$  since  $\beta X \in \Phi'$ . Let  $\phi'X = \bigcap \Phi'$ . To prove the theorem we shall show that  $\phi'X$  is  $\phi'$ -compact. So let  $f \in \bigcap_{p \in \beta X - \phi'X} M_{\phi'X}^p$  and let  $p \in \beta X - \phi'X$ . Then there is  $Y \in \Phi'$  such that  $p \in \beta X - Y$ . Now  $f \in C^*(\phi'X)$  and let  $f^Y$  be the continuous extension of  $f$  over  $Y$ . Let  $q \in \beta X - Y$ . Clearly  $q \in \beta X - \phi'X$ . So  $f \in M_{\phi'X}^q$ . Hence  $q \in \text{cl}_{\beta X} Z(f) \subseteq \text{cl}_{\beta X} Z(f^Y)$ . Thus  $f^Y \in M_Y^q$ . So  $f^Y \in \bigcap_{q \in \beta X - Y} M_Y^q$ . Since  $Y$  is  $\phi'$ -compact and  $p \in \beta X - Y$ , there is a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  such that  $f^Y \in P_Y$ . So by Theorem 2.6, there is a minimal prime ideal  $P_{\phi'X}$  of  $C(\phi'X)$  with  $P_{\phi'X} \supseteq O_{\phi'X}^p$  such that  $f \in P_{\phi'X}$ . So  $\mathcal{F} = \{P_{\phi'X} : P_{\phi'X} \text{ is a minimal prime ideal of } C(\phi'X) \text{ with } f \in P_{\phi'X}\}$  is adequate for  $\beta X - \phi'X$  and  $f \in \bigcap \mathcal{F}$ . Thus  $\phi'X$  is  $\phi'$ -compact.  $\square$

We now define nearly round subsets as follows.

**Definition 4.7.** A subset  $A$  of  $\beta X$  is said to be nearly round if  $f \in \bigcap_{p \in A} M^p$  implies  $f \in \bigcap \mathcal{F}$  for some family  $\mathcal{F}$  of minimal prime ideals of  $C(X)$ , adequate for  $A$ .

Obviously  $X$  is  $\phi'$ -compact if and only if  $\beta X - X$  is nearly round. We note that the union of any collection of nearly round subsets of  $\beta X$  is nearly round. We also note that a subset of  $\beta X$  is round if and only if it is both almost round and nearly round.

We now prove the following lemma.

**Lemma 4.8.** *Let  $X \subseteq Y \subseteq vX$ . Then  $Y$  is  $\phi'$ -compact if and only if  $\beta X - Y$  is nearly round (with respect to  $X$ ).*

PROOF: Let  $Y$  be  $\phi'$ -compact and let  $f \in \bigcap_{p \in \beta X - Y} M_X^p$ . Let  $f^Y$  be the continuous extension of  $f$  over  $Y$ . Then  $\text{cl}_{\beta X} Z(f^Y) = \text{cl}_{\beta X} Z(f)$  and thus  $f^Y \in \bigcap_{p \in \beta X - Y} M_Y^p$ . Suppose  $p \in \beta X - Y$ . Now  $\phi'$ -compactness of  $Y$  implies that there is a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^p$  such that



$f^Y \in P_Y$ . So by Theorem 2.6, there is a minimal prime ideal  $P_X$  of  $C(X)$  with  $P_X \supseteq O_X^p$  such that  $f \in P_X$ . Thus  $\mathcal{F} = \{P_X : P_X \text{ is a minimal prime ideal of } C(X) \text{ with } f \in P_X\}$  is adequate for  $\beta X - Y$  and  $f \in \bigcap \mathcal{F}$ . Consequently  $\beta X - Y$  is nearly round.

Conversely let  $\beta X - Y$  be nearly round and let  $f \in \bigcap_{p \in \beta X - Y} M_Y^p$ . Let  $f|_X = g$ . Then  $\text{cl}_{\beta X} Z(f) = \text{cl}_{\beta X} Z(g)$  and so  $g \in \bigcap_{p \in \beta X - Y} M_X^p$ . Let  $q \in \beta X - Y$ . Since  $\beta X - Y$  is nearly round, there is a minimal prime ideal  $P_X$  of  $C(X)$  with  $P_X \supseteq O_X^q$  such that  $g \in P_X$ . Hence by Theorem 2.6, there is a minimal prime ideal  $P_Y$  of  $C(Y)$  with  $P_Y \supseteq O_Y^q$  such that  $f \in P_Y$ . Thus  $\mathcal{F}' = \{P_Y : P_Y \text{ is a minimal prime ideal of } C(Y) \text{ with } f \in P_Y\}$  is adequate for  $\beta X - Y$  and  $f \in \bigcap \mathcal{F}'$ . Thus  $Y$  is  $\phi'$ -compact.  $\square$

**Corollary 4.9.** *For any space  $X$ ,  $\beta X - \phi'X$  is nearly round.*

We now use Lemma 4.8 to prove the following theorem.

**Theorem 4.10.** *For any space  $X$ ,  $\phi'X$  is the smallest subspace of  $\beta X$  containing  $X$  for which  $\beta X - \phi'X$  is nearly round.*

PROOF: Let  $X \subseteq Y \subseteq \beta X$  such that  $\beta X - Y$  is nearly round. Then  $(\beta X - \phi'X) \cup (\beta X - Y) = \beta X - (\phi'X \cap Y)$  is nearly round. Clearly  $X \subseteq \phi'X \cap Y \subseteq vX$  and so by Lemma 4.8,  $\phi'X \cap Y$  is  $\phi'$ -compact. Since  $\phi'X$  is the smallest  $\phi'$ -compact space between  $X$  and  $\beta X$ ,  $\phi'X \subseteq \phi'X \cap Y$ . So  $\phi'X \subseteq Y$  and the proof is complete.  $\square$

The following theorem gives a necessary and sufficient condition for an  $F$ -space to be a  $P$ -space.

**Theorem 4.11.** *An  $F$ -space  $X$  is a  $P$ -space if and only if every subset of  $\beta X$  is nearly round.*

PROOF: Let  $X$  be a  $P$ -space and  $A \subseteq \beta X$ . Suppose  $f \in \bigcap_{p \in A} M^p$ . Then  $f \in \bigcap_{p \in A} O^p$ . Thus  $\mathcal{F} = \{O^p : p \in A\}$  is a family of minimal prime ideals of  $C(X)$ , adequate for  $A$  with  $f \in \bigcap \mathcal{F}$ . So  $A$  is nearly round.

Conversely let  $X$  be an  $F$ -space and every subset of  $\beta X$  be nearly round. Let  $p \in \beta X$  and suppose  $f \in M^p$ . Since  $\{p\}$  is nearly round there is a minimal prime ideal  $P$  of  $C(X)$  with  $P \supseteq O^p$  such that  $f \in P$ . Also since  $X$  is an  $F$ -space,  $P = O^p$  and thus  $f \in O^p$ . So  $O^p = M^p$  and hence  $X$  is a  $P$ -space.

Let  $X$  be a  $\phi'$ -compact space. If  $\tau : X \rightarrow Y$  is a homeomorphism then  $\tau$  has an extension to a homeomorphism  $\tau_1 : \beta X \rightarrow \beta Y$  such that  $\tau_1|_{\beta X - X} : \beta X - X \rightarrow \beta Y - Y$  is also a homeomorphism. Also the map  $\psi : C(Y) \rightarrow C(X)$  defined by  $f \rightarrow f \circ \tau$  is an isomorphism. If  $f$  is in the intersection of all free maximal ideals of  $C(Y)$  then  $\psi(f)$  is in the intersection of all free maximal ideals of  $C(X)$ . Now  $\phi'$ -compactness of  $X$  implies that there is a family  $\mathcal{F}_X = \{P_X^\alpha : \alpha \in \Lambda\}$  of minimal prime ideals of  $C(X)$ , adequate for  $\beta X - X$  with  $\psi(f) \in \bigcap \mathcal{F}_X$ . Then  $\mathcal{F}_Y = \{\psi^{\leftarrow}(P_X^\alpha) : \alpha \in \Lambda\}$  becomes a family of minimal prime ideals of  $C(Y)$

adequate for  $\beta Y - Y$  and  $f \in \bigcap \mathcal{F}_Y$ . Thus  $Y$  is also  $\phi'$ -compact. So we have the following theorem. □

**Theorem 4.12.**  *$\phi'$ -compactness is a topological property.*

**Notation 4.13.** Let  $\omega_1$  denote the space of all countable ordinals. Let  $T^* = (\omega_1 + 1) \times (\omega_0 + 1)$  and  $T = T^* - \{(\omega_1, \omega_0)\}$  be the Tychonoff plank.

Let us denote for computational convenience,  $(\alpha, \omega_1) \times \{n\}$  ( $(\alpha, \omega_1] \times \{n\}$ ) by  $(\alpha, \{n\})$  ( $(\alpha, \{n\}]$ ), respectively, where  $\alpha \leq \omega_1$  and  $n \in (\omega_0 + 1)$ .

**Lemma 4.14.** *For each  $f \in M^t - O^t$ , there exists  $g \notin O^t$  such that  $fg = 0$  where  $t = \{(\omega_1, \omega_0)\}$ .*

PROOF: Since  $f \in M^t$ , i.e.  $t \in \text{cl}_{\beta T} Z(f)$ , every neighbourhood of  $t$  must meet  $Z(f)$ . Also  $f \notin O^t$  and so  $\text{cl}_{\beta T} Z(f)$  is not a neighbourhood of  $t$ . Now any neighbourhood of  $t$  is of the form  $(\alpha, \omega_1] \times N'$ , where  $N' \subseteq \omega_0 + 1$ ,  $\alpha \leq \omega_1$  and  $(\omega_0 + 1) - N'$  is at most a finite set. Thus there exist infinite subsets  $N_1, N_2$  of  $\omega_0$  with  $N_1 \cup N_2 = \omega_0$  and  $\alpha \leq \omega_1$ , such that, for each  $n \in N_1$ ,  $f((\alpha, \{n\})) = 0$  and for each  $n \in N_2$ ,  $f((\alpha, \{n\})) \neq 0$ . The choice of single  $\alpha$  is possible here because of the non-cofinality character of any denumerable subset of  $\omega_1$ . Also  $f((\alpha, \{\omega_0\})) = 0$ . Choose  $g : T \rightarrow \mathbb{R}$  by defining  $g((\alpha, \{n\})) = \frac{1}{n}$ , for each  $n \in N_1$ ,  $g((\alpha, \{n\})) = 0$  for each  $n \in N_2$  and assign 0 on rest of the region. Clearly,  $g$  is continuous in  $[0, \alpha] \times (\omega_0 + 1)$ . Choose  $(\gamma, n) \in (\alpha, \{n\}]$ ,  $n \in \omega_0$ . Then  $(\alpha, \{n\})$  is an open neighbourhood of  $(\gamma, n)$  and  $g((\alpha, \{n\}))$  is either  $= 0$  or  $\frac{1}{n}$ . Thus  $f$  is continuous at  $(\gamma, n)$ . If now  $(\gamma, \omega_0) \in (\alpha, \{\omega_0\})$ , then  $g((\gamma, \omega_0)) = 0$ . Choose any  $\epsilon \geq 0$ . Then there exists  $n \in \omega_0$  such that  $\frac{1}{n} \leq \epsilon$ . Take  $M = (\omega_0 + 1) - \{r \in \omega_0 : r \leq n\}$ . Then  $(\alpha, \omega_1] \times M - \{t\}$  is an open neighbourhood of  $(\gamma, \omega_0)$  and  $g(((\alpha, \omega_1] \times M) - \{t\})$  is contained in  $(-\epsilon, \epsilon)$ . Hence  $g$  is continuous at  $(\gamma, \omega_0)$ . Thus  $g$  is continuous on  $T$ . Also since  $T - Z(g)$  contains  $(\alpha, \omega_1] \times N_1$ ,  $g \notin O^t$ . Clearly,  $fg = 0$ . □

Using the above lemma, we now show that the Tychonoff plank  $T$  is  $\phi'$ -compact but not  $\mu$ -compact.

**Example 4.15.** Since  $T$  is not  $\phi$ -compact (Example 3.2(c)), it is neither  $\mu$ -compact. We now show that  $T$  is  $\phi'$ -compact. So let  $f \in \bigcap_{p \in \beta T - T} M^p$  i.e.  $f \in M^t$ . We have to produce a family  $\mathcal{F}$  of minimal prime ideals of  $C(T)$ , adequate for  $\beta T - T = \{t\}$  such that  $f \in \bigcap \mathcal{F}$ . If  $f \in O^t$ , then it becomes obvious, if not then  $fg = 0$  for some  $g \notin O^t$  by Lemma 4.14. Since  $O^t$  is the intersection of all minimal prime ideals containing it, there is a minimal prime ideal, say  $P$  containing  $O^t$  such that  $g \notin P$ . So  $f \in P$  since  $P$  is prime. Let  $\mathcal{F} = \{P\}$ . Clearly  $\mathcal{F}$  is adequate for  $\beta T - T$  and  $f \in \bigcap \mathcal{F}$ .

## REFERENCES

- [1] Gillman L., Jerison M., *Rings of Continuous Functions*, University Series in Higher Math., Van Nostrand, Princeton, New Jersey, 1960.
- [2] Henriksen M., Jerison M., *The space of minimal prime ideals of a commutative ring*, Trans. Amer. Math. Soc. **115** (1965), 110–130.
- [3] Johnson D.G., Mandelker M., *Functions with pseudocompact support*, General Topology Appl. **3** (1973), 331–338.
- [4] Mandelker M., *Round  $z$ -filters and round subsets of  $\beta X$* , Israel J. Math. **7** (1969), 1–8.
- [5] Mandelker M., *Supports of continuous functions*, Trans. Amer. Math. Soc. **156** (1971), 73–83.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF CALCUTTA, 35, BALLYGUNGE CIRCULAR ROAD, KOLKATA-700019, WEST BENGAL, INDIA

*E-mail:* swapan12345@yahoo.co.in

(Received January 31, 2006, revised April 21, 2006)