

Tran Van An; Luong Quoc Tuyen

Further properties of 1-sequence-covering maps

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 49 (2008), No. 3, 477--484

Persistent URL: <http://dml.cz/dmlcz/119737>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Further properties of 1-sequence-covering maps

TRAN VAN AN\*, LUONG QUOC TUYEN

*Abstract.* Some relationships between 1-sequence-covering maps and weak-open maps or sequence-covering  $s$ -maps are discussed. These results are used to generalize a result from Lin S., Yan P., *Sequence-covering maps of metric spaces*, Topology Appl. **109** (2001), 301–314.

*Keywords:* weak base,  $sn$ -network, sequence-covering, 1-sequence-covering, weak-open,  $\pi$ - $s$ -map

*Classification:* 54C10, 54D65, 54E40, 54E99

### 1. Introduction

To find internal characterizations of certain images of metric spaces is one of central problems in general topology. Arhangel'skii [1] showed that a space is an open compact image of a metric space if and only if it has a development consisting of point-finite open covers. In 1996, Lin [14] introduced the notion of 1-sequence-covering maps and proved that a space is a 1-sequence-covering and  $s$ -image of a metric space if and only if it has a point-countable  $sn$ -network, and a space is a 1-sequence-covering, quotient and  $s$ -image of a metric space if and only if it has a point-countable weak base. Then Lin and Yan [16] proved that every sequence-covering, quotient and  $s$ -image of a locally separable metric space is a local  $\aleph_0$ -space. In that paper they also show the following

**Theorem 1.1.** *Every sequence-covering and compact map of a metric space is a 1-sequence-covering map.*

Recently, Xia [25] introduced the concept of weak-open maps, and by using it, certain  $gf$ -countable spaces are characterized as images of metric spaces under various weak-open maps.  $\pi$ -map is another important map which was introduced by Ponomarev in 1960, and  $\pi$ -images of metric spaces attract attention again in [7], [11], [23].

The purpose of this paper is to establish some relationships between 1-sequence-covering maps and weak-open maps or sequence-covering  $s$ -maps, and also to give a generalization of Theorem 1.1.

---

\* Corresponding author.

We assume that all spaces are  $T_2$ , all maps are continuous and onto,  $\mathbb{N}$  denotes the set of all natural numbers,  $\omega = \mathbb{N} \cup \{0\}$ , and any convergent sequence includes its limit point. Let  $f : X \rightarrow Y$  be a map and  $\mathcal{P}$  be a collection of subsets of  $X$ . We denote  $\text{st}(x, \mathcal{P}) = \bigcup\{P \in \mathcal{P} : x \in P\}$ ,  $\bigcup \mathcal{P} = \bigcup\{P : P \in \mathcal{P}\}$ ,  $\bigcap \mathcal{P} = \bigcap\{P : P \in \mathcal{P}\}$ ,  $f(\mathcal{P}) = \{f(P) : P \in \mathcal{P}\}$ .

**Definition 1.2.** Let  $X$  be a space and  $P \subset X$ .

- (1) A sequence  $\{x_n\}$  in  $X$  is called *eventually* in  $P$ , if  $\{x_n\}$  converges to  $x$ , and there exists  $m \in \mathbb{N}$  such that  $\{x\} \cup \{x_n : n \geq m\} \subset P$ .
- (2)  $P$  is called a *sequential neighborhood* of  $x$  in  $X$  [4], if whenever  $\{x_n\}$  is a sequence converging to  $x$  in  $X$ , then  $\{x_n\}$  is eventually in  $P$ .
- (3)  $X$  is a *sequential space* [4], if whenever  $A$  is a non closed subset of  $X$ , then there is a sequence in  $A$  converging to a point not in  $A$ .

**Definition 1.3.** Let  $\mathcal{P}$  be a collection of subsets of  $X$ .

- (1)  $\mathcal{P}$  is *point-countable*, if each point  $x \in X$  belongs to only countably many members of  $\mathcal{P}$ .
- (2)  $\mathcal{P}$  is a *network at  $x$  in  $X$* , if  $x \in P$  for every  $P \in \mathcal{P}$ , and whenever  $x \in U$  with  $U$  open in  $X$ , then  $x \in P \subset U$  for some  $P \in \mathcal{P}$ .

**Definition 1.4.** Let  $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$  be a cover of a space  $X$ . Assume that  $\mathcal{P}$  satisfies the following (a) and (b) for every  $x \in X$ .

- (a)  $\mathcal{P}_x$  is a network at  $x$ .
  - (b) If  $P_1, P_2 \in \mathcal{P}_x$ , then there exists  $P \in \mathcal{P}_x$  such that  $P \subset P_1 \cap P_2$ .
- (1)  $\mathcal{P}$  is a *weak base* of  $X$  [1], if for  $G \subset X$ ,  $G$  is open in  $X$  if and only if for every  $x \in G$ , there exists  $P \in \mathcal{P}_x$  such that  $P \subset G$ ;  $\mathcal{P}_x$  is said to be a weak neighborhood base at  $x$ .
  - (2)  $\mathcal{P}$  is an *sn-network* for  $X$  [14], if every element of  $\mathcal{P}_x$  is a sequential neighborhood of  $x$  for every  $x \in X$ ;  $\mathcal{P}_x$  is said to be an *sn-network* at  $x$ .
  - (3) A space  $X$  is *gf-countable* [1] (resp., *snf-countable* [6]), if  $X$  has a weak base (resp., *sn-network*)  $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$  such that  $\mathcal{P}_x$  is countable for every  $x \in X$ .

*Remark 1.5* ([5]). (1) Weak base  $\implies$  *sn-network*, so *gf-countable*  $\implies$  *snf-countable*.

- (2) In a sequential space, weak bases  $\iff$  *sn-networks*, so *gf-countable*  $\iff$  sequential and *snf-countable*.

**Definition 1.6.** Let  $f : X \rightarrow Y$  be a map.

- (1)  $f$  is a *weak-open* map [25], if there exists a weak base  $\mathcal{B} = \bigcup\{\mathcal{B}_y : y \in Y\}$  for  $Y$ , and for  $y \in Y$ , there exists  $x_y \in f^{-1}(y)$  such that for each open neighborhood  $U$  of  $x_y$ ,  $B_y \subset f(U)$  for some  $B_y \in \mathcal{B}_y$ .

- (2)  $f$  is a 1-sequence-covering map [14], if for each  $y \in Y$ , there is  $x_y \in f^{-1}(y)$  such that whenever  $\{y_n\}$  is a sequence converging to  $y$  in  $Y$ , there is a sequence  $\{x_n\}$  converging to  $x_y$  in  $X$  with  $x_n \in f^{-1}(y_n)$  for every  $n \in \mathbb{N}$ .
- (3)  $f$  is a sequence-covering map [20], if every convergent sequence of  $Y$  is the image of some convergent sequence of  $X$ .
- (4)  $f$  is a quotient map [3], if whenever  $f^{-1}(U)$  is open in  $X$ , then  $U$  is open in  $Y$ .
- (5)  $f$  is an  $s$ -map [8] (resp., a Lindelöf map [22], a compact map), if  $f^{-1}(y)$  is separable (resp., Lindelöf, compact) for each  $y \in Y$ .

**Definition 1.7.** Let  $f : (X, d) \rightarrow Y$  be a map, and  $(X, d)$  be a metric space.

- (1)  $f$  is a  $\pi$ -map [1], if for each  $y \in Y$  and its open neighborhood  $U$  in  $Y$ ,  $d(f^{-1}(y), X \setminus f^{-1}(U)) > 0$ .
- (2)  $f$  is a  $\pi$ - $s$ -map, if  $f$  is both  $\pi$ -map and  $s$ -map.

*Remark 1.8.*

- (1) 1-sequence-covering map  $\implies$  sequence-covering map.
- (2) Sequence-covering map  $\implies$  quotient if  $Y$  is a sequential space.
- (3) Weak-open map  $\implies$  quotient map.

## 2. Main results

**Lemma 2.1.** Let  $f : X \rightarrow Y$  be a weak-open map. If  $X$  is first countable, then  $Y$  is  $gf$ -countable.

PROOF: Let  $f$  be a weak-open map, and  $X$  be first countable. Since  $f$  is a weak-open map, there exists a weak base  $\mathcal{B} = \bigcup\{\mathcal{B}_y : y \in Y\}$  for  $Y$  such that for each  $y \in Y$  there exists  $x_y \in f^{-1}(y)$  with the property that for every open neighborhood  $U$  of  $x_y$ , there exists  $B_y \in \mathcal{B}_y$  such that  $B_y \subset f(U)$ . Because  $X$  is first countable, there is a countable neighborhood base  $\mathcal{P}_x$  for every  $x \in X$ . Now, for each  $y \in Y$ , we put

$$\mathcal{F}_y = \{f(P) : P \in \mathcal{P}_{x_y}\}, \quad \mathcal{F} = \bigcup\{\mathcal{F}_y : y \in Y\}.$$

Since for each  $y \in Y$ ,  $x_y$  belongs to only countably many members of  $\mathcal{P}_{x_y}$ . It implies that  $\mathcal{F}_y$  is countable for every  $y \in Y$ . So, we only need to prove that  $\mathcal{F}$  is a weak base  $Y$ . Indeed,

(1) Note that  $f$  is a continuous map, and  $\mathcal{P}_{x_y}$  is a neighborhood base at  $x_y$ , it follows that  $\mathcal{F}_y$  is a network at  $y$  in  $Y$ .

(2) For each  $y \in Y$  and  $U, V \in \mathcal{F}_y$ , there exist  $P_1, P_2 \in \mathcal{P}_{x_y}$  such that  $x_y \in P_1 \cap P_2$ , and  $f(P_1) = U$ ,  $f(P_2) = V$ . Since  $\mathcal{P}_{x_y}$  is a neighborhood base at  $x_y$ , there exists  $P \in \mathcal{P}_{x_y}$  such that  $x_y \in P \subset P_1 \cap P_2$ . This implies that  $f(P) \in \mathcal{F}_y$ , and  $f(P) \subset f(P_1 \cap P_2) \subset U \cap V$ .

(3) Suppose that  $G$  is open in  $Y$ . Then for each  $y \in G$ ,  $x_y \in f^{-1}(G)$ . Since  $\mathcal{P}_{x_y}$  is a neighborhood base at  $x_y$ , there exists  $P \in \mathcal{P}_{x_y}$  such that  $x_y \in P \subset f^{-1}(G)$ . Thus,  $f(P) \in \mathcal{F}_y$ , and  $f(P) \subset G$ .

Conversely, suppose that  $G \subset Y$  is such that for each  $y \in G$ , there exists  $F \in \mathcal{F}_y$  satisfying  $F \subset G$ . Then there exists  $P \in \mathcal{P}_{x_y}$  such that  $x_y \in P$ , and  $F = f(P)$ . Since  $P$  is an open neighborhood of  $x_y$ , and  $f$  is a weak-open map, there exists  $B_y \in \mathcal{B}_y$  such that  $B_y \subset f(P)$ . So, for each  $y \in G$ , there exists  $B_y \in \mathcal{B}_y$  such that  $B_y \subset G$ . Because  $\mathcal{B}$  is a weak base,  $G$  is an open subset of  $Y$ .

By (1), (2), and (3),  $\mathcal{F}$  is a weak base for  $Y$ . Therefore  $Y$  is  $gf$ -countable.  $\square$

**Proposition 2.2.** *Let  $f : X \rightarrow Y$  be a sequence-covering map, and  $Y$  be  $snf$ -countable. If (1) or (2) holds, then  $f$  is a 1-sequence-covering map.*

- (1)  $f$  is an  $s$ -map, and  $X$  has a point-countable base.
- (2)  $f$  is a Lindelöf map, and  $X$  is first countable.

PROOF: Let  $\mathcal{P} = \bigcup\{\mathcal{P}_y : y \in Y\}$  be an  $sn$ -network for  $Y$  such that  $\mathcal{P}_y$  is countable for every  $y \in Y$ . We can suppose that  $\mathcal{P}_y$  is closed under finite intersections for every  $y \in Y$ .

(1) Let  $f$  be an  $s$ -map, and  $\mathcal{B}$  a point-countable base of  $X$ .

Firstly, we prove that for each  $y \in Y$ , there exists a point  $x_y \in f^{-1}(y)$  such that whenever  $U$  is an open neighborhood of  $x_y$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(U)$ . Otherwise, there exists  $y \in Y$  so that for every  $x \in f^{-1}(y)$ , there is an open neighborhood  $U_x$  of  $x$  such that  $P \not\subset f(U_x)$  for every  $P \in \mathcal{P}_y$ . Since  $\mathcal{B}$  is a base of  $X$ , for each  $x$ , there exists  $B_x \in \mathcal{B}$  such that  $x \in B_x \subset U_x$ . This implies that for every  $x \in f^{-1}(y)$ ,  $P \not\subset f(B_x)$  whenever  $P \in \mathcal{P}_y$ . On the other hand, because  $\mathcal{B}$  is a point-countable base and  $f^{-1}(y)$  is a separable subset of  $X$ , it follows that  $\{B_x : x \in f^{-1}(y)\}$  is countable. Assume that  $\{B_x : x \in f^{-1}(y)\} = \{B_m : m \in \mathbb{N}\}$ , and  $\mathcal{P}_y = \{F_n : n \in \mathbb{N}\}$ . Put  $\mathcal{R}_y = \{P_n = \bigcap_{i=1}^n F_i : n \in \mathbb{N}\}$ . It is easy to see that  $\mathcal{R}_y \subset \mathcal{P}_y$ , and  $P_{n+1} \subset P_n$ , for every  $n \in \mathbb{N}$ . Hence, for each  $m, n \in \mathbb{N}$ , there exists  $x_{n,m} \in P_n \setminus f(B_m)$ . For  $n \geq m$ , we denote  $y_k = x_{n,m}$  with  $k = m + n(n-1)/2$ . Since  $\mathcal{P}_y$  is a network at  $y$  and  $P_{n+1} \subset P_n$  for each  $n \in \mathbb{N}$ ,  $\{y_k\}$  is a sequence converging to  $y$  in  $Y$ . Because  $f$  is a sequence-covering map,  $\{y_k\}$  is an image of some sequence  $\{x_n\}$  converging to  $x \in f^{-1}(y)$  in  $X$ . Since  $x \in f^{-1}(y) \subset \bigcup\{B_m : m \in \mathbb{N}\}$ , there exists  $m_0 \in \mathbb{N}$  such that  $x \in B_{m_0}$ . So  $\{x\} \cup \{x_k : k \geq k_0\} \subset B_{m_0}$  for some  $k_0 \in \mathbb{N}$ . Thus,  $\{y\} \cup \{y_k : k \geq k_0\} \subset f(B_{m_0})$ . But if we take  $k \geq k_0$ , then there exists  $n \geq m_0$  such that  $y_k = x_{n,m_0}$ , and it implies that  $x_{n,m_0} \in f(B_{m_0})$ . This contradicts to  $x_{n,m_0} \in P_n \setminus f(B_{m_0})$ .

We now prove that  $f$  is a 1-sequence-covering map. Suppose that  $y \in Y$ . By the above proof there is  $x_y \in f^{-1}(y)$  such that whenever  $U$  is an open neighborhood of  $x_y$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(U)$ . Denote  $\{B_n : n \in \mathbb{N}\}$  a countable neighborhood base at  $x_y$  such that  $B_{n+1} \subset B_n$  for every  $n \in \mathbb{N}$ . Let  $\{y_n\}$  be any sequence in  $Y$ , which converges to  $y$ . Now, we choose a sequence  $\{z_n\}$  in  $X$  as follows.

Since  $B_n$  is an open neighborhood of  $x_y$ , by the above argument, there exists  $P_{k_n} \in \mathcal{P}_y$  satisfying  $P_{k_n} \subset f(B_n)$  for every  $n \in \mathbb{N}$ , and by assumption every  $P \in \mathcal{P}_y$  is a sequential neighborhood. It follows that for each  $n \in \mathbb{N}$ ,  $f(B_n)$  is a sequential neighborhood of  $y$  in  $Y$ . Hence, for each  $n \in \mathbb{N}$ , there exists  $i_n \in \mathbb{N}$  such that  $y_i \in f(B_n)$  for every  $i \geq i_n$ . Assume that  $1 < i_n < i_{n+1}$  for each  $n \in \mathbb{N}$ . Then for each  $j \in \mathbb{N}$ , we take

$$z_j = \begin{cases} z_j \in f^{-1}(y_j) & \text{if } j < i_1, \\ z_{j,n} \in f^{-1}(y_j) \cap B_n & \text{if } i_n \leq j < i_{n+1}. \end{cases}$$

Denote  $S = \{z_j : j \geq 1\}$ . Then  $S$  converges to  $x_y$  in  $X$  and  $f(S) = \{y_n\}$ .

(2) Let  $f$  be a Lindelöf map, and  $X$  be first countable. Firstly, we prove that for each  $y \in Y$ , there exists  $x_y \in f^{-1}(y)$  such that for every neighborhood  $U$  of  $x_y$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(U)$ . Indeed, if not, there exists  $y \in Y$  such that for each  $x \in f^{-1}(y)$ , there exists an open neighborhood  $U_x$  satisfying  $P \not\subset f(U_x)$  for every  $P \in \mathcal{P}_y$ . Since  $f$  is a Lindelöf map, and  $\{U_x : x \in f^{-1}(y)\}$  is an open cover of  $f^{-1}(y)$ , there exists a countable family  $\{U_n : n \in \mathbb{N}\} \subset \{U_x : x \in f^{-1}(y)\}$  such that  $f^{-1}(y) \subset \bigcup\{U_n : n \in \mathbb{N}\}$ . Now, using the argument from the proof of (1), this leads to a contradiction. Then, using again the proof of (1), we obtain that  $f$  is a 1-sequence-covering map.  $\square$

**Corollary 2.3.** *Let  $f : X \rightarrow Y$  be a map. If one of the following conditions is satisfied, then  $f$  is a 1-sequence-covering map.*

- (1)  $f$  is a sequence-covering  $s$ -map,  $X$  has a point-countable base and  $Y$  is  $gf$ -countable.
- (2)  $f$  is a sequence-covering Lindelöf map,  $X$  is first countable and  $Y$  is  $gf$ -countable.
- (3)  $f$  is a weak-open map and  $X$  is first countable.

PROOF: It follows from Proposition 2.2 and Remark 1.5 that both (1) and (2) imply that  $f$  is 1-sequence-covering. Assuming (3), because  $f$  is a weak-open map and  $X$  is first countable, it follows from Lemma 2.1 that  $Y$  is  $gf$ -countable. Since  $f$  is a weak-open map, for each  $y \in Y$ , there exists  $x_y \in f^{-1}(y)$  such that for every neighborhood  $U$  of  $x_y$ , there exists  $P \in \mathcal{P}_y$  satisfying  $P \subset f(U)$ . Then using the proof of Proposition 2.2(1), we have  $f$  is 1-sequence-covering.  $\square$

**Corollary 2.4** ([14]). *Every open map of a first countable space is 1-sequence-covering.*

**Theorem 2.5.** *If  $f : X \rightarrow Y$  is a sequence-covering  $\pi$ - $s$ -map, then  $f$  is a 1-sequence-covering map.*

PROOF: Firstly, we prove that  $Y$  is  $snf$ -countable. Let  $f : (X, d) \rightarrow Y$  be a sequence-covering  $\pi$ - $s$ -map, and  $(X, d)$  be a metric space. For each  $n \in \mathbb{N}$ , denote

$$\mathcal{F}_n = \left\{ f \left[ B \left( z, \frac{1}{n} \right) \right] : z \in X \right\}, \quad \text{with } B \left( z, \frac{1}{n} \right) = \left\{ y \in X : d(z, y) < \frac{1}{n} \right\}.$$

It is clear that  $\mathcal{F}_{n+1}$  refines  $\mathcal{F}_n$  for each  $n \in \mathbb{N}$ . Now, for each  $x \in Y$ , we put  $\mathcal{P}_x = \{\text{st}(x, \mathcal{F}_n) : n \in \mathbb{N}\}$ ,  $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in Y\}$ . Then,

(i) It is obvious that  $\mathcal{P}_x$  is countable for every  $x \in Y$ .

(ii) Let  $U$  be an open neighborhood of  $x$ . Since  $f$  is a  $\pi$ -map, there exists  $n \in \mathbb{N}$  such that  $d(f^{-1}(x), X \setminus f^{-1}(U)) > \frac{1}{n}$ . Take  $m \in \mathbb{N}$  such that  $m \geq 2n$ . It is easy too see that if  $x \in f[B(z, \frac{1}{m})]$  for some  $z \in X$ , then  $B(z, \frac{1}{m}) \subset f^{-1}(U)$ , so  $f[B(z, \frac{1}{m})] \subset U$ . Since

$$\text{st}(x, \mathcal{F}_m) = \bigcup \left\{ f \left[ B \left( z, \frac{1}{m} \right) \right] : B \left( z, \frac{1}{m} \right) \cap f^{-1}(x) \neq \emptyset \right\},$$

this implies that  $\text{st}(x, \mathcal{F}_m) \subset U$ . Therefore  $\mathcal{P}_x$  is a network at  $x$ .

(iii) Let  $P_1, P_2 \in \mathcal{P}_x$ . Then there exist  $m, n \in \mathbb{N}$  such that  $P_1 = \text{st}(x, \mathcal{F}_m)$ , and  $P_2 = \text{st}(x, \mathcal{F}_n)$ . Pick  $k \in \mathbb{N}$  such that  $k > \max\{m, n\}$  and put  $P = \text{st}(x, \mathcal{F}_k)$ . It is obvious that  $P \in \mathcal{P}_x$ . Suppose  $y \in P = \text{st}(x, \mathcal{F}_k)$ ; then there exists  $z_1$  such that  $y \in f[B(z_1, \frac{1}{k})] \in \mathcal{F}_k$ . Since  $\mathcal{F}_k$  refines  $\mathcal{F}_m$  and  $\mathcal{F}_n$ , there exist  $f[B(z_2, \frac{1}{m})] \in \mathcal{F}_m$  and  $f[B(z_3, \frac{1}{n})] \in \mathcal{F}_n$  such that  $f[B(z_1, \frac{1}{k})] \subset f[B(z_2, \frac{1}{m})]$ , and  $f[B(z_1, \frac{1}{k})] \subset f[B(z_3, \frac{1}{n})]$ . Thus  $y \in f[B(z_2, \frac{1}{m})] \cap f[B(z_3, \frac{1}{n})]$ , so  $y \in \text{st}(x, \mathcal{F}_m) \cap \text{st}(x, \mathcal{F}_n)$ . Therefore  $P \subset P_1 \cap P_2$ .

(iv) Suppose  $P \in \mathcal{P}_x$  and let  $\{x_n\}$  be any sequence in  $Y$  which converges to  $x$  in  $Y$ . Because  $f$  is a sequence-covering map,  $\{x_n\}$  is the image of some sequence  $\{z_n\}$  converging to  $z \in f^{-1}(x)$  in  $X$ . Since  $P \in \mathcal{P}_x$ , there exists  $m \in \mathbb{N}$  such that

$$P = \text{st}(x, \mathcal{F}_m) = \bigcup \left\{ f \left[ B \left( z, \frac{1}{m} \right) \right] : f^{-1}(x) \cap B \left( z, \frac{1}{m} \right) \neq \emptyset \right\}.$$

Since  $\{z_n\}$  converges to  $z \in f^{-1}(x)$  and  $\bigcup\{B(z, \frac{1}{m}) : f^{-1}(x) \cap B(z, \frac{1}{m}) \neq \emptyset\}$  is a neighborhood of  $f^{-1}(x)$ ,  $\{z_n\}$  is eventually in  $\bigcup\{B(z, \frac{1}{m}) : f^{-1}(x) \cap B(z, \frac{1}{m}) \neq \emptyset\}$ . This implies that  $\{x_n\}$  is eventually in  $P$ . Hence,  $P$  is a sequential neighborhood of  $x$  in  $Y$ . Therefore,  $Y$  is  $snf$ -countable.

Then, it follows from Proposition 2.2(1) that  $f$  is a 1-sequence-covering map. □

*Remark 2.6.* Since every compact map of a metric space is a  $\pi$ -s-map, Theorem 2.5 is a generalization of Theorem 1.1. Furthermore, this generalization is proper.

**Example 2.7.** There is a sequence-covering  $\pi$ -s-map  $f : X \rightarrow Y$  of a metric space  $X$  which is not a sequence-covering compact map.

PROOF: Indeed, let  $\mathbb{R}$  be the real line with usual Euclidean topology. Denote  $X = \mathbb{R}$  and  $Y = (-\infty; 0]$ . Now, we define  $f : X \rightarrow Y$  by

$$f(t) = \begin{cases} t & \text{if } t \in (-\infty; 0] \\ 0 & \text{if } t \in [0; +\infty). \end{cases}$$

Then, we have

(1)  $f$  is a surjective, continuous  $s$ -map. It is obvious.

(2)  $f$  is a  $\pi$ -map. Indeed, let  $y \in (-\infty; 0]$  and  $U$  be an open neighborhood of  $y$ .

If  $y = 0$ , then  $f^{-1}(0) = [0, +\infty)$  and there is  $\varepsilon > 0$  such that  $0 \in (-\varepsilon; 0] \subset U$ . Hence, we have

$$\begin{aligned} d\left(f^{-1}(y), \mathbb{R} \setminus f^{-1}(U)\right) &\geq d\left([0, +\infty), \mathbb{R} \setminus f^{-1}[(-\varepsilon; 0]]\right) \\ &= d\left([0, +\infty), (-\infty; -\varepsilon]\right) \geq \varepsilon > 0. \end{aligned}$$

If  $y \neq 0$ , then  $f^{-1}(y) = \{y\}$  and there is  $\varepsilon > 0$  such that  $(y - \varepsilon, y + \varepsilon) \subset U$ . So, we have

$$\begin{aligned} d\left(f^{-1}(y), \mathbb{R} \setminus f^{-1}(U)\right) &\geq d\left(\{y\}, \mathbb{R} \setminus f^{-1}[(y - \varepsilon; y + \varepsilon)]\right) \\ &= d\left(y, (-\infty; y - \varepsilon] \cup [y + \varepsilon; 0]\right) \geq \varepsilon > 0. \end{aligned}$$

Therefore,  $f$  is a  $\pi$ -map.

(3)  $f$  is a sequence-covering map. In fact, suppose  $S = \{y_n : n \in \omega\}$  is any sequence converging to  $y_0$  in  $Y$ . Since  $Y \subset X$ ,  $S = \{y_n : n \in \omega\} \subset X$ . By definition of  $f$ , we have  $f(y_n) = y_n$  for every  $n \in \omega$ . Therefore  $f$  is a 1-sequence-covering map.

(4)  $f$  is not a compact map. Since  $f^{-1}(0) = [0, +\infty)$ , and  $[0, +\infty)$  is not compact in  $\mathbb{R}$ , it follows that  $f$  is not compact.

By Corollary 3.6 in [25], Remark 1.8 and Theorem 2.5, we get □

**Corollary 2.8.** *Let  $f : M \rightarrow X$  be a map. If  $M$  is a metric space, then the following are equivalent:*

- (1)  $f$  is a weak-open  $\pi$ - $s$ -map;
- (2)  $f$  is a 1-sequence-covering, quotient  $\pi$ - $s$ -map;
- (3)  $f$  is a sequence-covering, quotient  $\pi$ - $s$ -map.

**Corollary 2.9.** *Let  $f : M \rightarrow X$  be a map. If  $M$  is a metric space, then the following are equivalent:*

- (1)  $f$  is a weak-open compact map;
- (2)  $f$  is a 1-sequence-covering, quotient compact map;
- (3)  $f$  is a sequence-covering, quotient compact map.

PROOF: It follows immediately by Corollary 2.8. □

**Acknowledgment.** The authors would like to express their thanks to referee for his/her helpful comments and valuable suggestions.



## REFERENCES

- [1] Arhangel'skii A.V., *Mappings and spaces*, Russian Math. Surveys **21** (1966), no. 4, 115–162.
- [2] Davis S.W., *More on Cauchy conditions*, Topology Proc. **9** (1984), 31–36.
- [3] Engelking R., *General Topology*, PWN–Polish Scientific Publishers, Warszawa, 1977.
- [4] Franklin S.P., *Spaces in which sequences suffice*, Fund. Math. **57** (1965), 107–115.
- [5] Ge X., *Spaces with a locally countable sn-network*, Lobachevskii J. Math. **26** (2007), 33–49.
- [6] Ge Y., *Characterizations of sn-metrizable spaces*, Publ. Inst. Math. (Beograd) (N.S.) **74** (88) (2003), 121–128.
- [7] Ge Y., *Spaces with countable sn-networks*, Comment. Math. Univ. Carolin. **45** (2004), no. 1, 169–176.
- [8] Gruenhagen G., Michael E., Tanaka Y., *Spaces determined by point-countable covers*, Pacific J. Math. **113** (1984), no. 2, 303–332.
- [9] Ikeda Y., Tanaka Y., *Spaces having star-countable k-networks*, Topology Proc. **18** (1993), 107–132.
- [10] Ikeda Y., Liu C., Tanaka Y., *Quotient compact images of metric spaces, and related matters*, Topology Appl. **122** (2002), no. 1–2, 237–252.
- [11] Li Z., *On  $\pi$ -s-images of metric spaces*, Int. J. Math. Math. Sci. **7** (2005), 1101–1107.
- [12] Li Z., Lin S., *On the weak-open images of metric spaces*, Czechoslovak Math. J. **54** (2004), 939–400.
- [13] Lin S., *Generalized Metric Spaces and Mappings*, Chinese Science Press, Beijing, 1995 (Chinese).
- [14] Lin S., *On sequence-covering s-mappings*, Adv. Math. (China) **25** (1996), no. 6, 548–551.
- [15] Lin S., *Point-Countable Covers and Sequence-Covering Mappings*, Chinese Science Press, Beijing, 2002.
- [16] Lin S., Yan P., *Sequence-covering maps of metric spaces*, Topology Appl. **109** (2001), 301–314.
- [17] Liu C., *On weak bases*, Topology Appl. **150** (2005), 91–99.
- [18] O'Meara P., *On paracompactness in function spaces with the compact-open topology*, Proc. Amer. Math. Soc. **29** (1971), 183–189.
- [19] Michael E.,  $\aleph_0$ -spaces, J. Math. Mech. **15** (1966), 983–1002.
- [20] Siwiec F., *On defining a space by a weak base*, Pacific J. Math. **52** (1974), 233–245.
- [21] Tanaka Y., *Point-countable covers and k-networks*, Topology Proc. **12** (1987), 327–349.
- [22] Tanaka Y., *Theory of k-networks II*, Questions Answers Gen. Topology **19** (2001), 27–46.
- [23] Tanaka Y., Ge Y., *Around quotient compact images of metric spaces, and symmetric spaces*, Houston J. Math. **32** (2006), no. 1, 99–117.
- [24] Yan P., *On strong sequence-covering compact mappings*, Northeastern Math. J. **14** (1998), 341–344.
- [25] Xia S., *Characterizations of certain g-first countable spaces*, Adv. Math. (China) **29** (2000), 61–64.

DEPARTMENT OF MATHEMATICS, VINH UNIVERSITY, VIETNAM

E-mail: andhv@yahoo.com

luongtuyench12@yahoo.com

(Received December 2, 2007, revised March 13, 2008)