

Harry Levy

Conformal invariants in two dimensions. [I.]

Časopis pro pěstování matematiky a fysiky, Vol. 69 (1940), No. 2, 50--56

Persistent URL: <http://dml.cz/dmlcz/121984>

## Terms of use:

© Union of Czech Mathematicians and Physicists, 1940

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Conformal Invariants in Two Dimensions I.

Harry Levy, Urbana, Illinois.

(Received February 4, 1938.)

The conformal geometry of Riemannian spaces has been studied by Hlavatý, Cartan, T. Y. Thomas, Schouten, and others who have developed various algorithms for  $n > 2$ . In particular, Hlavatý<sup>1</sup>) has found the generalized conformal curvatures of a curve. But for  $n = 2$  a single curve has no conformal invariants since any two surfaces are conformally equivalent. Here we shall obtain some invariants of two or more curves in the two dimensional case.

1. Let  $V_n$  and  $V'_n$  be two Riemannian spaces in conformal correspondence so that the coefficients of their linear elements

$$ds^2 = g_{\lambda\mu} dx^\lambda dx^\mu \quad ds'^2 = g'_{\lambda\mu} dx^\lambda dx^\mu \quad (1)$$

can be taken in the relation

$$g'_{\lambda\mu} = \sigma g_{\lambda\mu} \quad (\lambda, \mu = 1, 2, \dots, n). \quad (2)$$

The reciprocal elements  $g^{\lambda\mu}$  and the determinant  $g = |g_{\lambda\mu}|$  (which we assume to be different from zero) then satisfy

$$g'^{\lambda\mu} = \sigma^{-1} g^{\lambda\mu} \quad g' = \sigma^n g. \quad (3)$$

The Christoffel symbols of the second kind, which we designate by  $\left\{ \begin{smallmatrix} \lambda \\ \mu\nu \end{smallmatrix} \right\}$  will satisfy the relation

$$\left\{ \begin{smallmatrix} \lambda \\ \mu\nu \end{smallmatrix} \right\}' = \left\{ \begin{smallmatrix} \lambda \\ \mu\nu \end{smallmatrix} \right\} + \sigma_\mu \delta_\nu^\lambda + \sigma_\nu \delta_\mu^\lambda - g_{\mu\nu} \sigma^\lambda, \quad (4)$$

where  $\sigma^\lambda = g^{\lambda\mu} \sigma_\mu$  and  $\sigma_\mu = \frac{1}{2} \frac{\partial}{\partial x^\mu} \log \sigma^2$ .

We assume that  $V_n$  and  $V'_n$  are analytic so that  $g_{\lambda\mu}$  and  $g'_{\lambda\mu}$

<sup>1</sup>) Zur Conformgeometrie, Akad. Wetensch. Amsterdam, Proc. 88 (1935), pp. 281, 738, 1006.

<sup>2</sup>) Cf. Eisenhart, Riemannian Geometry, Princeton University Press (1926), p. 89.

(and therefore  $\sigma$  too) are continuous and have continuous derivatives of all orders.

Let  $C$  be an analytic curve and  $C'$  its transform. The unit components of the tangent vectors,  $\lambda_1^\mu$  and  $\lambda'_1{}^\mu$  respectively, satisfy

$$\lambda'_1{}^\mu = \sigma^{-\frac{1}{2}} \lambda_1^\mu \quad (5)$$

and we obtain by a direction computation that the components of the principle normal (if defined) satisfy

$$k'_1 \lambda'_2{}^\mu = \sigma^{-1} (k_1 \lambda_2{}^\mu + \sigma_r \lambda_1{}^r \lambda_1{}^\mu - \sigma^\mu),^3 \quad (6)$$

where  $k_1$  and  $k'_1$  are the first curvatures of  $C$  and  $C'$  respectively.

From equations (3) we have that  $\sigma = (g'/g)^{1/n}$  and therefore that

$$\sigma_\mu = \frac{1}{2n} \frac{\partial}{\partial x^\mu} \log (g'/g). \quad (7)$$

If we eliminate  $\sigma$  from (6) by means of (7) we obtain that  $C^\mu$  defined by

$$C^\mu = g^{1/n} \left\{ k_1 \lambda_2{}^\mu + \frac{1}{2n} (g^{\mu r} - \lambda_1{}^\mu \lambda_1{}^r) \frac{\partial}{\partial x^r} \log g \right\} \quad (8)$$

is invariant under conformal transformations.

2. Henceforth we assume  $n = 2$ . In this case there is a unique direction normal to a given direction, and consequently we must have

$$\lambda'_2{}^r = \sigma^{-\frac{1}{2}} \lambda_2{}^r \quad (r, s = 1, 2), \quad (9)$$

where the ambiguity in sign can be regarded as incorporated in  $\sigma^{-\frac{1}{2}}$  itself. From (2) and (3) it then follows that

$$g'^{-\frac{1}{2}} \lambda'_2{}^r g'_{rs} = g^{-\frac{1}{2}} \lambda_2{}^r g_{rs} \quad (10)$$

and multiplying the right hand sides of (8) and (10) and contracting we obtain that the expression  $I_1$ , defined by

$$I_1 = g^{\frac{1}{2}} \left( k + \frac{1}{4} \lambda_2{}^r \frac{\partial}{\partial x^r} \log g \right) \quad (11)$$

is invariant under conformal transformations.

It will be convenient to write  $\mu^r$  for  $\lambda_2{}^r$  and simply  $\lambda^r$  for  $\lambda_1{}^r$  and

$$\varphi = g^{\frac{1}{2}}, \quad \varphi_r = \frac{\partial}{\partial x^r} \varphi$$

so that (11) may be written

$$I_1 = \varphi k + \varphi_r \mu^r. \quad (12)$$

<sup>3)</sup> Cf. Modesitt, Some Singular Properties of Conformal Transformations between Riemannian Spaces, Am. Journ. of Math., 1938.

We obtain a sequence of functions  $I_\alpha$  ( $\alpha = 1, 2, 3, \dots$ ) from (12) by differentiation with respect to the arc  $s$  of the given curve and multiplication by  $\varphi$ ,

$$I_\alpha = \varphi \frac{d}{ds} I_{\alpha-1} \quad (\alpha = 2, 3, 4, \dots) \quad (13)$$

and clearly  $I_\alpha$  is (for all values of  $\alpha$ ) invariant under conformal transformations. We can readily obtain by means of the Frenet equations that the values of  $I_\alpha$  for  $\alpha = 2, 3$  are given by

$$I_2 = \varphi^2 \frac{d}{ds} k + \varphi \varphi_{rs} \mu^r \lambda^s, \quad (14)$$

$$I_3 = \varphi^3 \frac{d^2}{ds^2} k + 2\varphi^2 \varphi_r \lambda^r \frac{d}{ds} k + \varphi^2 \varphi_{rs} (\mu^r \mu^s - \lambda^r \lambda^s) k + \\ + \varphi^2 (\varphi_{rst} + \varphi \varphi_t \varphi_{rs}) \lambda^r \mu^s \lambda^t, \quad (15)$$

where  $\varphi_{r\dots st}$  is obtained from  $\varphi_{r\dots s}$  by the formal process of covariant differentiation with respect to the  $g's$ ,

$$\varphi_{rs} = \frac{\partial}{\partial x^s} \varphi_r - \varphi_u \left\{ \begin{matrix} u \\ rs \end{matrix} \right\}$$

etc.

We can prove by induction that  $I_\alpha$ , for an arbitrary value of  $\alpha$ , has an expansion of the following form

$$I_\alpha = \varphi^\alpha k^{(\alpha-1)} + a_\alpha \varphi^{\alpha-1} \varphi_r \lambda^r k^{(\alpha-2)} \\ + \{b_\alpha \varphi^{\alpha-2} (\varphi_r \lambda^r)^2 + (c_\alpha + 1) \varphi^{\alpha-1} \varphi_{rs} \lambda^r \lambda^s + \varphi^{\alpha-1} \varphi_{rs} \mu^r \mu^s\} \\ + d_\alpha \varphi^{\alpha-1} \varphi_r \mu^r k\} k^{\alpha-1} + *$$

where  $a_\alpha, b_\alpha, c_\alpha, d_\alpha$  are the constants given by

$$a_\alpha = \frac{1}{2} (\alpha - 2) (\alpha + 1), \quad b_\alpha = \frac{1}{4!} (\alpha - 1) (\alpha - 2) (\alpha - 3) (3\alpha + 4), \\ c_\alpha = \frac{1}{3!} \alpha (\alpha + 1) (\alpha - 4), \quad d_\alpha = \frac{1}{3!} (\alpha - 3) (\alpha^2 - 4), \quad (17)$$

and the \* represents a polynomial in  $\lambda^r, \mu^r, \varphi, \varphi_r, \dots, \varphi_{r_1 \dots r_\alpha}, k, k', k'', \dots, k^{(\alpha-4)}$ .

We can prove further by induction by means of (13) that  $I_\alpha$  regarded as a function of  $\mu^r, k, k', \dots, k^{(\alpha-1)}$  is an odd function of these variables taken together, that is

$$I_\alpha (-\mu^r, -k, -k', \dots, -k^{(\alpha-1)}) = -I_\alpha (\mu^r, k, k', \dots, k^{(\alpha-1)}).$$

3. Let  $C_i$  ( $i = 1, 2, 3$ ) be three curves concurrent in a point  $P_0$  of the surface  $V_3$ , and let  $\lambda_i^a$  be the contravariant components of their unit tangent vectors and  $\mu_i^r$  of their unit normal vectors.

Choose a positive direction of rotation about  $P_0$  and let  $\Theta_i$  be the directed angle from  $\mu_{i+1}$  to  $\mu_{i+2}$ .<sup>4)</sup>

Three directions at a point in two dimensions are necessarily linearly dependent, and it follows directly that the coefficients of dependence for the normal directions are the sines of their angles, so that

$$\mu_1^r \sin \Theta_1 + \mu_2^r \sin \Theta_2 + \mu_3 \sin \Theta_3 = 0. \quad (18)$$

If we write (12) for each of the three curves and evaluate at their common point  $P$ , multiply each of these three equations by the corresponding  $\sin \Theta_i$  and add, the terms in  $\mu$  drop out by virtue of (18) and we obtain

$$\Sigma I_1 \sin \Theta = \varphi \Sigma k \sin \Theta. \quad (19)$$

The left hand side is invariant under conformal transformations and the right under coordinate transformations. We have consequently the following theorem:

If three curves  $C_i$  on a surface are concurrent and if  $K_i$  are the values, at their common point, of their (geodesic) curvatures and  $\Theta_1, \Theta_2, \Theta_3$  the directed angles from the normals to  $C_2, C_3, C_1$  to the normals to  $C_3, C_1, C_2$  respectively, then  $\sqrt{g} (k_1 \sin \Theta_1 + k_2 \sin \Theta_2 + k_3 \sin \Theta_3)$  is an absolute conformal invariant.

This theorem has a particularly simple geometric interpretation in the Euclidean plane; if we assume that  $k_1 k_2 k_3 \neq 0$ , the sum with which we are concerned differs by a factor from

$$R_2 R_3 \sin \Theta_1 + R_1 R_3 \sin \Theta_2 + R_1 R_2 \sin \Theta_3,$$

where the  $R$ 's are the radii of curvature, reciprocals of the  $K$ 's and we observe that this expression represents the area of the triangle formed by the three centers of curvature. In particular, it then follows that if the centers of curvature for a common point of three curves in the Euclidean plane are collinear, they remain collinear under conformal transformations of the plane into itself.

The above results can be extended by writing in place of (18) the linear equations satisfied by the  $\lambda_S$  and by the products  $\lambda_\mu$ . Let us assume for the sake of definiteness that the positive direction

of rotation has been chosen so that the angle from  $\lambda_1$  to  $\mu_1$  is  $+\frac{\pi}{2}$ .

Then  $\Theta_3$  and  $-\Theta_3$  are the directed angles from  $\mu_1$  to  $\mu_2$  and  $\mu_3$  respectively and we have that

<sup>4)</sup> It is of course understood that by  $\mu_k$  we mean  $\mu_{k-3}$  if  $k > 3$ .

$$\begin{aligned}
\lambda_2^r &= e (\cos \Theta_3 \lambda_1^r + \sin \Theta_3 \mu_1^r), \\
\lambda_3^r &= \bar{e} (\cos \Theta_2 \lambda_1^r - \sin \Theta_2 \mu_1^r), \\
\mu_2^r &= -\sin \Theta_3 \lambda_1^r + \cos \Theta_3 \mu_1^r, \\
\mu_3^r &= \sin \Theta_2 \lambda_1^r + \cos \Theta_2 \mu_1^r,
\end{aligned} \tag{20}$$

where  $e(\bar{e})$  is  $\pm 1$  according as  $(\lambda_2, \mu_2)$  ( $\lambda_3, \mu_3$ ) and  $\lambda_1, \mu_1$  have the same or opposite orientation.

It follows by direct computation, by means of (20), that

$$\sin \Theta_1 \lambda_1^r + e \sin \Theta_2 \lambda_2^r + \bar{e} \sin \Theta_3 \lambda_3^r = 0 \tag{21}$$

and, denoting by  $S$  the symmetric part, that is

$$S \{ \sin 2\Theta_1 \lambda_1^r \mu_1^r + e \sin 2\Theta_2 \lambda_2^r \mu_2^r + \bar{e} \sin 2\Theta_3 \lambda_3^r \mu_3^r \} = 0. \tag{22}$$

We can now write (14) for each of the three curves, multiply each  $I_2$  by the appropriate  $\pm \sin 2\Theta$  and add the three products. The terms in  $\lambda\mu$  drop out by virtue of (22) leaving

$$\begin{aligned}
I_2^{(1)} \sin 2\Theta_1 + e I_2^{(2)} \sin 2\Theta_2 + \bar{e} I_2^{(3)} \sin 2\Theta_3 &= \\
&= \varphi^2 (k'_1 \sin 2\Theta_1 + e K'_2 \sin 2\Theta_2 + \bar{e} K'_3 \sin 2\Theta_3).
\end{aligned}$$

Hence we have the theorem:

If three curves  $C_i$  on a surface have a common point and if  $k'_i$  is the value there of the derivative of the (geodesic) curvature of  $C_i$  with respect to its arc, if  $\Theta_i$  is the directed angle from the normal to  $C_{i+1}$ <sup>4)</sup> to the normal to  $C_{i+2}$  and  $e$  and  $\bar{e}$  are  $\pm 1$  (as above defined), then  $\sqrt{g} (k'_1 \sin 2\Theta_1 + e k'_2 \sin 2\Theta_2 + \bar{e} k'_3 \sin 2\Theta_3)$  is invariant under conformal maps of the surface.<sup>5)</sup>

4. In the determination of the curvature of a curve an ambiguity may arise in the choice of sign. Let us assume that we take the (geodesic) curvature as always non-negative and that thereby a positive normal is determined in accordance with the Frenet equations.<sup>6)</sup> If we introduce normal coordinates at  $P_0$  and find the expansion of the coordinates of the points of a curve in a neighborhood of  $P_0$  we see that the given curve and its positive principal normal lie on the same side of the geodesic tangent at  $P_0$  to the curve. Hence if two curves are tangent and lie on the same side of their common tangent geodesic they have the same principal normal whereas if they lie on opposite sides of their common tangent geodesic their normals are directed oppositely.

<sup>5)</sup> If a conformal transformation of  $V_2$  into itself interchanges  $C_1$  and  $C_2$  and leaves  $C_3$  invariant this and the preceding theorem reduce to special theorems obtained by Kasner in the plane. Cf. *Geometry of Conformal Symmetry*, *Annals of Math.*, 2nd series, vol. 38 (1937) pp. 876—877.

<sup>6)</sup> See, for example, Eisenhart, l. c., p. 106.

Suppose  $C_2$  and  $C_3$  are tangent to each other but not to  $C_1$  and that their tangent vectors have the same direction. Then  $e$  and  $\bar{e}$  of equations (20) will have the same or opposite signs according as  $C_2$  and  $C_3$  lie on the same or opposite sides of their common tangent geodesic. But in either case  $\sin \theta_1 = 0$  and  $\sin 2\theta_2 = -\sin 2\theta_3$  while  $\sin \theta_2 = \mp \sin \theta_3$ . The preceding theorem reduces to the following special case:

$\sqrt{g}(k_1 \mp k_2)$  and  $\sqrt{g}(k'_1 \mp k'_2)$  are absolute conformal invariants of two tangent curves where the  $\mp$  is to be taken according as the curves lie on the same or opposite sides of their common tangent geodesic.

The derivation of the result that  $\sqrt{g}(k_1 \mp k_2)$  and  $\sqrt{g}(k'_1 \mp k'_2)$  are absolute conformal invariants was communicated to the writer by Professor Hlavaty who obtained it by methods only different from the above. Others have obtained somewhat similar results.<sup>7)</sup>

If two curves are tangent and have contact of order  $h > 1$  either  $k_1 = k_2, k'_1 = k'_2, \dots, k_1^{(h-2)} = k_2^{(h-2)}$  and they lie on the same side of their common geodesic tangent so that the positive direction of their normals coincide, or  $k_1 = k_2 = k'_1 = \dots = k_2^{(h-2)} = 0$  in which case the positive directions of their normals need not coincide. It follows from (16) and the remarks in the paragraph following (17) that the functions  $I_\alpha$  for  $\alpha = 1, 2, \dots, h-1$  formed for one curve are equal, except possibly for sign, to the corresponding functions for the other curve and that the functions  $I_h$  differ only in the first term so that

$$I_h^{(1)} \mp I_h^{(2)} = \varphi^h (k_1^{(h-1)} \mp k_2^{(h-1)}),$$

where the  $\mp$  sign is to be taken according as the normals coincide or are oppositely directed. Hence we have the result:

If two curves have contact of order  $h, \sqrt{g^h}(k_1^{(h-1)} \mp k_2^{(h-1)})$  is an absolute conformal invariant.

Equations (14) yield one final result. If  $C$  and  $C'$  are two curves intersecting orthogonally, the pairs of directions  $\lambda_1, \mu_1$

<sup>7)</sup> Cf. Kasner, *Conformal Geometry*, Proceedings, Fifth Int. Cong. 2 (1912) p. 81; Ostrowski, *Berührungsmaße, nullwinklige Kreisbogendreiecke und die Modulfigur*, Jahresb. Deut. Math.-Verein. 44 (1934) p. 56; and Kasner and Comenetz, *Conformal Geometry of Horn Angles*, Proceedings, Nat. Ac. Sciences (Washington) 22 (1936) p. 303. These writers restrict themselves to conformal transformations of the Euclidean plane into itself and naturally find only the relative invariants  $k_1 - k_2$  and  $k'_1 - k'_2$ . Comenetz, *Conformal Geometry on a Surface*, Bull. Am. Math. Soc. 42 (1936) p. 806 extends the results of Kasner and Comenetz to surfaces and notes that  $(k'_1 - k'_2)/(k_1 - k_2)^2$  is an absolute invariant.

for  $C$  and  $\lambda_2, \mu_2$  for  $C'$  differ at most in orientation. If they have the same orientation

$$\lambda_1 = \pm \mu_2, \mu_1 = \mp \lambda_2$$

whereas the signs are reversed if they have opposite orientation. Writing equations (14) for each of the two curves and adding in the first case, subtracting in the second gives the following theorem:

If two curves intersect orthogonally,  $\sqrt{g}(k'_1 \pm k'_2)$  is an absolute conformal invariant.<sup>8)</sup>

*Institute for Advanced Study, Princeton, N. J., and University of Illinois, Urbana, Illinois.*

\*

### Konformní invarianty ve dvou dimensích I.

(Obsah předešlého článku.)

Autor nalézá algoritmus, kterým lze sestrojiti konformní invarianty křivek na ploše rekurentními formullemi. To mu dovoluje, mezi jiným, studovati konformní invarianty dvou křivek na ploše, jež v daném bodě mají styk libovolného řádu.

\*

### Konforminvarianten in zwei Dimensionen I.

(Auszug aus dem vorstehenden Artikel.)

Der Verfasser findet einen Algorithmus, mit welchem es gelingt, die Konforminvarianten der Kurven auf einer Fläche durch rekurrente Formeln anzugeben. Dies ermöglicht ihm, die Konforminvarianten von zwei Kurven zu studieren, wenn sich diese Kurven in einem Punkte in beliebiger Ordnung berühren.

---

<sup>8)</sup> Kasner, *Annals* l. c. p. 879 shows that in the plane the sign of  $k'_1 + k'_2$  is a conformal invariant of a general right angle.