

Bohdan Zelinka

Point-set domatic numbers of graphs

Mathematica Bohemica, Vol. 124 (1999), No. 1, 77–82

Persistent URL: <http://dml.cz/dmlcz/125976>

Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

POINT-SET DOMATIC NUMBERS OF GRAPHS

BOHDAN ZELINKA, Liberec

(Received September 10, 1997)

Abstract. A subset D of the vertex set $V(G)$ of a graph G is called point-set dominating, if for each subset $S \subseteq V(G) - D$ there exists a vertex $v \in D$ such that the subgraph of G induced by $S \cup \{v\}$ is connected. The maximum number of classes of a partition of $V(G)$, all of whose classes are point-set dominating sets, is the point-set domatic number $d_p(G)$ of G . Its basic properties are studied in the paper.

Keywords: dominating set, point-set dominating set, point-set domatic number, bipartite graph

MSC 1991: 05C35

The point-set domatic number of a graph is a variant of the domatic number $d(G)$ of a graph, which was introduced by E. J. Cockayne and S. T. Hedetniemi [1], and of the point-set domination number $\gamma_p(G)$, which was introduced by E. Sampathkumar and L. Pushpa Latha in [3] and [4]. We will describe its basic properties. All graphs considered are finite undirected graphs without loops and multiple edges.

A subset D of the vertex set $V(G)$ of a graph G is called dominating, if for each vertex $x \in V(G) - D$ there exists a vertex $y \in D$ adjacent to x . It is called point-set dominating (or shortly *ps*-dominating), if for each subset $S \subseteq V(G) - D$ there exists a vertex $v \in D$ such that the set $S \cup \{v\}$ induces a connected subgraph of G . A partition of $V(G)$ is called domatic (or point-set domatic), if all of its classes are dominating (or *ps*-dominating, respectively) sets in G . The maximum number of classes of a domatic (or point-set domatic) partition of $V(G)$ is called the domatic (or point-set domatic, respectively) number of G . The domatic number of G is denoted by $d(G)$, the point-set domatic number of G is denoted by $d_p(G)$. Instead of “point-set domatic” we will say shortly “*ps*-domatic”.

For every graph G there exists at least one *ps*-domatic partition of $V(G)$, namely $\{V(G)\}$. Therefore $d_p(G)$ is well-defined for every graph G .

Evidently each ps -dominating set in G is a dominating set in G and thus we have a proposition.

Proposition 1. For every graph G the inequality

$$d_p(G) \leq d(G)$$

holds.

Each vertex of a complete graph K_n forms a one-element ps -dominating set and therefore the following proposition holds.

Proposition 2. For every complete graph K_n its ps -domatic number satisfies

$$d_p(K_n) = n.$$

A similar assertion holds for a complete bipartite graph $K_{m,n}$.

Proposition 3. Let $K_{m,n}$ be a complete bipartite graph with $2 \leq m \leq n$. Then

$$d_p(K_{m,n}) = m.$$

Proof. Let U, V be the bipartition classes of $K_{m,n}$. Let $u \in U, v \in V$ and consider the set $D = \{u, v\}$. Let $S \subseteq V(K_{m,n}) - D$. If $S \subseteq U$, then $S \cup \{v\}$ induces a subgraph which is a star and thus it is connected. If $S \subseteq V$, then so is $S \cup \{u\}$. Suppose that $S \cap U \neq \emptyset, S \cap V \neq \emptyset$. The set S itself induces a connected subgraph, namely a complete bipartite graph. The vertex u is adjacent to a vertex of $S \cap V$ and thus also $S \cup \{u\}$ induces a connected subgraph; the set $D = \{u, v\}$ is ps -dominating. If $U = \{u_1, \dots, u_m\}, V = \{v_1, \dots, v_n\}$, we take $D_i = \{u_i, v_i\}$ for $i = 1, \dots, m-1$ and $D_m = \{u_m, v_m, \dots, v_n\}$. Then $\{D_1, \dots, D_m\}$ is a ps -domatic partition of $K_{m,n}$ and $d_p(K_{m,n}) \geq m$. On the other hand, $d_p(K_{m,n}) \leq d(K_{m,n}) = m$ and thus $d_p(K_{m,n}) = m$. \square

Proposition 4. Let n be an even integer, let G be obtained from the complete graph K_n by deleting edges of a linear factor. Then

$$d_p(G) = n/2.$$

Proof. Evidently each pair of non-adjacent vertices in G is ps -dominating and there exists a partition of $V(G)$ into $n/2$ such sets. On the other hand, no one-vertex ps -dominating set exists. This implies the assertion. \square

Now we will prove some theorems. By $d_G(x, y)$ we denote the distance between vertices x, y in a graph G . By $\text{diam}(G)$ we denote the diameter of G .

Theorem 1. *Let G be a graph. If $d_p(G) \geq 3$, then $\text{diam}(G) \leq 2$.*

Proof. Let $d_p(G) = k \geq 3$. Then there exists a ps -domatic partition $\{D_1, \dots, D_k\}$ of G . Let x, y be two vertices of G . As $k \geq 3$, at least one of the sets D_1, \dots, D_k contains neither x nor y . Without loss of generality let it be D_1 . We have $\{x, y\} \subseteq V(G) - D_1$ and therefore there exists a vertex $v \in D_1$ such that $\{v, x, y\}$ induces a connected subgraph of G . If x, y are adjacent, then $d_G(x, y) = 1$. If x, y are not adjacent, then v must be adjacent to both x and y and $d_G(x, y) = 2$. As x, y were chosen arbitrarily, we have $\text{diam}(G) \leq 2$. \square

Theorem 2. *Let G be a graph. If $d_p(G) = 2$, then $\text{diam}(G) \leq 3$.*

Proof. Let $d_p(G) = 2$. There exists a ps -domatic partition $\{D_1, D_2\}$ of $V(G)$. Let x, y be two vertices of G . If both x, y are in D_1 , then $\{x, y\} \subseteq V(G) - D_2$ and $d_G(x, y) \leq 2$ analogously as in the proof of Theorem 1. Similarly in the case when both x, y are in D_2 . Now let $x \in D_1, y \in D_2$. As $\{y\} \subseteq V(G) - D_1$, there exists $v \in D_1$ adjacent to y . As both x, v are in D_1 , we have $d_G(x, v) \leq 2, d_G(v, y) = 1$ and thus $d_G(x, y) \leq 3$. As x, y were chosen arbitrarily, we have $\text{diam}(G) \leq 3$. \square

Now we shall consider bipartite graphs.

Corollary. *Let G be a bipartite graph. If $d_p(G) \geq 3$, then G is a complete bipartite graph.*

This follows from the fact that every non-complete bipartite graph has the diameter at least 3.

Theorem 3. *Let G be a non-complete bipartite graph. Then $d_p(G) = 2$ if and only if G has a spanning tree T with $\text{diam}(T) \leq 3$.*

Proof. Let T be a tree with $\text{diam}(T) \leq 3$. If D_1, D_2 are the bipartition classes of T , then $\{D_1, D_2\}$ is a ps -domatic partition of T and $d_p(T) \leq 2$ and thus $d_p(G) = 2$. If G is a graph such that T is its spanning tree and G is a non-complete bipartite graph, then obviously also $d_p(G) = 2$.

Now suppose that $d_p(G) = 2$ and let $\{D_1, D_2\}$ be a ps -domatic partition. Let V_1, V_2 be the bipartition classes of G . First suppose that D_1 is a proper subset of V_1 . Then $V_1 - D_1 \subseteq V(G) - D_1$ and for each $v \in D_1$ the set $(V_1 - D_1) \cup \{v\}$ is independent, i.e. it does not induce a connected subgraph of G . Hence this case is impossible and moreover D_1 cannot be a proper subset of V_2 and D_2 cannot be a proper subset of V_1

or of V_2 . Now consider the case $D_1 = V_1$. Then $D_2 = V_2$. We have $V_2 \subseteq V(G) - D_1$ and there exists a vertex $v_1 \in V_1$ adjacent to all vertices of V_2 . Analogously, there exists a vertex $v_2 \in V_2$ adjacent to all vertices of V_1 . All edges joining v_1 with vertices of V_2 and all edges joining v_2 with vertices of V_1 form the spanning tree T ; its central edge is v_1v_2 and its diameter is 3. The case $D_1 = V_2, D_2 = V_1$ is analogous. Now the case remains when $D_1 \cap V_1 \neq \emptyset, D_1 \cap V_2 \neq \emptyset, D_2 \cap V_1 \neq \emptyset, D_2 \cap V_2 \neq \emptyset$. Let $V_1 \in D_1 \cap V_1, x_2 \in D_1 \cap V_2$. We have $\{x_1, x_2\} \subseteq V(G) - D_2$ and there exists a vertex $v \in D_2$ such that $\{v, x_1, x_2\}$ induces a connected subgraph of G . As x_1, x_2 belong to distinct bipartition classes of G , the vertex v cannot be adjacent to both of them and thus x_1, x_2 are adjacent. Therefore D_2 induces a complete bipartite subgraph on the sets $D_2 \cap V_1, D_2 \cap V_2$ and analogously, D_1 induces a complete bipartite subgraph on the sets $D_1 \cap V_1, D_1 \cap V_2$. We have $D_1 \cap V_1 \subseteq V(G) - D_2$ and therefore there exists a vertex $w_2 \in D_2$ adjacent to all vertices of $D_2 \cap V_1$; evidently $w_2 \in D_2 \cap V_2$. Analogously, there exists a vertex $w_1 \in D_1 \cap V_1$ adjacent to all vertices of $D_1 \cap V_2$. The vertex w_1 is adjacent to all vertices of V_2 and the vertex w_2 is adjacent to all vertices of V_1 . Obviously w_1, w_2 are adjacent. There exists a spanning tree T with the central edge w_1w_2 which has the diameter 3. \square

Now we turn to circuits. By C_n we denote the circuit of the length n .

Theorem 5. *For the circuits we have*

$$\begin{aligned} d_p(C_3) &= 3, \\ d_p(C_4) &= 2, \\ d_p(C_5) &= 2, \\ d_p(C_n) &= 1 \quad \text{for } n \geq 6. \end{aligned}$$

Proof. The circuit C_3 is the complete graph K_3 and thus $d_p(C_3) = 3$. The circuit C_4 contains a spanning tree which is a path P_3 of length 3 and therefore $d_p(C_4) = 2$; note that C_4 is a bipartite graph. Consider C_5 and let its vertices be u_1, \dots, u_5 and edges u_iu_{i+1} for $i = 1, \dots, 4$ and u_5u_1 . There exists a ps -domatic partition $\{D_1, D_2\}$, where $D_1 = \{u_1, u_2, u_4\}, D_2 = \{u_3, u_5\}$; thus $d_p(C_5) \geq 2$. As the domatic number $d(C_5) = 2$, we have $d_p(C_5) = 2$ as well. The circuit C_6 is a bipartite graph and does not contain any spanning tree of diameter 3, therefore $d_p(C_6) = 1$. Now consider C_7 . Suppose that in C_7 there exists a ps -domatic partition $\{D_1, D_2\}$ and denote its vertices by u_1, \dots, u_7 in the usual way. Any two vertices with the distance 3 are in distinct classes of $\{D_1, D_2\}$; this follows from the proofs of Theorem 1 and of Theorem 2. If $u_1 \in D_1$ (without loss of generality), then $u_4 \in D_2, u_7 \in D_1, u_3 \in D_2, u_6 \in D_1, u_2 \in D_2, u_5 \in D_1, u_1 \in D_2$, which is a contradiction and thus $d_p(C_7) = 1$. For $n \geq 8$ we have $\text{diam}(C_n) \geq 4$ and thus $d_p(C_n) = 1$. \square

Theorem 6. For the complement \bar{C}_n of a circuit C_n we have

$$\begin{aligned}d_p(\bar{C}_3) &= 1, \\d_p(\bar{C}_4) &= 1, \\d_p(\bar{C}_n) &= \lfloor n/2 \rfloor \quad \text{for } n \geq 5.\end{aligned}$$

Proof. The graphs \bar{C}_3 and \bar{C}_4 are disconnected and therefore they have the ps -domatic number 1. If $n \geq 5$, then any pair of non-adjacent vertices in \bar{C}_n is a ps -dominating set, which can be easily verified by the reader. There exists a partition of $V(\bar{C}_n)$ into $\lfloor n/2 \rfloor$ sets, each of which is a pair of non-adjacent vertices, except at most one which has three vertices from which only two are adjacent. There exists no one-element ps -dominating set, therefore $d_p(\bar{C}_n) = \lfloor n/2 \rfloor$. \square

In the end we will prove an existence theorem.

Theorem 7. Let V be a finite set, let k be an integer, $1 \leq k \leq |V|$, let $\{D_1, \dots, D_k\}$ be a partition of V . Then there exists a graph G such that $V(G) = V$, $d_p(G) = k$ and $\{D_1, \dots, D_k\}$ is a ps -domatic partition of G .

Proof. For $i = 1, \dots, k$ choose a vertex $v_i \in D_i$ and join it by edges with all vertices not belonging to D_i . The resulting graph is the graph G . For each subset $S \subseteq V(G) - D_i$ there exists a vertex of D_i which is adjacent to all vertices of S , namely v_i . Therefore $\{D_1, \dots, D_k\}$ is a ps -domatic partition of G and $d_p(G) \geq k$. If $|D_i| = 1$ for all i , then G is K_k and $d_p(G) = k$. If $|D_i| \geq 2$ for some i , then a vertex $u \in D_i - \{v_i\}$ has the degree $k - 1$ and thus the domatic number satisfies $d(G) \leq k$ (by [1]) and $d_p(G) \leq d(G) \leq k$. This implies $d_p(G) = k$. \square

In the end we will give a motivation for introducing the point-set domination. The concept of a dominating set is usually motivated by the displacement of certain service stations (medical, police, fire-brigade) which have to provide service for certain places (vertices of a graph). In the case of the point-set dominating set we want that for any chosen region (set of vertices) there might exist a station providing services for the whole region. Note that the point-set domination number is also a variant of the set domination number introduced in [5] and mentioned in [2].



References

- [1] *Cockayne E. J., Hedetniemi S. T.*: Towards the theory of domination in graphs. *Networks* 7 (1977), 247–261.
- [2] *Haynes T. W., Hedetniemi S. T., Slater P. J.*: *Fundamentals of Domination in Graphs*. Marcel Dekker, Inc., New York, 1998.
- [3] *Pushpa Latha L.*: The global point-set domination number of a graph. *Indian J. Pure Appl. Math.* 28 (1997), 47–51.
- [4] *Sampathkumar E., Pushpa Latha L.*: Point-set domination number of a graph. *Indian J. Pure Appl. Math.* 24 (1993), 225–229.
- [5] *Sampathkumar E., Pushpa Latha L.*: Set domination in graphs. *J. Graph Theory* 18 (1994), 489–495.

Author's address: Bohdan Zelinka, Katedra aplikované matematiky Technické univerzity, Voroněžská 13, 461 17 Liberec 1, Czech Republic.