

Aleksander Maliszewski

Averages of quasi-continuous functions

Mathematica Bohemica, Vol. 124 (1999), No. 1, 29–34

Persistent URL: <http://dml.cz/dmlcz/125978>

Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

AVERAGES OF QUASI-CONTINUOUS FUNCTIONS

ALEKSANDER MALISZEWSKI, Bydgoszcz¹

(Received May 19, 1997)

Abstract. The goal of this paper is to characterize the family of averages of comparable (Darboux) quasi-continuous functions.

Keywords: cliquishness, quasi-continuity, Darboux property, comparable functions, average of functions

MSC 1991: 26A15, 54C08

PRELIMINARIES

The letters \mathbb{R} , \mathbb{Q} and \mathbb{N} denote the real line, the set of rationals and the set of positive integers, respectively. The word *function* denotes a mapping from \mathbb{R} into \mathbb{R} . We say that functions φ and ψ are *comparable* if either $\varphi < \psi$ on \mathbb{R} or $\varphi > \psi$ on \mathbb{R} . For each $A \subset \mathbb{R}$ we use the symbols $\text{cl } A$ and $\text{bd } A$ to denote the closure and the boundary of A , respectively.

Let f be a function. If $A \subset \mathbb{R}$ is nonvoid, then let $\omega(f, A)$ be the *oscillation of f on A* , i.e., $\omega(f, A) = \sup\{|f(x) - f(t)| : x, t \in A\}$. For each $x \in \mathbb{R}$ let $\omega(f, x)$ be the *oscillation of f at x* , i.e., $\omega(f, x) = \lim_{\delta \rightarrow 0^+} \omega(f, (x - \delta, x + \delta))$. The symbol \mathcal{C}_f denotes the set of points of continuity of f .

We say that a function f is *quasi-continuous* in the sense of Kempisty [4] (*cliquish* [10]) at a point $x \in \mathbb{R}$ if for each $\varepsilon > 0$ and each open set $U \ni x$ there is a nonvoid open set $V \subset U$ such that $\omega(f, \{x\} \cup V) < \varepsilon$ ($\omega(f, V) < \varepsilon$ respectively). We say that f is *quasi-continuous (cliquish)* if it is quasi-continuous (cliquish) at each point $x \in \mathbb{R}$. Cliquis functions are also known as *pointwise discontinuous*.

¹ Supported by BW grant, WSP.

Let I be an interval and $f: I \rightarrow \mathbb{R}$. We say that f is *Darboux* if it has the intermediate value property. We say that f is *strong Świątkowski* [6] if whenever $a, b \in I$, $a < b$, and y is a number between $f(a)$ and $f(b)$, there is an $x \in (a, b) \cap \mathcal{C}_f$ with $f(x) = y$. One can easily verify that strong Świątkowski functions are both Darboux and quasi-continuous, and that the converse is not true.

For brevity, if f is a cliquish function and $x \in \mathbb{R}$, then we define

$$\underline{\text{LIM}}(f, x) = \underline{\lim}_{t \rightarrow x, t \in \mathcal{C}_f} f(t).$$

The symbols $\underline{\text{LIM}}(f, x^-)$ and $\underline{\text{LIM}}(f, x^+)$ are defined analogously.

INTRODUCTION

In 1974 A. M. Bruckner, J. G. Ceder, and T. L. Pearson characterized the averages of comparable Darboux functions [1, Theorem 2]. In this paper we solve an analogous problem, namely we characterize the averages of comparable quasi-continuous functions.

A similar problem is to determine a necessary and sufficient condition that for a function f there exists a quasi-continuous function ψ such that $\psi > f$ on \mathbb{R} . (The answer to this question for Darboux functions can be easily obtained using the proof of [1, Theorem 2].) In both cases we ask whether there is a positive function g such that both $f + g$ and $-f + g$ are quasi-continuous (the first problem) or such that $f + g$ is quasi-continuous (the second problem). This suggests a similar problem for larger classes of functions. Theorem 4.1 contains a solution of this problem for finite classes of cliquish functions. Recall that by [5, Example 2], we cannot in general allow infinite families in Theorem 4.1. Unlike [7, Theorem 4], we cannot conclude in condition (ii) of Theorem 4.1 that g is a Baire one function; actually, we cannot even conclude that g is Borel measurable (Corollary 4.5).

The Baire class one case makes no difficulty if we require only quasi-continuity of the sums, but it needs a separate argument if we require both the Darboux property and the quasi-continuity. Notice that by Proposition 4.3, the necessary and sufficient condition for Darboux quasi-continuous Baire one functions is essentially stronger.

AUXILIARY LEMMAS

Lemma 3.1. *If f is a cliquish function, then the mapping $x \mapsto \underline{\text{LIM}}(f, x)$ is lower semicontinuous, while the mapping $x \mapsto \underline{\text{LIM}}(f, x^-)$ belongs to Baire class two.*

Proof. Let $y \in \mathbb{R}$. For every $x \in \mathbb{R}$, if $\underline{\text{LIM}}(f, x) > y$, then there exist an open interval $I_x \ni x$ and a rational $q_x > y$ such that $f > q_x$ on $\mathcal{C}_f \cap I_x$, whence $\underline{\text{LIM}}(f, t) \geq q_x > y$ for each $t \in I_x$. Thus the set $\{x \in \mathbb{R} : \underline{\text{LIM}}(f, x) > y\}$ is open.

To prove the other assertion put $A_y = \{x \in \mathbb{R} : \underline{\text{LIM}}(f, x^-) > y\}$ for each $y \in \mathbb{R}$. Let $y \in \mathbb{R}$. If $x \in A_y$, then proceeding as above we can find a closed interval $I_x \subset A_y$ with $x \in I_x$. So $A_y \cap \text{bd } A_y$ is at most countable. Hence A_y is an F_σ set, while $\{x \in \mathbb{R} : \underline{\text{LIM}}(f, x^-) < y\} = \bigcup \{\mathbb{R} \setminus A_q : q < y, q \in \mathbb{Q}\}$ is the difference of an F_σ set and a countable one. \square

Lemma 3.2. *Let $I = [a, b]$ and $n \in \mathbb{N}$. Suppose that functions f_1, \dots, f_k are cliquish and $\max\{\omega(f_1, I), \dots, \omega(f_k, I)\} < 1$. There is a positive Baire one function g such that $g = 1$ on $\text{bd } I$, $\mathcal{C}_g \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$, and for each i the function $(f_i + g) \upharpoonright I$ is strong Świątkowski and*

$$(f_i + g) \left[I \cap \bigcap_{i=1}^k \mathcal{C}_{f_i} \right] \supset [\inf f_i[I] + 1, \max\{\inf f_i[I] + 1, n\}].$$

Proof. Put $T = \max\{n - \inf f_i[I] : i \in \{1, \dots, k\}\} + 1$. Construct a nonnegative continuous function φ such that $\varphi[I] = [0, T]$ and $\varphi = 0$ outside of I . For each i define $\tilde{f}_i(x) = (f_i + \varphi)(x)$ if $x \in I$, and let \tilde{f}_i be constant on $(-\infty, a)$ and $[b, \infty)$. By [7, Theorem 4], there is a Baire one function \tilde{g} such that $\tilde{f}_i + \tilde{g}$ is strong Świątkowski for each i (see condition (8) in the proof of [7, Theorem 4]), $\mathcal{C}_{\tilde{g}} \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$, and $|\tilde{g}| < 1$ on \mathbb{R} ; by its proof, we can conclude that $\tilde{g} = 0$ on $\{a, b\}$. Put $g = \varphi + \tilde{g} + 1$. Then for each i , since $\tilde{f}_i + \tilde{g}$ is strong Świątkowski and $f_i + g = \tilde{f}_i + \tilde{g} + 1$ on I , we have

$$\begin{aligned} (f_i + g) \left[I \cap \bigcap_{i=1}^k \mathcal{C}_{f_i} \right] &\supset (\inf(f_i + g)[I], \sup(f_i + g)[I]) \\ &\supset (f_i(a), \inf f_i[I] + \sup g[I]) \\ &\supset [\inf f_i[I] + 1, \max\{\inf f_i[I] + 1, n\}]. \end{aligned}$$

The other requirements are evident. \square

MAIN RESULTS

Theorem 4.1. *Let \mathcal{F} be one of the following classes of functions: all cliquish functions, Lebesgue measurable cliquish functions, cliquish functions in Baire class α ($\alpha \geq 1$), and suppose $f_1, \dots, f_k \in \mathcal{F}$. The following are equivalent:*

- (i) *there is a positive function g such that $f_i + g$ is quasi-continuous for each i ;*
- (ii) *there is a positive function $g \in \mathcal{F}$ such that $\mathcal{C}_g \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$ and $f_i + g$ is quasi-continuous for each i ;*
- (iii) *for each $x \in \mathbb{R}$ and each i we have $\underline{\text{LIM}}(f_i, x) < \infty$.*

Proof. The implication (ii) \Rightarrow (i) is obvious.

(i) \Rightarrow (iii). Let $x \in \mathbb{R}$ and $i \in \{1, \dots, k\}$. Since $f_i + g$ is quasi-continuous, so by [2] (see also [3, Lemma 2]) we obtain

$$\underline{\text{LIM}}(f_i, x) \leq \underline{\text{LIM}}(f_i + g, x) \leq (f_i + g)(x) < \infty.$$

(iii) \Rightarrow (ii). Put $A = \bigcup_{i=1}^k \{x \in \mathbb{R} : \omega(f_i, x) \geq 1\}$. Then A is closed and nowhere dense. Find a family $\{I_n : n \in \mathbb{N}\}$ consisting of nonoverlapping compact intervals, such that $\bigcup_{n \in \mathbb{N}} I_n = \mathbb{R} \setminus A$ and each $x \notin A$ is an interior point of $I_n \cup I_m$ for some $n, m \in \mathbb{N}$. Since each I_n is compact and $\omega(f_i, x) < 1$ for each $x \in I_n$ and $i \in \{1, \dots, k\}$, so we may assume that $\omega(f_i, I_n) < 1$ for each i and n . For each $n \in \mathbb{N}$ use Lemma 3.2 to construct a positive Baire one function g_n such that $g_n = 1$ on $\text{bd } I_n$, $\mathcal{C}_{g_n} \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$, and for each i the function $(f_i + g_n) \upharpoonright I_n$ is strong Świątkowski and

$$(*) \quad (f_i + g_n) \left[I_n \cap \bigcap_{i=1}^k \mathcal{C}_{f_i} \right] \supset [\inf f_i|_{I_n} + 1, \max\{\inf f_i|_{I_n} + 1, n\}].$$

Define $g(x) = g_n(x)$ if $x \in I_n$ for some $n \in \mathbb{N}$, and

$$g(x) = \max\{\max\{\underline{\text{LIM}}(f_i, x) - f_i(x) : i \in \{1, \dots, k\}\}, 0\} + 1$$

if $x \in A$. By Lemma 3.1, each mapping $x \mapsto \underline{\text{LIM}}(f_i, x)$ is Baire one, so $g \in \mathcal{F}$.

Fix an $i \in \{1, \dots, k\}$. Clearly $f_i + g$ is quasi-continuous outside of A . On the other hand, if $x \in A$, then by $(*)$, for each $\delta > 0$ we have

$$(f_i + g)[(x - \delta, x + \delta) \cap \mathcal{C}_{f_i + g}] \supset (\underline{\text{LIM}}(f_i, x) + 1, \infty).$$

Hence $f_i + g$ is quasi-continuous. □

Theorem 4.2. Let \mathcal{F} be one of the following classes of functions: all cliquish functions, Lebesgue measurable cliquish functions, cliquish functions in Baire class α ($\alpha \geq 2$), and suppose $f_1, \dots, f_k \in \mathcal{F}$. The following are equivalent:

- (i) there is a positive function g such that $f_i + g$ is both Darboux and quasi-continuous for each i ;
- (ii) there is a positive function $g \in \mathcal{F}$ such that $\mathcal{C}_g \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$ and $f_i + g$ is strong \acute{S} wiątkowski for each i ;
- (iii) for each $x \in \mathbb{R}$ and each i we have $\max\{\underline{\text{LIM}}(f_i, x^-), \underline{\text{LIM}}(f_i, x^+)\} < \infty$.

Proof. The proof of the implication (iii) \Rightarrow (ii) is a repetition of the argument used in Theorem 4.1, and the implication (ii) \Rightarrow (i) is obvious.

(i) \Rightarrow (iii). Let $x \in \mathbb{R}$ and $i \in \{1, \dots, k\}$. Since $f_i + g$ is both Darboux and quasi-continuous, so by [9, Lemma 2] we obtain

$$\underline{\text{LIM}}(f_i, x^-) \leq \underline{\text{LIM}}(f_i + g, x^-) \leq (f_i + g)(x) < \infty.$$

Similarly $\underline{\text{LIM}}(f_i, x^+) < \infty$. □

Proposition 4.3. There is a Baire one function f such that $f + g$ is strong \acute{S} wiątkowski for some positive function g in Baire class two, but $f + g$ is Darboux for no positive Baire one function g .

Proof. Let F be the Cantor ternary set and let $\mathcal{S} = \{(a_n, b_n) : n \in \mathbb{N}\}$ and \mathcal{J} be disjoint families of components of $\mathbb{R} \setminus F$ such that $F = (\text{cl} \bigcup \mathcal{S}) \cap (\text{cl} \bigcup \mathcal{J})$. Define $f(x) = n$ if $x \in (a_n, b_n)$ for some $n \in \mathbb{N}$ and $f(x) = 0$ otherwise. Clearly f belongs to Baire class one.

Let $x \in \mathbb{R}$. If $x \in (a_n, b_n]$ for some $n \in \mathbb{N}$, then $\underline{\text{LIM}}(f, x^-) = n$, otherwise $\underline{\text{LIM}}(f, x^-) = 0$. Similarly $\underline{\text{LIM}}(f, x^+) < \infty$. By Theorem 4.2 there is a positive Baire two function g such that $f + g$ is strong \acute{S} wiątkowski.

On the other hand, by [8, Proposition 6.10], $f + g$ is Darboux for no positive Baire one function g . □

In Proposition 4.4 the symbol \mathfrak{c} denotes the first ordinal equipollent with \mathbb{R} .

Proposition 4.4. Given a family of positive functions, $\{g_\xi : \xi < \mathfrak{c}\}$, we can find a cliquish function f which fulfils condition (iii) of Theorem 4.2 and such that $f + g_\xi$ is not quasi-continuous for each $\xi < \mathfrak{c}$.

Proof. Let F be the Cantor ternary set and let $\{x_\xi : \xi < \mathfrak{c}\}$ be an enumeration of F . Define $f(x) = -g_\xi(x) - 1$ if $x = x_\xi$ for some $\xi < \mathfrak{c}$, and $f(x) = 0$ otherwise. Clearly f is cliquish, and for each $x \in \mathbb{R}$ we have $\underline{\text{LIM}}(f, x^-) = \underline{\text{LIM}}(f, x^+) = 0$.

Let $\xi < \mathfrak{c}$. Then $(f + g_\xi)(x_\xi) = -1$ and $f + g_\xi$ is positive on a dense open set. Thus $f + g_\xi$ is not quasi-continuous at x_ξ . □

Corollary 4.5. *There is a cliquish function f which fulfils condition (iii) of Theorem 4.2 and such that $f + g$ is not quasi-continuous for each positive Borel measurable function g .*

Theorem 4.6. *Let f_1, \dots, f_k be Baire one functions. The following are equivalent:*

- (i) *there is a positive Baire one function g such that $f_i + g$ is both Darboux and quasi-continuous for each i ;*
- (ii) *there is a positive Baire one function g such that $\mathcal{C}_g \supset \bigcap_{i=1}^k \mathcal{C}_{f_i}$ and $f_i + g$ is strong Świątkowski for each i ;*
- (iii) *there is a Baire one function h such that for each $x \in \mathbb{R}$ and each i we have $\max\{\underline{\text{LIM}}(f_i, x^-), \underline{\text{LIM}}(f_i, x^+)\} \leq h(x)$.*

Proof. The implication (i) \Rightarrow (iii) can be proved similarly as in Theorem 4.2 (we let $h = \max\{f_1, \dots, f_k\} + g$), and the implication (ii) \Rightarrow (i) is obvious.

(iii) \Rightarrow (ii). The proof of this implication is a repetition of the argument used in Theorem 4.1. The only difference is in the definition of the function g on the set A . More precisely, we put

$$g(x) = \max\{\max\{h(x) - f_i(x) : i \in \{1, \dots, k\}\}, 0\} + 1$$

if $x \in A$. Then clearly g is a Baire one function. □

References

- [1] *A. M. Bruckner, J. G. Ceder, T. L. Pearson: On Darboux functions. Rev. Roumaine Math. Pures Appl. 19 (1974), 977–988.*
- [2] *Z. Grande: Sur la quasi-continuité et la quasi-continuité approximative. Fund. Math. 129 (1988), 167–172.*
- [3] *Z. Grande, T. Natkaniec: Lattices generated by \mathcal{F} -quasi-continuous functions. Bull. Polish Acad. Sci. Math. 34 (1986), 525–530.*
- [4] *S. Kempisty: Sur les fonctions quasicontinues. Fund. Math. 19 (1932), 184–197.*
- [5] *A. Maliszewski: Sums of bounded Darboux functions. Real Anal. Exchange 20 (1994–95), 673–680.*
- [6] *A. Maliszewski: On the limits of strong Świątkowski functions. Zeszyty Nauk. Politech. Łódź. Mat. 27 (1995), 87–93.*
- [7] *A. Maliszewski: On theorems of Pu & Pu and Grande. Math. Bohem. 121 (1996), 83–87.*
- [8] *A. Maliszewski: On the averages of Darboux functions. Trans. Amer. Math. Soc. 350 (1998), 2833–2846.*
- [9] *T. Natkaniec: On quasi-continuous functions having Darboux property. Math. Pann. 3 (1992), 81–96.*
- [10] *H. P. Thielman: Types of functions. Amer. Math. Monthly 60 (1953), 156–161.*

Author's address: Aleksander Maliszewski, Department of Mathematics, Pedagogical University, pl. Weyssenhoffa 11, 85-072 Bydgoszcz, Poland, e-mail: AMal@wp.bydgoszcz.pl.