

Rastislav Jurga

The cross-ratio in Hjelmslev planes

Mathematica Bohemica, Vol. 122 (1997), No. 3, 243–247

Persistent URL: <http://dml.cz/dmlcz/126149>

Terms of use:

© Institute of Mathematics AS CR, 1997

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE CROSS-RATIO IN HJELMSLEV PLANES

RASTISLAV JURGA, Košice

(Received January 31, 1996)

Summary. The cross-ratio in Hjelmslev planes is defined. The cross-ratio in the Hjelmslev plane $H(R)$ is independent of the choice of a coordinate system on a line.

Keywords: Hjelmslev plane over a special local ring, cross-ratio in Hjelmslev plane

MSC 1991: 51C05, 51E30

1. INTRODUCTION

A special local ring is a finite commutative local ring R the ideal I of divisors of zero of which is principal. Suppose that R is not a field and that the characteristic of R is odd. Denote the factor ring R/I by the symbol \bar{R} . Further denote the set of all regular elements of R by the symbol R^* , thus $R^* = R - I$.

Definition 1.1. A projective Hjelmslev plane (we will denote it by $H(R)$) over R is an incidence structure $H(R) = (\mathcal{B}; \mathcal{P}; \mathcal{I})$ defined in the following way:

- the elements of \mathcal{B} —the points of $H(R)$ are classes of ordered triples $(\lambda x_1; \lambda x_2; \lambda x_3)$ where $\lambda \in R^*$, $x_1, x_2, x_3 \in R$ and at least one x_i is regular;
- the elements of \mathcal{P} —the lines of $H(R)$ are classes of ordered triples $(\alpha a_1; \alpha a_2; \alpha a_3)$ where $\alpha \in R^*$, $a_1, a_2, a_3 \in R$ and at least one a_i is regular.

A point $X = [x_1; x_2; x_3]$ is incident to the line $a = [a_1; a_2; a_3]$ if and only if

$$(1.1) \quad a_1 x_1 + a_2 x_2 + a_3 x_3 = 0.$$

Remark 1.1. The canonical homomorphism $\Phi: R \rightarrow R/I = \bar{R}$ induces a homomorphism of $H(R)$ onto the projective plane $\pi(\bar{R})$.

We will call the points $X, Y \in H(R)$ *neighbouring* if $\overline{X} = \overline{Y}$ where $\Phi(X) = \overline{X}$ $\Phi(Y) = \overline{Y}$. Similarly we will call points $X, Y \in H(R)$ *substantially different* if $\overline{X} \neq \overline{Y}$. Two lines are neighbouring if there are points $A_1, A_2 \in \mathcal{B}$, $A_1 \neq A_2$ such that $A_1 \mathcal{I}a, b$ and $A_2 \mathcal{I}a, b$. Let X be a subset of the R -modul M and let $j: X \rightarrow M$ be an insertion of the subset X into M . Then $M(R)$ is called the free modul over X if for an arbitrary function $f: X \rightarrow A$ into the R -modul A there is exactly one linear mapping $t: M(R) \rightarrow A$ such that $t \circ j = f$.

Remark 1.2. The analytic model of the Hjelmslev plane, introduced by definition 1.1 is really a free modul over R with a factorization defined in the following way: triples $(x_1; x_2; x_3)$ and $(x'_1; x'_2; x'_3)$ are identical if there is $\lambda \in R^*$ such that $x'_i = \lambda x_i$ for $i = 1, 2, 3$ and we do not consider the zero triple.

2. THE CONSTRUCTION AND PROOF OF THEOREM

Definition 2.1. A coordinate system in $H(R)$ is an ordered quadruple of points E_1, E_2, E_3, E_4 such that the points $\overline{E}_1, \overline{E}_2, \overline{E}_3, \overline{E}_4$ generate a coordinate system in $\pi(\overline{R})$.

If a point $X = [x_1; x_2; x_3]$ is given by the vector $x = (x_1; x_2; x_3)$, we write $X = \langle x \rangle$.

Lemma 2.1. Let $M(R)$ be a free modul over R and let e_1, e_2, e_3 be a basis of $M(R)$. Then the points $E_1 = \langle e_1 \rangle$, $E_2 = \langle e_2 \rangle$, $E_3 = \langle e_3 \rangle$, $E_4 = \langle e_1 + e_2 + e_3 \rangle$ generate the coordinate system in the Hjelmslev plane $H(R)$ corresponding to the modul $M(R)$.

Proof. It is necessary to prove that the points $\overline{E}_1, \overline{E}_2, \overline{E}_3, \overline{E}_4$ generate a coordinate system in $\pi(\overline{R})$. Obviously $\overline{e}_1, \overline{e}_2, \overline{e}_3$ form a basis of a vector space over \overline{R} and thus the vectors $\overline{e}_1, \overline{e}_2, \overline{e}_3$ are linearly independent. It follows that the points $\overline{E}_1 = \langle \overline{e}_1 \rangle$, $\overline{E}_2 = \langle \overline{e}_2 \rangle$, $\overline{E}_3 = \langle \overline{e}_3 \rangle$ and $\overline{E}_4 = \langle \overline{e}_1 + \overline{e}_2 + \overline{e}_3 \rangle$ are not on a unique line. \square

Conversely, we have

Lemma 2.2. Let E_1, E_2, E_3, E_4 be a coordinate system in $H(R)$. Then there is a basis of the modul $M(R)$ such that $\langle e_1 \rangle = E_1$, $\langle e_2 \rangle = E_2$, $\langle e_3 \rangle = E_3$, $\langle e_1 + e_2 + e_3 \rangle = E_4$.

Proof. Let $E_1 = \langle b_1 \rangle$, $E_2 = \langle b_2 \rangle$, $E_3 = \langle b_3 \rangle$ and $E_4 = \langle b_4 \rangle$. Because $\{b_1, b_2, b_3\}$ is a basis of $M(R)$ the vector b_4 can be expressed in the form

$$b_4 = \beta_1 b_1 + \beta_2 b_2 + \beta_3 b_3.$$

If we denote $e_1 = \beta_1 b_1$, $e_2 = \beta_2 b_2$, $e_3 = \beta_3 b_3$ then e_1, e_2, e_3 are the vectors from the statement of the lemma.

Let E_1, E_2, E_3, E_4 and E'_1, E'_2, E'_3, E'_4 be coordinate systems in $H(R)$. If e_1, e_2, e_3 and e'_1, e'_2, e'_3 are the corresponding bases of the modul $M(R)$ then there is a regular matrix $A = [a_{ij}]$ such that

$$e'_i = \sum_j a_{ij} e_j, \quad i = 1, 2, 3.$$

Let $X_E = [x_1; x_2; x_3]$, $X'_E = [x'_1; x'_2; x'_3]$. Then

$$x = \sum_i x'_i e'_i = \sum_i x'_i \sum_j a_{ij} e_j = \sum_j \left(\sum_i x'_i a_{ij} \right) e_j = \sum_j x_j e_j.$$

Comparing the two identities, we get

$$(2.1) \quad x_j = \sum_i x'_i a_{ij}.$$

The relation (2.1) can be written also in the form

$$(2.2) \quad X_E = X'_E A, \quad X'_E = X_E A^{-1}.$$

Let an invertible matrix A and a coordinate system E_1, E_2, E_3, E_4 be given, then points E'_1, E'_2, E'_3, E'_4 generate a coordinate system and the corresponding vectors of the point $X \in H(R)$ satisfy

$$X_E = X'_E A.$$

Let the special local ring R be given. We introduce a set Ω by

$$(2.3) \quad \Omega \cap R = \emptyset, \quad |\Omega| = |I|.$$

Thus there is a bijective mapping ω such that

$$(2.4) \quad \omega: I \rightarrow \Omega, \quad \omega: i \rightarrow \omega_i = \omega(i), \quad i \in I$$

where ω_i are "inverse" elements of elements $i \in I$, thus $\omega_i \sim 1/i$. Ω is the set of "infinities" corresponding to singular elements. Define an extension of the canonical homomorphism Φ to the set $R \cup \Omega$, let us put

$$(2.5) \quad \Phi(\Omega) = \infty.$$

Let A, B, E be three substantially different points generating a coordinate system on a line. Then every point X of this line can be expressed uniquely (the single-valuedness guarantees the point E) in the form

$$(2.6) \quad X = sA + tB$$

and hence the point $X = [s; t]$ is determined by the pair $(s; t)$.

On the line with the coordinate system A, B, E let us have points P_1, P_2, P_3, P_4 where $P_i = s_i A + t_i B$ thus $P_i[s_i; t_i]$. \square

Definition 2.2. The cross-ratio of an ordered quadruple of points P_1, P_2, P_3, P_4 on a line in $H(R)$, of which at least three are substantially different is an element $(P_1 P_2, P_3 P_4) \in R \cup \Omega$ which is defined by relations

$$(2.7) \quad (P_1 P_2, P_3 P_4) = \frac{\begin{vmatrix} s_1 & t_1 \\ s_3 & t_3 \end{vmatrix} \begin{vmatrix} s_2 & t_2 \\ s_4 & t_4 \end{vmatrix}}{\begin{vmatrix} s_2 & t_2 \\ s_3 & t_3 \end{vmatrix} \begin{vmatrix} s_1 & t_1 \\ s_4 & t_4 \end{vmatrix}}$$

if points $P_1 P_4$ and $P_2 P_3$ are substantially different,

$$(2.8) \quad (P_1 P_2, P_3 P_4) = \omega(P_1 P_2, P_3 P_4)$$

if points P_1, P_4 and P_2, P_3 are neighbouring. Suppose that points P_1, P_3 and P_2, P_4 are substantially different.

Remark. If R is a field, $I = \{0\}$ then Definition 2.2 is the same as the definition of the cross-ratio in a projective plane.

Theorem 2.3. The cross-ratio introduced by relations 2.7 and 2.8 is independent of the choice of a coordinate system on the line.

Proof. Let a line $p \in H(R)$ be given and on this line let us have coordinate systems A, B, E and A', B', E' . Let P_1, P_2, P_3, P_4 be points on the given line p whose the cross-ratio we want to investigate. There is obviously a linear transformation which maps the points A, B to the points A', B' on p . We want to verify that the cross-ratio is independent of the choice of the coordinate points on the line. Thus

$$(P_1 P_2, P_3 P_4)_{AB} = (P_1 P_2, P_3 P_4)_{A'B'}$$

We have

$$\begin{aligned} A' &= a_1 A + a_2 B \\ B' &= b_1 A + b_2 B \end{aligned}$$

and thus

$$P_i = s_i' A' + t_i' B'$$

and after a substitution we get

$$P_i = (s_i' a_1 + t_i' b_1) A + (s_i' a_2 + t_i' b_2) B = s_i A + t_i B, \quad i = 1, 2, 3, 4.$$

By direct calculation we obtain $(P_1 P_2, P_3 P_4)_{AB} = (P_1 P_2, P_3 P_4)_{A'B'}$ which was to be proved. \square

References

- [1] *Dembowski P.*: Finite Geometries. Springer-Verlag, New York Inc., 1968.
- [2] *Ewald G.*: Geometry: An Introduction. Wadsworth Publishing Company, Inc., Belmont, California, 1971.
- [3] *Hjelmslev J.*: Die natürliche Geometrie. Leipzig, 1923.
- [4] *Hughes D. R., Piper F. C.*: Projective Planes. Springer-Verlag, New York, 1973.
- [5] *Jurga R.*: Some combinatorial properties of conics in the Hjelmslev plane. *Math. Slovaca* 45 (1995), no. 3, 219–226.

Author's address: Rastislav Jurga, Katedra aplikovanej matematiky, Podnikovohospodárska fakulta v Košiciach Ekonomickej univerzity v Bratislave, Tajovského 11, 041 30 Košice, Slovakia, e-mail: jurga@phf.euke.sk.