

Zhibo Chen

On minimum locally  $n$ -(arc)-strong digraphs

*Czechoslovak Mathematical Journal*, Vol. 46 (1996), No. 2, 317–323

Persistent URL: <http://dml.cz/dmlcz/127293>

## Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON MINIMUM LOCALLY  $n$ -(ARC)-STRONG DIGRAPHSZHIBO CHEN<sup>1</sup>, McKeesport

(Received September 9, 1994)

## 1. INTRODUCTION

Extensive studies have been devoted to the (global) connectedness in graphs and digraphs, one of the most important properties that a graph or digraph can possess (see, for instance, the surveys [2] and [8]). In 1974, G. Chartrand and R.E. Pippert [4] first defined locally connected and locally  $n$ -connected graphs and obtained some interesting results. Following [4], a variety of research [9–14] has been devoted to locally connected graphs. Recently, we first extended the study of local connectedness to digraphs (see [5] and [6]). In [5], we defined the locally  $n$ -strong digraphs and the locally  $n$ -arc-strong digraphs (See section 2 for definitions.), generalized some results of Chartrand and Pippert, and established relationships between local connectedness and global connectedness in digraphs, among which are the following theorems:

**Theorem A.** *Any weakly connected and locally  $n$ -arc-strong digraph is  $(n + 1)$ -arc-strong.*

**Theorem B.** *Any weakly connected and locally  $n$ -strong digraph is  $(n + 1)$ -strong.*

The aim of this paper is to further the study of locally  $n$ -(arc)-strong digraphs. We shall determine the minimum locally  $n$ -(arc)-strong digraphs and the minimum locally  $n$ -(arc)-strong oriented graphs. [Note: A minimum digraph with some property  $\mathcal{P}$  is a digraph with minimum number of arcs in the digraphs with the property  $\mathcal{P}$  which have minimum number of vertices.] Moreover, some results concerning tournaments are obtained, and the converses of the above Theorems A and B are shown to be not true.

---

<sup>1</sup> This research is supported in part by the RDG grant of the Pennsylvania State University.

## 2. DEFINITIONS

We follow the standard terminology and notation. A digraph  $D = (V(D), A(D))$  is a finite nonempty set  $V(D)$  of vertices together with a (possibly empty) set  $A(D)$  of ordered pairs of distinct vertices of  $D$  called arcs. An ordered pair  $(u, v) \in A(D)$  is also called an arc from  $u$  to  $v$ . A digraph  $D$  is said to be weakly connected if its underlying undirected graph is connected. If there is a dipath from  $u$  to  $v$  for any pair  $u$  and  $v$  of vertices in  $D$ , then the digraph  $D$  is said to be strongly connected, or simply said to be strong. The subdigraph induced by a nonempty subset  $W \subset V(D)$  is denoted  $\langle W \rangle_D$ . Let  $u, v \in V(D)$ . We say  $u$  is a neighbor of  $v$  if  $(u, v) \in A(D)$  or  $(v, u) \in A(D)$ . The set of neighbors of  $v$  in  $D$  is denoted  $N_D(v)$ . The induced subdigraph  $\langle N_D(v) \rangle_D$  is said to be the neighborhood of  $v$ . The outdegree of  $v$  is denoted as  $\text{od } v$  and the indegree of  $v$  is denoted as  $\text{id } v$ . Let  $\delta(D) = \min_{v \in V(D)} \{\text{id } v, \text{od } v\}$ . If  $\text{id } v = \text{od } v = \delta(D)$  for all  $v \in V(D)$ ,  $D$  is said to be diregular. Let  $S$  and  $T$  be two disjoint proper subsets of  $V(D)$ . We use  $(S, T)_D$  to denote the set of arcs  $(s, t)$  in  $D$  with  $s \in S$  and  $t \in T$ . When there is no confusion, we may simply use  $\langle W \rangle$ ,  $\langle N(v) \rangle$  and  $(S, T)$  to denote the corresponding  $\langle W \rangle_D$ ,  $\langle N_D(v) \rangle_D$  and  $(S, T)_D$ , respectively.

Let  $n \geq 1$ . A digraph  $D$  is said to be  $n$ -strong [ $n$ -arc-strong, resp.] if the removal of fewer than  $n$  vertices [arcs, resp.] always results in a nontrivial strong digraph. Clearly, every  $n$ -strong digraph is  $n$ -arc-strong. Every  $n$ -strong [ $n$ -arc-strong, resp.] digraph is also  $m$ -strong [ $m$ -arc-strong, resp.] for  $1 \leq m < n$ . It should also be noted that  $D$  is 1-strong iff  $D$  is 1-arc-strong iff  $D$  is a nontrivial strong digraph. The trivial strong digraph consisting of a single vertex is the only digraph that is strong but not 1-strong (or not 1-arc-strong).

A digraph  $D$  is said to be locally strong [locally  $n$ -strong, locally  $n$ -arc-strong, resp.] if the neighborhood of every vertex of  $D$  is strong [ $n$ -strong,  $n$ -arc-strong, resp.].

The associated digraph of a graph  $G$ , denoted as  $D(G)$ , is the digraph obtained from  $G$  when each edge  $e$  of  $G$  is replaced by a pair of oppositely oriented arcs with the same ends as  $e$ .

For other terminologies not defined here we refer the reader to the book [3].

## 3. MAIN RESULTS

**Theorem 1.** *The associated digraph  $D(K_{n+2})$  of the complete graph  $K_{n+2}$  is both the unique minimum locally  $n$ -strong digraph and the unique minimum locally  $n$ -arc-strong digraph.*

Before giving the proof of Theorem 1, we list some needed simple facts as the following propositions.

**Proposition 1.** *Let  $D$  be an  $n$ -(arc)-strong digraph. Then  $\delta(D) \geq n$ ,  $|V(D)| \geq n + 1$ , and  $|A(D)| \geq n(n + 1)$ .*

The proof is easy and is omitted here.

From Proposition 1, we immediately get

**Proposition 2.** *The associated digraph  $D(K_{n+1})$  is both the unique minimum  $n$ -strong digraph and the unique minimum  $n$ -arc-strong digraph.*

*Proof.* Clearly,  $D(K_{n+1})$  is  $n$ -strong and  $n$ -arc-strong. Both the vertex number and the arc number reach the lower bounds given in Proposition 1. □

**Proposition 3.** *Let  $D$  be a locally  $n$ -(arc)-strong digraph. Then  $\delta(D) \geq n + 1$ ,  $|V(D)| \geq n + 2$ , and  $|A(D)| \geq (n + 1)(n + 2)$ .*

*Proof.* By Theorem A and Proposition 1. □

Now the proof of Theorem 1 goes as follows.

*Proof of Theorem 1.* From Proposition 2,  $D(K_{n+2})$  is locally  $n$ -strong and locally  $n$ -arc-strong. Since both the vertex number and the arc number of  $D(K_{n+2})$  reach the lower bounds given in Proposition 3,  $D(K_{n+2})$  is a minimum locally  $n$ -strong and minimum locally  $n$ -(arc)-strong digraph.

The uniqueness is easily seen from the following:

If  $D$  is a minimum locally  $n$ -(arc)-strong digraph, then by Proposition 3,  $\delta(D) \geq n + 1$ . Note that  $|V(D)|$  must be not greater than the vertex number of  $D(K_{n+2})$ . Then,  $|V(D)| = n + 2$ . Thus we must have  $od v = id v = n + 1$  for all vertices in  $D$ . Therefore,  $D = D(K_{n+2})$ . □

Now we turn to determine the minimum locally  $n$ -(arc)-strong oriented graphs. Recall that a digraph is said to be an oriented graph if its underlying graph is a simple graph. Such digraphs are widely used in applications of graph theory.

**Theorem 2.** *A digraph  $D$  is a minimum locally  $n$ -arc-strong oriented graph if and only if  $D$  is a diregular tournament of  $2n + 3$  vertices.*

In the proof, we need the following lemmas where Lemma 1 is a rewritten version of a known result in [1].

**Lemma 1.** *Let  $D$  be an oriented graph. If  $\delta(D) \geq \left\lfloor \frac{|V(D)|+2}{4} \right\rfloor$ , then  $D$  is  $\delta(D)$ -arc-strong.*

**Lemma 2.** *Let  $D$  be a locally  $n$ -arc-strong oriented graph. Then  $\delta(D) \geq n + 1$ ,  $|V(D)| \geq 2n + 3$ , and  $|A(D)| \geq (n + 1)(2n + 3)$ .*

*Proof.* By Propositions 3,  $\delta(D) \geq n + 1$ . Then the other two inequalities immediately follow since  $D$  is an oriented graph.  $\square$

Now the proof of Theorem 2 goes as follows.

*Proof of Theorem 2.* We first prove the sufficiency. Let  $D$  be a diregular tournament of  $2n + 3$  vertices. By Lemma 1, it is easy to see that every neighborhood of a vertex in  $D$  is  $n$ -arc-strong. So,  $D$  is locally  $n$ -arc-strong. Since  $|V(D)| = 2n + 3$  and  $A(D) = (n + 1)(2n + 3)$ ,  $D$  is a minimum locally  $n$ -arc-strong oriented graph by Lemma 2.

Now we prove the necessity. Let  $D$  be a minimum locally  $n$ -arc-strong oriented graph. Since we have proved that a diregular tournament of  $2n + 3$  vertices is a minimum locally  $n$ -arc-strong oriented graph, we have  $|V(D)| = 2n + 3$ ,  $|A(D)| = (n + 1)(2n + 3)$ . By Lemma 2,  $\delta(D) \geq n + 1$ . Then we must have  $\text{id } v = \text{od } v = n + 1$  for any vertex  $v$  in  $D$ . Therefore,  $D$  is a diregular tournament of  $2n + 3$  vertices.  $\square$

For the minimum locally  $n$ -strong oriented graphs, we have the following result.

**Theorem 3.** *Every minimum locally  $n$ -strong oriented graph is a diregular tournament of  $2n + 3$  vertices.*

Before giving the proof we need to give a lemma, which also has its own interest.

**Lemma 3.** *Let  $D$  be a tournament. Then  $D$  is locally  $n$ -strong if and only if  $D$  is  $(n + 1)$ -strong.*

*Proof.* The necessity is immediately seen from Theorem B. We only need to show the sufficiency.

Assume there is a tournament  $D$  which is  $(n + 1)$ -strong but not locally  $n$ -strong. Then, there is a vertex  $v$  in  $D$  such that  $\langle N(v) \rangle$  is not  $n$ -strong. Thus, we can find a proper subset  $S$  of  $N(v)$  such that  $|S| \leq n - 1$  and  $\langle N(v) \rangle - S$  is not strong. Let  $S' = S \cup \{v\}$ . Then  $|S'| \leq n$ , and  $D - S' = \langle N(v) \rangle - S$  since  $D$  is a tournament. Thus,  $D - S'$  is not strong, which contradicts the assumption that  $D$  is  $(n + 1)$ -strong.

It completes the proof of Lemma 3.  $\square$

Now we prove Theorem 3 as follows.

*Proof of Theorem 3.* First we claim that for any positive integer  $n$ , there exists a diregular tournament of  $2n + 3$  vertices which is locally  $n$ -strong. For instance, we may consider the right Cayley digraph  $L(Z_{2n+3}, \{1, 2, \dots, n + 1\})$  which

is a diregular tournament of  $2n+3$  vertices. (Recall that for an additive group  $G$  and  $S \subseteq G \setminus \{0\}$ , the right Cayley digraph  $L(G, S)$  is a digraph  $D$  with  $V(D) = G$  and  $A(D) = \{(x, x+y) : y \in S\}$ .) By a result of Y. O. Hamidoune [7, Proposition 5.1],  $L(\mathbb{Z}_{2n+3}, \{1, 2, \dots, n+1\})$  is  $(n+1)$ -strong. Then it is locally  $n$ -strong by Lemma 3. So, our claim is true.

Let  $D$  be a minimum locally  $n$ -strong oriented graph. By the above claim,  $|V(D)| \leq 2n+3$  and  $|A(D)| \leq (n+1)(2n+3)$ . Then by Lemma 2, we must have  $|V(D)| = 2n+3$  and  $|A(D)| = (n+1)(2n+3)$ . Moreover, from Lemma 2,  $\delta(D) \geq n+1$ . Then we must have  $\text{id } v = \text{od } v = n+1$  for every vertex  $v$  in  $D$ . Therefore,  $D$  is a diregular tournament of  $2n+3$  vertices.  $\square$

**Remark 1.** From Lemma 3, it seems natural to pose the following conjecture:

Let  $D$  be a tournament. Then  $D$  is locally  $n$ -arc-strong if and only if  $D$  is  $(n+1)$ -arc-strong.

However, this conjecture is false, which can be seen from Proposition 4 given at the end of this paper.

Note that Theorem 3 only gives a result parallel to the necessity part of Theorem 2. In fact, the converse of Theorem 3 does not hold for  $n \geq 3$ . It can be seen from the following result.

**Theorem 4.** *For any integer  $n \geq 3$ , there exists a diregular tournament of  $2n+3$  vertices which is not locally  $n$ -strong.*

*Proof.* We proceed in two steps.

Step 1. By induction on  $n$ , show that there is a diregular tournament  $D_{2n+3}$  of  $2n+3$  vertices satisfying the following conditions:  $V(D_{2n+3}) = X_n \cup Y_n \cup C$  where  $X_n, Y_n$  and  $C$  are pairwise disjoint,  $|X_n| = |Y_n| = n$  and  $\langle C \rangle$  is a dicycle of length 3; and  $A(D_{2n+3}) \supset (X_n, C) \cup (C, Y_n)$ .

For  $n = 3$ , the desired  $D_9$  can be constructed as follows. Take three pairwise disjoint dicycles of length 3 and denote their vertex sets as  $X_3, Y_3$  and  $C$ . Then add all arcs in  $(X_3, C) \cup (C, Y_3) \cup (Y_3, X_3)$ . It can be easily verified that this digraph is the desired  $D_9$ .

Now, assume that  $D_{2k+3}$  has been constructed for  $k \geq 3$ . We construct a new tournament of two more vertices as follows. First, we add two new vertices  $x$  and  $y$  and add arcs  $(y, x) \cup (\{x\}, C) \cup (C, \{y\})$  so that we have  $\text{od } x - \text{id } x = 2$ ,  $\text{id } y - \text{od } y = 2$  and  $\text{od } v = \text{id } v$  for every  $v \in C$ . Then, we arbitrarily take a subset  $S \subset X_k \cup Y_k$  with  $|S| = k+1$ , and let  $\bar{S} = (X_k \cup Y_k) - S$ . Clearly,  $|S| - |\bar{S}| = 2$ . Then we add arcs  $(S, \{x\}) \cup (\{x\}, \bar{S}) \cup (\{y\}, S) \cup (\bar{S}, \{y\})$ . Let  $X_{k+1} = X_k \cup \{x\}$  and  $Y_{k+1} = Y_k \cup \{y\}$ . Then it is easily seen that the obtained digraph is the desired  $D_{2n+3}$ . This completes the induction.

Step 2. Show that  $D_{2n+3}$  is not locally  $n$ -strong.

Let  $D = D_{2n+3} - X_n$ . Then  $V(D)$  can be decomposed as two disjoint subsets  $C$  and  $Y_n$ . Since  $(Y_n, C) = \emptyset$ ,  $D$  is not strong. Note that  $|X_n| = n$ . Then we see that  $D_{2n+3}$  is not  $(n+1)$ -strong. Hence, it is not locally  $n$ -strong by Theorem B.

It completes the proof of Theorem 4. □

**Remark 2.** The condition  $n \geq 3$  in Theorem 4 is necessary since any diregular tournament of 5 (7, resp.) vertices is easily seen to be locally 1-strong (locally 2-strong, resp.). Therefore, the converse of Theorem 3 only holds for  $n = 1, 2$ .

**Remark 3.** It should be noted that the conclusion in Lemma 3 is not true for general digraphs, i.e., the converses of Theorems A and B are not true, which can be seen from the associated digraphs  $D(K_{n+1, n+k})$  of the complete bipartite graphs  $K_{n+1, n+k}$  (with  $k \geq 1$ ).

It is easy to see the following facts:

- (a)  $G$  is connected iff  $D(G)$  is strong;
- (b)  $G$  is  $n$ -connected iff  $D(G)$  is  $n$ -strong;
- (c)  $G$  is  $n$ -edge-connected iff  $D(G)$  is  $n$ -arc-strong;
- (d)  $G$  is locally  $n$ -connected iff  $D(G)$  is locally  $n$ -strong;
- (e)  $G$  is locally  $n$ -edge-connected iff  $D(G)$  is locally  $n$ -arc-strong (Note:  $G$  is said to be locally  $n$ -edge-connected if the neighborhood of every vertex of  $G$  is  $n$ -edge-connected.)

From these relationships between  $G$  and  $D(G)$ , we can easily see that  $D(K_{n+1, n+k})$  is  $(n+1)$ -strong and  $(n+1)$ -arc-strong but not locally  $n$ -(arc)-strong, since  $K_{n+1, n+k}$  ( $k \geq 1$ ) is  $(n+1)$ -connected and  $(n+1)$ -edge-connected but not locally  $n$ -(edge)-connected.

Finally, let us go back to the conjecture mentioned earlier. It is disproved by the following result:

**Proposition 4.** *For any integer  $n \geq 1$ , there is a tournament which is  $(n+1)$ -arc-strong but not locally  $n$ -arc-strong.*

**Proof.** Let  $D$  be a diregular tournament of  $2n+3$  vertices. Let  $S$  be a subset of  $V(D)$  with  $|S| = n-1$ , and let  $\bar{S} = V(D) - S$ . Then  $|\bar{S}| = n+4$ . Let  $D_1$  be an isomorphic copy of  $D$  under the isomorphism  $\varphi: V(D) \rightarrow V(D_1)$ . Let  $S_1 = \varphi(S)$  and  $\bar{S}_1 = \varphi(\bar{S})$ . Then we extend the digraph  $D \cup D_1$  to a tournament  $H$  by adding arcs between  $V(D)$  and  $V(D_1)$  so that it satisfies the condition  $(V(D), V(D_1)) = \{(x, \varphi(x)) | x \in S\}$ . Then by Lemma 1 of [5] (which says that a digraph  $D$  is  $n$ -arc-strong if and only if  $|(S, \bar{S})_D| \geq n$  for every nonempty proper subset  $S$  of  $V(D)$  (where  $\bar{S} = V(D) - S$ ), we see that  $H$  is not  $n$ -arc-strong since  $|(V(D), V(D_1))| =$

$|S| = n - 1 < n$ . Now we construct the desired tournament  $T$  from  $H$  by adding a new vertex  $x$  and adding all the arcs in  $(\{x\}, S \cup \bar{S}_1) \cup (\bar{S} \cup S_1, \{x\})$ . It is easily seen that  $\delta(T) = n + 2$ .

Note that  $\lfloor \frac{v(T)+2}{4} \rfloor = \lfloor \frac{(2(2n+3)+1)+2}{4} \rfloor = n + 2$ . Then by Lemma 1,  $T$  is  $\delta(T)$ -arc-strong, implying that  $T$  is  $(n + 1)$ -arc-strong. However, since  $N_T(x) = H$ ,  $T$  is not locally  $n$ -arc-strong.  $\square$

### Acknowledgement.

I would like to thank Professor Gary Chartrand for his kindly sending me a copy of his joint paper [4] with Professor Raymond E. Pippert which inspired this work.

### References

- [1] *J.N. Ayoub and I.T. Frish*: On the smallest-branch cuts in directed graphs. IEEE Trans. Circuit Theory *CT-17* (1970), 249–250. (MR 42#1717).
- [2] *J.C. Bermond, N. Homobono and C. Peyrat*: Large fault-tolerant interconnection networks. Graphs and Combinatorics *5* (1989), 107–123.
- [3] *G. Chartrand and L. Lesniak*: Graphs and Digraphs. Second edition, Wadsworth, 1986.
- [4] *G. Chartrand and R.E. Pippert*: Locally connected graphs. Časopis Pěst. Mat. *99* (1974), 158–163. (MR 53#2723).
- [5] *Zhibo Chen*: On locally  $n$ -(arc)-strong digraphs. To appear in Ars Combinatoria.
- [6] *Zhibo Chen*: Local connectedness of digraphs. Graph Theory, Combinatorics, and Applications: Proceedings of the Seventh Quadrennial International Conference on the Theory and Applications of Graphs, Vol. 1 (Y. Alavi and A. Schwenk, eds.). John Wiley & Sons, New York, 1995, pp. 195–200.
- [7] *Y.O. Hamidoune*: On the connectivity of Cayley digraphs. Europ. J. Combinatorics *5* (1984), 309–312.
- [8] *W. Mader*: Connectivity and edge-connectivity in finite graphs. Surveys in Combinatorics: Proc. Seventh British Combinatorial Conference, Cambridge 1979. London, Math. Soc. Lec. Note Ser. 38, 1979, pp. 66–95.
- [9] *L. Nebeský*: Every connected, locally connected graph in upper embeddable. J. Graph Theory *5* (1981), 205–207.
- [10] *L. Nebeský*: On locally quasicomnected graphs and their upper embeddability. Czechoslovak Math. J. *35* (1985), no. 110, 162–166.
- [11] *L. Nebeský*:  $N_2$ -locally connected graphs and their upper embeddability. Czechoslovak Math. J. *41* (1991), no. 116, 731–735.
- [12] *Z. Ryjáček*: On graphs with isomorphic, non-isomorphic and connected  $N_2$ -neighborhoods. Časopis Pěst. Mat. *112* (1987), 66–79.
- [13] *J. Sedláček*: Local properties of graphs. Časopis Pěst. Mat. *106* (1981), 290–298. (In Czech.)
- [14] *D.W. Vanderjagt*: Sufficient conditions for locally connected graphs. Časopis Pěst. Mat. *99* (1974), 400–404.

*Author's address*: Department of Mathematics, Pennsylvania State University, McKeesport, PA 15132, USA.