

Bedřich Pondělíček

Inverse semirings and their lattice of congruences

Czechoslovak Mathematical Journal, Vol. 46 (1996), No. 3, 513–522

Persistent URL: <http://dml.cz/dmlcz/127312>

Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

INVERSE SEMIRINGS AND THEIR LATTICE OF CONGRUENCES

BEDŘICH PONDĚLÍČEK, Praha

(Received August 16, 1994)

To the memory of Otakar Borůvka

Universal Algebra Theory, algebras whose congruences form a modular (distributive, boolean) lattice with respect to inclusion have attracted great attention. For example, in semigroup theory see [1]. The aim of this paper is to describe all inverse semirings having a modular (distributive, boolean) congruence lattice. The notion of an associative inverse semiring can be found in [2]. The congruence lattices of these semirings have also been studied in [3].

1. INTRODUCTION

We shall fix the type $\tau = (t, \text{ar})$ with $t = (+, \cdot, -)$, $\text{ar}(+) = \text{ar}(\cdot) = 2$ and $\text{ar}(-) = 1$. An *inverse semiring* is a τ -algebra $\mathcal{S} = (S, \tau)$ satisfying the axioms

(1.1) $(S, +)$ is a commutative semigroup,

(1.2) $x(y + z) = xy + xz$, $(y + z)x = yx + zx$,

(1.3) $x = (x - x) + x$, $-x = -x + (x - x)$ where $xy = x \cdot y$, $xy + z = (xy) + z$ and $x - y = x + (-y)$,

(1.4) $(x - x) + (y - y) = (x - x)(y - y)$.

Note that we need not use and suppose the associativity of the multiplication.

By $S(\mathcal{S})$ we denote the set of all elements of an inverse semiring \mathcal{S} . We put $E(\mathcal{S}) = \{x \in S(\mathcal{S}), x = x + x\}$. It follows from (1.3) that $x - x \in E(\mathcal{S})$ for every $x \in S(\mathcal{S})$. Let $0: S(\mathcal{S}) \rightarrow E(\mathcal{S})$ be a mapping such that $0x = x - x$ for all $x \in S(\mathcal{S})$. According to (1.1) and (1.3) we have

$$x + 0x = x = 0x + x.$$

In some proofs the following implication will be used:

$$(1.5) \quad x = x + y + x, y = y + x + y \Rightarrow y = -x \quad \text{for every } x, y \in S(\mathcal{S}).$$

Proof. Suppose that $x = x + y + x$ and $y = y + x + y$. Then according to (1.1) and (1.3) we have $y = y + x + y = y + x + (-x) + x + y = (x + y + x) + y + (-x) = x + y + (-x) = x + y + (-x) + x + (-x) = (-x) + (x + y + x) + (-x) = (-x) + x + (-x) = -x$. \square

From (1.1), (1.2) and (1.5) it is easy to show the following:

$$(1.6) \quad -(-x) = x,$$

$$(1.7) \quad -(x + y) = (-x) + (-y),$$

$$(1.8) \quad -(xy) = (-x)y = x(-y).$$

It follows from (1.5) and (1.4) that

$$(1.9) \quad e = -e = 0e = e^2 \quad \text{for every } e \in E(\mathcal{S}) \quad \text{and}$$

$$(1.10) \quad e + f = ef \in E(\mathcal{S}) \quad \text{for every } e, f \in E(\mathcal{S}).$$

This implies that $\mathcal{E}(\mathcal{S}) = (E(\mathcal{S}), \tau)$ is an inverse subsemiring of \mathcal{S} which is a *semilattice* and so for $e, f \in E(\mathcal{S})$ we can put

$$(1.11) \quad e \leq f \quad \text{if and only if } ef = e,$$

$$(1.12) \quad e < f \quad \text{if and only if } e \leq f \quad \text{and } e \neq f,$$

$$(1.13) \quad e \parallel f \quad \text{if and only if } e \neq ef \neq f.$$

According to (1.2), (1.9), (1.8) and (1.10) for every $c \in E(\mathcal{S})$ and every $x \in S(\mathcal{S})$ we have

$$(1.14) \quad xe = (0x)e = e(0x) = ex \in E(\mathcal{S}).$$

Indeed, we have $xe = x(e + c) = x(e - e) = xe - xc = (x - x)e = (0x)e = (0x) = \dots = ex$. It follows from (1.1), (1.6), (1.7), (1.9) and (1.14) that

$$(1.15) \quad 0 \text{ is the homomorphism projection of } \mathcal{S} \text{ onto } \mathcal{E}(\mathcal{S}).$$

2. PROPERTY (M)

Let \mathcal{S} be an inverse semiring. By $(\text{Con}(\mathcal{S}), \wedge, \vee)$ (or briefly $\text{Con}(\mathcal{S})$) we denote the lattice of all congruences on \mathcal{S} with respect to set inclusion. For $x, y \in S(\mathcal{S})$ we denote by $\Theta_{\mathcal{S}}(x, y)$ (or briefly $\Theta(x, y)$) the least congruence on \mathcal{S} containing (x, y) .

Recall that $\text{Ker } 0 = \{(x, y) : x, y \in S(\mathcal{S}) \text{ and } 0x = 0y\} \in \text{Con}(\mathcal{S})$. By $[\text{Ker } 0]$ we denote the principal filter of $\text{Con}(\mathcal{S})$ generated by $\text{Ker } 0$, i.e. $[\text{Ker } 0] = \{X \in \text{Con}(\mathcal{S}) ; \text{Ker } 0 \subseteq X\}$. For every $X \in [\text{Ker } 0]$ we put $\varphi(X) = X \cap (E(\mathcal{S}) \times E(\mathcal{S}))$. It is clear that $\varphi : [\text{Ker } 0] \rightarrow \text{Con}(\mathcal{E}(\mathcal{S}))$.

Lemma 2.1. *The mapping φ is an isomorphism of the lattice $[\text{Ker } 0]$ onto the lattice $\text{Con}(\mathcal{E}(\mathcal{S}))$.*

PROOF. First we shall show that $X_1 \subseteq X_2$ if and only if $\varphi(X_1) \subseteq \varphi(X_2)$ for all $X_1, X_2 \in [\text{Ker } 0]$. It is clear that $X_1 \subseteq X_2$ implies $\varphi(X_1) \subseteq \varphi(X_2)$. Suppose that $\varphi(X_1) \subseteq \varphi(X_2)$. Let $(x, y) \in X_1$. Then $(0x, 0y) \in X_1$ and so $(0x, 0y) \in \varphi(X_1) \subseteq \varphi(X_2) \subseteq X_2$. It is easy to show that $(x, 0x), (0y, y) \in \text{Ker } 0 \subseteq X_2$ and so $(x, y) \in X_2$. We have $X_1 \subseteq X_2$.

Now, we shall prove that φ is a surjective mapping. Let $Y \in \text{Con}(\mathcal{E}(\mathcal{S}))$. Put $X = \{(x, y); x, y \in S(\mathcal{S}) \text{ and } (0x, 0y) \in Y\}$. According to (1.15), we have $X \in \text{Con}(\mathcal{S})$. If $(x, y) \in \text{Ker } 0$, then $0x = 0y$ and so $(x, y) \in X$. Therefore we have $\text{Ker } 0 \subseteq X$ and so $X \in [\text{Ker } 0]$. Finally, we obtain $\varphi(X) = X \cap (E(\mathcal{S}) \times E(\mathcal{S})) \subseteq Y \subseteq \varphi(X)$. Hence we have $\varphi(X) = Y$. \square

Recall that a semilattice \mathcal{E} is a tree if for any pair of elements $e, f \in S(\mathcal{E})$ with $e \parallel f$ there is no element $g \in S(\mathcal{E})$ such that $e \leq g$ and $f \leq g$.

Lemma 2.2. *Let \mathcal{S} be an inverse semiring.*

If $\text{Con}(\mathcal{S})$ is modular, then $\mathcal{E}(\mathcal{S})$ is a tree.

If $\mathcal{E}(\mathcal{S})$ is a tree, then $[\text{Ker } 0]$ is distributive.

The proof follows from Lemma 2.1 and Theorem 4.4 of [1]. \square

Lemma 2.3. *If the lattice $\text{Con}(\mathcal{S})$ is modular, then $a + f = f$ for all elements a, f of an inverse semiring \mathcal{S} , where $f = 0f < 0a$.*

PROOF. Suppose that $\text{Con}(\mathcal{S})$ is modular and $a + f \neq f$, where $f = 0f < 0a = e$. It follows from (1.4) that $a \neq e$. Put $A = \Theta(a + f, f)$, $B = \Theta(e, f)$ and $C = \Theta(a, e)$. We have $(a + f, f) = (a, e) + (f, f) \in C$ and so $A \subseteq C$.

Let $g \in E(\mathcal{S})$ and put $G_g = \{(x, y); x, y \in S(\mathcal{S}), \text{ where } x = y \text{ or } 0x = 0y \leq g\}$. We shall show that $G_g \in \text{Con}(\mathcal{S})$. Evidently G_g is an equivalence on $S(\mathcal{S})$. Suppose that $(x, y) \in G_g$ and $x \neq y$. By (1.15) we have $(-x, -y) \in G_g$. If $z \in S(\mathcal{S})$, then by virtue of (1.4) we obtain $0(x + z) = 0(xz) = (0x)(0z) = (0y)(0z) = 0(yz) = 0(y + z) \leq g$ and $0(xz) = 0(yz) = 0(zx) = 0(zy) \leq g$. Therefore $G_g \in \text{Con}(\mathcal{S})$.

We have $(a + f, f) \in G_f$ and so $A \subseteq G_f$. From $(a, e) \in G_e$ it follows that $C \subseteq G_e$.

Let $D = \{(x, y); x, y \in S(\mathcal{S}) \text{ and } x + f = y + f\}$. We shall prove that $D \in \text{Con}(\mathcal{S})$. It is clear that D is an equivalence on $S(\mathcal{S})$. Assume that $(x, y) \in D$. By (1.7) and (1.9) we have $(-x, -y) \in D$. If $z \in S(\mathcal{S})$, then (1.1) implies $(x + z, y + z) \in D$. It remains to show that $(xz, yz) \in D$ (and dually $(zx, zy) \in D$) for every $z \in S(\mathcal{S})$. We have $x + f = y + f$ and so, by (1.15) and (1.2), we obtain

$$(2.1) \quad 0x + f = 0y + f \quad \text{and} \quad xz + fz = yz + fz.$$

Using (1.14) and (1.10) we have

$$(2.2) \quad xz + f + 0z = yz + f + 0z.$$

From (2.1) and (2.2) it follows that $xz + f = xz + 0(rz) + f = xz + 0x + 0z + f = yz + 0x + 0z + f = yz + 0y + 0z + f = yz + 0(yz) + f = yz + f$. Thus we have $D \in \text{Con}(\mathcal{S})$.

Since $(e, f) \in D$, we have $B \subseteq D$ and so $B \cap C \subseteq D \cap G_e = D \wedge G_e$. We have $(a, a + f) = (a, a) + (e, f) \in B$, $(a + f, f) \in A$ and $(f, e) \in B$. This implies $(a, e) \in (A \vee B) \wedge C = A \vee (B \wedge C) \subseteq G_f \vee (D \wedge G_e)$. Then there exists a finite sequence $a = x_0, x_1, \dots, x_n = e$ of elements from $S(\mathcal{S})$ such that $(x_{i-1}, x_i) \in G_f$ or $(x_{i-1}, x_i) \in D \cap G_e$. We can suppose that the length $n \geq 1$ is minimal.

If $(x_0, x_1) \in G_f$, then $x_0 = x_1$, which is a contradiction. We have $x_0 \neq x_1$ and $(x_0, x_1) \in D \cap G_e$. Then $0x_1 = 0x_0 = e$, $a + f = x_0 + f = x_1 + f$ and so $x_1 \neq e$. Therefore $n \geq 2$ and $(x_1, x_2) \in G_f$. This implies that $x_1 = x_2$, a contradiction.

Consequently, we have $a + f = f$. □

Definition 2.1. We shall say that an inverse semiring \mathcal{S} has property (M) if

$$a + f = f$$

for all $a, f \in S(\mathcal{S})$, where $f = 0f < 0a$.

Lemma 2.4. If an inverse semiring \mathcal{S} has property (M), then for $a, b \in S(\mathcal{S})$ we have

- (i) $a + b = b$ for $0b < 0a$,
- (ii) $ab = ba = 0b$ for $0b < 0a$,
- (iii) $a + b = ab = 0(ab)$ for $0a \parallel 0b$.

Proof. (i) and (ii). If $0b < 0a$, then by (1.3) and Definition 2.1 we obtain $a + b = a + 0b + b = 0b + b = b$. Further we have $ab + b^2 = b^2$ (see (1.2)) and so $ab + 0b^2 = 0b^2$. It is easy to show that $0(ab) = 0b = 0b^2$. Hence we have $ab = 0b$. Analogously we can show that $ba = 0b$.

(iii). Suppose that $0a \parallel 0b$. According to (1.3), (1.4) and Definition 2.1, we have $a + b = a + b + 0(a + b) = a + b + 0(ab) = a + 0(ab) = 0(ab)$. This implies that $a^2 + ab = a0(ab)$ and so, by (1.14), (1.15) and (i), we have $ab = a^2 + ab = (0a)0(ab) = 0(ab)$. □

3. PROPERTY (D)

Let \mathcal{S} be an inverse semiring and let $e \in E(\mathcal{S})$. By \mathcal{S}_e we denote the inverse subsemiring of \mathcal{S} satisfying $S(\mathcal{S}_e) = \{x \in S(\mathcal{S}); 0x = e\}$. By virtue of (1.15) it is easy to show that $E(\mathcal{S}_e) = \{e\}$ and so \mathcal{S}_e is a subring of \mathcal{S} .

Put $H_e = \{(x, y); x, y \in S(\mathcal{S}), \text{ where } x = y \text{ or } 0x = e = 0y\}$.

Lemma 3.1. *Let \mathcal{S} be an inverse semiring having property (M) and let $e \in E(\mathcal{S})$. Then $H_e \in \text{Con}(\mathcal{S})$ and the principal ideal $(H_e]$ of $\text{Con}(\mathcal{S})$ generated by H_e is isomorphic to $\text{Con}(\mathcal{S}_e)$.*

Proof. Suppose that an inverse semiring \mathcal{S} has property (M) and $e \in E(\mathcal{S})$. Let $X \in \text{Con}(\mathcal{S}_e)$ and put $\psi(X) = X \cup \text{id}_{S(\mathcal{S})}$.

First we shall show that $\psi(X) \in \text{Con}(\mathcal{S})$. Evidently, $\psi(X)$ is an equivalence on $S(\mathcal{S})$. Suppose that $(x, y) \in \psi(X)$ and $x \neq y$. Then $(x, y) \in X$ and so $0x = e = 0y$. It is clear that $(-x, -y) \in X \subseteq \psi(X)$, Let $z \in S(\mathcal{S})$. If $0z = e$, then evidently $(x+z, y+z), (xz, yz), (zx, zy) \in X \subseteq \psi(X)$. From Lemma 2.4 we obtain the following implications. If $0z < e$, then $(x+z, y+z) = (z, z)$ and $(xz, yz) = (zx, zy) = (0z, 0z)$ belong to $\psi(X)$. If $e < 0z$, then $(x+z, y+z) = (x, y)$ and $(xz, yz) = (zx, zy) = (e, e)$ belong to $X \subseteq \psi(X)$. If $e \parallel 0z$, then $(x+z, y+z) = (xz, yz) = (zx, zy) = (ez, ez) \in \psi(X)$. Hence $\psi(X) \in \text{Con}(\mathcal{S})$ and so $\psi: \text{Con}(\mathcal{S}_e) \rightarrow \text{Con}(\mathcal{S})$.

It is easy to see that $X \subseteq Y$ if and only if $\psi(X) \subseteq \psi(Y)$ for all $X, Y \in \text{Con}(\mathcal{S}_e)$. Clearly $\psi(S(\mathcal{S}_e) \times S(\mathcal{S}_e)) = H_e$ and so $H_e \in \text{Con}(\mathcal{S})$ and $\psi(\text{Con}(\mathcal{S}_e)) \subseteq (H_e] = \{X \in \text{Con}(\mathcal{S}); X \subseteq H_e\}$. It remains to prove that $\psi(\text{Con}(\mathcal{S}_e)) = (H_e]$. Let $Y \in (H_e]$. Then $Y \subseteq H_e$. Put $X = Y \cap (S(\mathcal{S}_e) \times S(\mathcal{S}_e))$. Clearly $X \in \text{Con}(\mathcal{S}_e)$. Suppose that $(x, y) \in \psi(X)$. Then $(x, y) \in X$ or $x = y$ and so $(x, y) \in Y$. Hence we have $\psi(X) \subseteq Y$. Assume that $(x, y) \in Y$. Then $(x, y) \in H_e$ and this implies $x, y \in S(\mathcal{S}_e)$ or $x = y$. Consequently, we obtain that $(x, y) \in X \cup \text{id}_{S(\mathcal{S})} = \psi(X)$. Thus we have $Y \subseteq \psi(X)$ and so $Y = \psi(X)$. □

Definition 3.1. We shall say that an inverse semiring \mathcal{S} has property (D) if for each $e \in E(\mathcal{S})$ the lattice $\text{Con}(\mathcal{S}_e)$ is distributive.

Lemma 3.2. *If the lattice $\text{Con}(\mathcal{S})$ is distributive, then an inverse semiring \mathcal{S} has property (D).*

The proof follows from Lemma 2.3 and Lemma 3.1. □

4. MODULARITY AND DISTRIBUTIVITY

Lemma 4.1. *Let $X, Y \in \text{Con}(\mathcal{S})$, where \mathcal{S} is an inverse semiring. Then $(e, f) \in X \vee Y$ for $e, f \in E(\mathcal{S})$ if and only if there is a finite sequence $e = x_0, x_1, \dots, x_n = f$ of elements from $E(\mathcal{S})$ such that $(x_{i-1}, x_i) \in X \cup Y$ for $i = 1, 2, \dots, n$.*

Proof. It is well known that $(e, f) \in X \vee Y$ if and only if there is a finite sequence $e = y_0, y_1, \dots, y_n = f$ of elements from $S(\mathcal{S})$ such that $(y_{i-1}, y_i) \in X \cup Y$ for $i = 1, 2, \dots, n$. Suppose that $e, f \in E(\mathcal{S})$ and put $x_i = 0y_i$ for $i = 0, 1, \dots, n$. Then we have $(x_{i-1}, x_i) \in X \cup Y$ for $i = 1, 2, \dots, n$ and $x_0 = e, x_n = f$ and $x_i \in E(\mathcal{S})$. □

Lemma 4.2. *Let $X \in \text{Con}(\mathcal{S})$, where S is an inverse semiring. If for $e, f \in E(\mathcal{S})$ we have $(e, f) \in X \vee \text{Ker } 0$, then $(e, f) \in X$.*

Proof. According to Lemma 4.1, there is a finite sequence $e = x_0, x_1, \dots, x_n = f$ of elements from $E(\mathcal{S})$ such that $(x_{i-1}, x_i) \in X \cup \text{Ker } 0$ for $i = 1, 2, \dots, n$. If $(x_{i-1}, x_i) \in \text{Ker } 0$, then $x_{i-1} = 0x_{i-1} = 0x_i = x_i$ and so $(x_{i-1}, x_i) \in X$. Therefore we have $(e, f) \in X$. □

Lemma 4.3. *Let $A, B, C \in \text{Con}(\mathcal{S})$, where \mathcal{S} is an inverse semiring in which $\mathcal{E}(\mathcal{S})$ is a tree. If for $e, f \in E(\mathcal{S})$ we have $(e, f) \in (A \vee B) \wedge C$, then $(e, f) \in (A \wedge C) \vee (B \wedge C)$.*

Proof. Suppose that $(e, f) \in (A \vee B) \wedge C$, where $e, f \in E(\mathcal{S})$. Put $A' = A \vee \text{Ker } 0, B' = B \vee \text{Ker } 0$ and $C' = C \vee \text{Ker } 0$. Clearly we have $(e, f) \in (A' \vee B') \wedge C'$. It follows from Lemma 2.2 that $(e, f) \in (A' \wedge C') \vee (B' \wedge C')$. By Lemma 4.1 there is a finite sequence $e = x_0, x_1, \dots, x_n = f$ of elements from $E(\mathcal{S})$ such that $(x_{i-1}, x_i) \in (A' \wedge C') \cup (B' \wedge C')$ for $i = 1, 2, \dots, n$. According to Lemma 4.2, we have $(x_{i-1}, x_i) \in (A \wedge C) \cup (B \wedge C)$. Consequently, $(e, f) \in (A \wedge C) \vee (B \wedge C)$. □

Lemma 4.4. *Let $X, Y \in \text{Con}(\mathcal{S})$, where \mathcal{S} is an inverse semiring. If $(x, z) \in X, (z, y) \in Y$ and $0x = 0z = 0y$, then there exists w such that $(x, w) \in Y, (w, y) \in X$ and $0w = 0z$.*

Proof. Put $w = x - z + y$. It follows from (1.4) that $0w = 0z$ and $(x, w) = (x, x) - (z, z) + (z, y) \in Y, (w, z) = (x, z) - (z, z) + (y, y) \in X$. □

Lemma 4.5. *Let $X \in \text{Con}(\mathcal{S})$, where \mathcal{S} is an inverse semiring having property (M). Let $(x, y) \in X$. If $0x < 0y$ or $0x \parallel 0y$, then $(u, v) \in X$ for all $u, v \in S(\mathcal{S})$ with $0u = 0y = 0v$.*

Proof. Suppose that $0x < 0y$. Then according to Lemma 2.4, we have $(u, x) = (u - y, u - y) + (y, x) \in X$ and $(x, v) = (v - y, v - y) + (x, y) \in X$. Therefore $(u, v) \in X$.

Assume that $0x \parallel 0y$. Then we have $0(xy) < 0y$. Using (1.3), (1.4) and (1.14) we obtain $(0(xy), y) = (0y, 0y) + (x, y) \in X$. The rest of the proof follows from its first part. \square

Lemma 4.6. *Let $A, B, C \in \text{Con}(\mathcal{S})$, where \mathcal{S} is an inverse semiring having property (M), and $\mathcal{E}(\mathcal{S})$ is a tree.*

- (i) *If $A \subseteq C$, then $(A \vee B) \wedge C \subseteq A \vee (B \wedge C)$.*
- (ii) *If \mathcal{S} has property (D), then $(A \vee B) \wedge C \subseteq (A \vee C) \vee (B \wedge C)$.*

Proof. Let $(u, v) \in (A \vee B) \wedge C$. Then there exists a finite sequence $u = x_0, x_1, \dots, x_n = v$ of elements from $S(\mathcal{S})$ such that $(x_{i-1}, x_i) \in A \cup B$ for $i = 1, 2, \dots, n$. Further, we have $(u, v) \in C$. Put $e = 0u$ and $f = 0v$. Clearly $(e, f) \in (A \vee B) \wedge C$. We have the following possibilities:

Case 1. $e = f$. Put $y_i = x_i + e$. It follows from (1.4) that $0y_i \leq e, u = y_0, v = y_n$ and $(y_{i-1}, y_i) \in A \cup B$ for $i = 1, 2, \dots, n$.

Subcase 1a. $0y_i = e$ for all $i = 1, 2, \dots, n$. It follows from Lemma 4.4 that there is an element w of $S(\mathcal{S})$ such that $(u, w) \in A, (w, v) \in B$ and $0w = e$.

If $A \subseteq C$ then $(w, v) \in C$ and so $(u, v) \in A \vee (B \wedge C)$.

If \mathcal{S} has property (D), then according to Lemma 3.1, the lattice $(H_e]$ is distributive. Clearly $(u, w), (w, v), (u, v) \in H_e$ and so $(u, w) \in A', (w, v) \in B', (u, v) \in C'$, where $A' = A \wedge H_e, B' = B \wedge H_e$ and $C' = C \wedge H_e$. It is clear that $A', B', C' \in (H_e]$ and so we have $(u, v) \in (A' \vee B') \wedge C' = (A' \wedge C') \vee (B' \wedge C') \subseteq (A \wedge C) \vee (B \wedge C)$.

Subcase 1b. $0y_i < e$ for some i . It follows from Lemma 4.5 that $(u, v) \in A$ or $(u, v) \in B$. Thus we have $(u, v) \in (A \wedge C) \vee (B \wedge C)$.

Case 2. $f < e$. It follows from Lemma 4.3 that $(f, e) \in (A \wedge C) \vee (B \wedge C)$. According to Lemma 4.5, we have $(u, e) \in (A \wedge C) \vee (B \wedge C)$ and so $(u, f) \in (A \wedge C) \vee (B \wedge C) \subseteq (A \vee B) \wedge C$. This implies that $(f, v) \in (A \vee B) \wedge C$. Using Case 1 we can continue our proof.

If $A \subseteq C$, then $(f, v) \in A \vee (B \wedge C)$ and so $(u, v) \in A \vee (B \wedge C)$. If \mathcal{S} has property (D), then $(f, v) \in (A \wedge C) \vee (B \wedge C)$ and so $(u, v) \in (A \wedge C) \vee (B \wedge C)$.

Case 3. $e < f$. This is dual to Case 2.

Case 4. $e \parallel f$. According to Lemma 4.3, we have $(e, f) \in (A \wedge C) \vee (B \wedge C)$. It follows from Lemma 4.5 that $(u, e), (f, v) \in (A \wedge C) \vee (B \wedge C)$. Therefore $(u, v) \in (A \wedge C) \vee (B \wedge C)$. \square

Theorem 4.1. *Let \mathcal{S} be an inverse semiring. Then*

- (i) $\text{Con}(\mathcal{S})$ is modular if and only if $\mathcal{E}(\mathcal{S})$ is a tree and \mathcal{S} has property (M);
- (ii) $\text{Con}(\mathcal{S})$ is distributive if and only if $\mathcal{E}(\mathcal{S})$ is a tree and \mathcal{S} has properties (M) and (D).

The proof follows from Lemmas 2.2, 2.3, 3.2 and 4.6. □

5. PROPERTY (B)

Definition 5.1. We shall say that an inverse semiring has property (B) if $\text{card } S(\mathcal{S}_e) > 1$ implies that e is the zero of $\mathcal{E}(\mathcal{S})$ and $\text{Con}(\mathcal{S}_e)$ is boolean.

Lemma 5.1. *If the lattice $\text{Con}(\mathcal{S})$ is boolean, then the inverse semiring \mathcal{S} has property (B).*

Proof. Assume that $\text{Con}(\mathcal{S})$ is boolean and $e \in E(\mathcal{S})$. It follows from Lemma 2.3 that \mathcal{S} has property (M). According to Lemma 3.1, the lattice $\text{Con}(\mathcal{S}_e)$ is isomorphic to the principal ideal $(H_e]$ of $\text{Con}(\mathcal{S})$. It is well known that $(H_e]$ is boolean and so $\text{Con}(\mathcal{S}_e)$ is boolean. Suppose by way of contradiction that $\text{card } S(\mathcal{S}_e) > 1$ and e is no zero of $\mathcal{E}(\mathcal{S})$. Then there exists some f of $E(\mathcal{S})$ such that $f < e$. Since $\text{Con}(\mathcal{S})$ is boolean, there is $Y \in \text{Con}(\mathcal{S})$ such that $H_e \wedge Y = \text{id}_{S(\mathcal{S})}$ and $H_e \vee Y = S(\mathcal{S}) \times S(\mathcal{S})$. We have $(e, f) \in H_e \vee Y$ and so, by Lemma 4.1, there is a finite sequence $e = x_0, x_1, \dots, x_n = f$ of elements from $E(\mathcal{S})$ such that $(x_{i-1}, x_i) \in H_e \cup Y$ for $i = 1, 2, \dots, n$. If $(x_{i-1}, x_i) \in H_e$, then $x_{i-1} = x_i$ and so $(x_{i-1}, x_i) \in Y$. Thus we have $(e, f) \in Y$. Let $a \in S(\mathcal{S}_e)$. By (M), we obtain $(a, f) = (e, f) + (a, a) \in Y$. Hence $(a, e) \in Y \cap H_e = \text{id}_{S(\mathcal{S})}$. This implies that $a = e$ and so $\text{card } S(\mathcal{S}_e) = 1$, a contradiction.

Consequently, \mathcal{S} has property (B). □

Recall that a tree \mathcal{E} is said to be *locally finite* if every interval of \mathcal{E} is a finite chain.

Lemma 5.2. *Let E be a semilattice. Then $\text{Con}(E)$ is boolean if and only if E is a locally finite tree.*

Proof. See Theorem 4.5 of [1]. □

Theorem 5.1. *Let \mathcal{S} be an inverse semiring. Then the lattice $\text{Con}(\mathcal{S})$ is boolean if and only if $\mathcal{E}(\mathcal{S})$ is a locally finite tree and \mathcal{S} has property (B).*

Proof. 1. Suppose that $\text{Con}(\mathcal{S})$ is boolean. According to Lemma 5.1, \mathcal{S} has property (B). It follows from Lemma 2.1 that the lattice $\text{Con}(\mathcal{E}(\mathcal{S}))$ is isomorphic

to the principal filter $[\text{Ker } 0]$ of $\text{Con}(\mathcal{S})$, which is boolean. Thus $\text{Con}(\mathcal{E}(\mathcal{S}))$ is boolean. It follows from Lemma 5.2 that $\mathcal{E}(\mathcal{S})$ is a locally finite tree.

2. Now we assume that $\mathcal{E}(\mathcal{S})$ is a locally finite tree and \mathcal{S} has property (B). Lemma 2.1 and Lemma 5.2 imply that the principal filter $[\text{Ker } 0]$ of $\text{Con}(\mathcal{S})$ is boolean.

If $E(\mathcal{S}) = S(\mathcal{S})$, then $\text{Ker } 0 = \text{id}_{S(\mathcal{S})}$ and so $[\text{Ker } 0] = \text{Con}(\mathcal{S})$, which is boolean.

Assume that $E(\mathcal{S}) \neq S(\mathcal{S})$. Property (B) implies that $\mathcal{E}(\mathcal{S})$ has the zero e , $\text{Con}(\mathcal{S}_e)$ is boolean and according to (1.10), \mathcal{S} has property (M). Lemma 3.1 implies that the principal ideal $(\text{Ker } 0]$ of $\text{Con}(\mathcal{S})$ is boolean, because $H_e = \text{Ker } 0$. Hence $(\text{Ker } 0] \times [\text{Ker } 0]$ is a boolean lattice.

Now, we shall prove that $\text{Ker } 0$ has a complement in the lattice $\text{Con}(\mathcal{S})$. Put $P = \{(x, y); x = y \in S(\mathcal{S}) \text{ or } x, y \in E(\mathcal{S})\}$. We shall show that $P \in \text{Con}(\mathcal{S})$. Clearly P is an equivalence on $S(\mathcal{S})$. Assume that $(x, y) \in P, x \neq y$, and $z \in S(\mathcal{S})$. Then $x, y \in E(\mathcal{S})$ and so, by (1.14), we have $xz, yz, zx, zy \in E(\mathcal{S})$. This means that $(xz, yz), (zx, zy) \in P$. If $z \in E(\mathcal{S})$, then it follows from (1.15) that $x + z, y + z \in E(\mathcal{S})$. Thus we have $(x + z, y + z) \in P$. Suppose that $z \notin E(\mathcal{S})$. Then $z \in S(\mathcal{S}_e)$ and so, by (1.3), (1.4), we have $x + z = x + z + e = z = y + z + e = y + z$. This implies that $(x + z, y + z) \in P$. Consequently, $P \in \text{Con}(\mathcal{S})$. It is easy to show that $P \wedge \text{Ker } 0 = \text{id}_{S(\mathcal{S})}$ and $P \vee \text{Ker } 0 = S(\mathcal{S}) \times S(\mathcal{S})$.

Finally, if \mathcal{S} has property (B), then it has properties (M) and (D) and so according to Theorem 4.1, $\text{Con}(\mathcal{S})$ is distributive. It follows from Theorem 6 (Section 7) of [4] that $\text{Con}(\mathcal{S})$ is isomorphic to the boolean lattice $(\text{Ker } 0] \times [\text{Ker } 0]$. Hence $\text{Con}(\mathcal{S})$ is boolean. □

6. INVERSE Δ -SEMIRINGS

Following the semigroup theory, an inverse semiring \mathcal{S} is called an inverse Δ -semiring if the lattice $\text{Con}(\mathcal{S})$ is a chain.

Lemma 6.1. *If \mathcal{S} is an inverse Δ -semiring, then $\text{card } E(\mathcal{S}) \leq 2$.*

Proof. It follows from Lemma 2.1 that $\text{Con}(E(\mathcal{S}))$ is a chain and so, by Lemma 3 of [5], we obtain $\text{card } E(\mathcal{S}) \leq 2$. □

Lemma 6.2. *Let \mathcal{S} be an inverse Δ -semiring. If $E(\mathcal{S}) = \{e, f\}, e < f$, then $\text{card } S(\mathcal{S}_e) = 1$.*

Proof. Put $Q = \{(x, y); x, y \in S(\mathcal{S}) \text{ and } x + e = y + e\}$. It is easy to show that $Q \in \text{Con}(\mathcal{S})$. Since $\text{Con}(\mathcal{S})$ is a chain we have the following two possibilities:

Case 1. $Q \subseteq \text{Ker } 0$. We have $(e, f) \in Q$ and so $(c, f) \in \text{Ker } 0$. This means that $e = f$, a contradiction.

Case 2. $\text{Ker } 0 \subseteq Q$. For $x, y \in S(\mathcal{S}_e)$ we have $(x, y) \in \text{Ker } 0$ and so $(x, y) \in Q$. Thus $x = x + e = y + e = y$. Consequently, $\text{card } S(\mathcal{S}_e) = 1$. \square

Let \mathcal{R} be a ring. By \mathcal{R}^0 we denote the τ -algebra, where $S(\mathcal{R}^0) = S(\mathcal{R}) \cup \{h\}$, $h \notin S(\mathcal{R})$, and

$$-x = \begin{cases} x & \text{for } x = h, \\ -x & \text{for } x \in S(\mathcal{R}). \end{cases}$$

The addition and the multiplication on $S(\mathcal{R}^0)$ are defined as follows:

If $x, y \in S(\mathcal{R})$, then $x + y$ (xy , respectively) is the same as in \mathcal{R} .

If $x \in S(\mathcal{R}^0)$, then $x + h = h + x = xh = hx = h$.

It is easy to show that \mathcal{R}^0 is an inverse semiring with $\text{card } E(\mathcal{R}^0) = 2$.

For every $X \in \text{Con}(\mathcal{R})$ we put $X^0 = X \cup \{(h, h)\}$. Clearly we have $X^0 \in \text{Con}(\mathcal{R}^0)$.

Lemma 6.3. *Let \mathcal{R} be a ring. Then $\text{Con}(\mathcal{R}^0) = \{X^0; X \in \text{Con}(\mathcal{R})\} \cup \{S(\mathcal{R}^0) \times S(\mathcal{R}^0)\}$.*

Proof. The former statement “ \supseteq ” is obvious. To prove the latter statement “ \subseteq ”, let $Y \in \text{Con}(\mathcal{R}^0)$ and $(h, z) \in Y$ for some $z \in S(\mathcal{R})$. Then for arbitrary $x, y \in S(\mathcal{R}^0)$ we have $(h, x) = (h, z) - (h, z) + (x, x) \in Y$ and similarly $(h, y) \in Y$. Therefore $(x, y) \in Y$, which means that $Y = S(\mathcal{R}^0) \times S(\mathcal{R}^0)$. \square

Theorem 6.1. *Let \mathcal{S} be an inverse semiring. Then $\text{Con}(\mathcal{S})$ is a chain if and only if \mathcal{S} is isomorphic to either \mathcal{R} or \mathcal{R}^0 , where \mathcal{R} is a ring whose $\text{Con}(\mathcal{R})$ is a chain.*

The proof follows from Lemmas 6.1, 6.2 and 6.3. \square

References

- [1] *H. Mitsch:* Semigroups and their lattice of congruences. *Semigroup Forum* 26 (1983), 1-63.
- [2] *V. N. Salij:* To the theory of inverse semirings (Russian). *Izv. vuzov. Matem.* (1969), 52-60.
- [3] *B. B. Kovalenko:* On the theory of generalized modules (Russian). *Izdat. Saratov. Univ. Saratov, Studies in algebra* (1977), no. No. 5, 30-43.
- [4] *G. Grätzer:* Lattice theory, first concepts and distributive lattices. San Francisco, 1971.
- [5] *T. Tamura:* Commutative semigroups whose lattice of congruences is a chain. *Bull. Soc. Math. France* 97 (1969), 369-380.

Author's address: 166 27 Praha 6, Technická 2, Czech Republic (Fakulta elektrotechnická ČVUT).