

Wiesław Aleksander Dudek; Xiaohong Zhang
On ideals and congruences in BCC-algebras

Czechoslovak Mathematical Journal, Vol. 48 (1998), No. 1, 21–29

Persistent URL: <http://dml.cz/dmlcz/127395>

Terms of use:

© Institute of Mathematics AS CR, 1998

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON IDEALS AND CONGRUENCES IN BCC-ALGEBRAS

WIESŁAW A. DUDEK and XIAOHONG ZHANG

(Received January 18, 1995)

Abstract. We introduce a new concept of ideals in BCC-algebras and describe connections between such ideals and congruences.

MSC 2000: Primary: 06F35, 03G25, 08A30

Keywords: BCC-algebra, BCK-algebra, ideal, congruence

1. INTRODUCTION

By an algebra $\mathbf{G} = (G, \cdot, 0)$ we mean a non-empty set G together with a binary multiplication and a distinguished element 0 . In the sequel a multiplication will be denoted by juxtaposition. Dots we use only to avoid repetitions of brackets. For example, the formula $((xy)(zy))(xz) = 0$ will be written as $(xy \cdot zy) \cdot xz = 0$.

Definition. An algebra $(G, \cdot, 0)$ is called a *BCC-algebra* if it satisfies the following axioms:

- (1) $(xy \cdot zy) \cdot xz = 0,$
- (2) $xx = 0,$
- (3) $0x = 0,$
- (4) $x0 = x,$
- (5) $xy = yx = 0$ implies $x = y.$

The above definition is a dual form of the ordinary definition (cf. [1], [6], [7]). In our convention *any BCK-algebra is a BCC-algebra*, but *there are BCC-algebras which are not BCK-algebras* (cf. [2]). Such BCC-algebras are called *proper*. Some methods of construction of BCC-algebras from BCK-algebras are given in [3]. Note

that (cf. [2]) a BCC-algebra is a BCK-algebra iff it satisfies

$$(6) \quad xy \cdot z = xz \cdot y.$$

2. IDEALS

As is well-known (cf. for example [4], [5]) a non-empty subset A of a BCK-algebra $(G, \cdot, 0)$ is called an ideal if

- (i) $0 \in A$,
- (ii) $xy \in A$ and $y \in A$ imply $x \in A$.

In the sequel this ideal will be called a BCK-ideal and will be considered also in BCC-algebras.

If A is a BCK-ideal of a BCK-algebra \mathbf{G} then the relation \sim defined on \mathbf{G} by

$$(7) \quad x \sim y \text{ iff } xy, yx \in A$$

is a congruence (cf. [4]). We say that this relation is defined by the ideal A .

This result is not true for BCC-algebras.

Example 2.1. Let $G = \{0, 1, 2, 3, 4\}$ and let the multiplication be defined by the table

\cdot	0	1	2	3	4
0	0	0	0	0	0
1	1	0	1	0	0
2	2	2	0	0	0
3	3	3	1	0	0
4	4	3	4	3	0

First we prove that this algebra is a BCC-algebra. It is clear that such algebra satisfies (2), (3), (4) and (5). We prove (1). If x, y, z are not different, then obviously (1) holds. For different x, y, z we verify only the case when one of elements x, y, z is equal to 4, because $S = \{0, 1, 2, 3\}$ is a BCC-algebra (cf. Table 14 in [2]). Since $xy \in S$, $4y \in \{3, 4\}$ and $u3 = u4 = 0$ for all $x, y, u \in S$, then (1) holds for $z = 4$. For $y = 4$ it holds, too. For $x = 4$ the left hand side of (1) has the form $(4y \cdot zy) \cdot 4z$, which for $y = 1$ and $y = 3$ is equal to 0 since $4y \cdot zy = 3 \cdot zy \in S$ and $u3 = u4 = 0$ for $u \in S$. The case $y = 0$ is obvious. If $y = 2$ then $(42 \cdot z2) \cdot 4z = (4 \cdot z2) \cdot 4z$, which for $z = 0$ trivially gives 0. For $z \in \{1, 3\}$ we obtain $(4 \cdot z2) \cdot 4z = 41 \cdot 3 = 0$. This completes the proof that G is a BCC-algebra.

It is not difficult to verify that $A = \{0, 1\}$ is a BCK-ideal of this BCC-algebra, but the relation \sim defined by this ideal is not a congruence. Indeed, $4 \sim 4$, $2 \sim 3$ but *not* $(42 \sim 43)$ since $42 \cdot 43 = 3 \notin A$.

In connection with this fact we introduce a new concept of ideals.

Definition. A non-empty subset A of a BCC-algebra \mathbf{G} is called a *BCC-ideal*, if

$$(8) \quad 0 \in A,$$

$$(9) \quad xy \cdot z \in A \text{ and } y \in A \text{ imply } xz \in A.$$

Lemma 2.2. *In a BCC-algebra any BCC-ideal is a BCK-ideal.*

Indeed, putting $z = 0$ in (9) we obtain (ii).

On the other hand, using (6) we have

Lemma 2.3. *In a BCK-algebra any BCK-ideal is a BCC-ideal.*

Lemma 2.4. *In a BCC-algebra any BCK-ideal is a BCC-subalgebra.*

Proof. Let A be a BCK-ideal. Then $0 \in A$ and $xy \cdot x = 0$ for all $x, y \in G$ (cf. [2]). Thus for $x, y \in A$ we have $xy \cdot x \in A$, which implies $xy \in A$. \square

Corollary 2.5. *Any BCC-ideal of a BCC-algebra is a BCC-subalgebra.*

The following example shows that a BCC-ideal is not a BCK-subalgebra, in general.

Example 2.6. Let $G = \{0, 1, 2, 3, 4, 5\}$ and let the multiplication be defined by the table

·	0	1	2	3	4	5
0	0	0	0	0	0	0
1	1	0	0	0	0	1
2	2	2	0	0	1	1
3	3	2	1	0	1	1
4	4	4	4	4	0	1
5	5	5	5	5	5	0

Since $S = \{0, 1, 2, 3, 4\}$ is a BCC-algebra (cf. Table 2 in [1]), then \mathbf{G} is a BCC-algebra by Proposition 4 in [2] (cf. also Construction 3 in [3]). It is easy to see that S is a BCC-ideal of \mathbf{G} . It is not a BCK-algebra since $21 \cdot 4 \neq 24 \cdot 1$.

On the other hand, in Example 2.1 $S = \{0, 1, 2, 3\}$ is a BCC-subalgebra which is not a BCK-ideal, because $43 = 3 \in S$ but $4 \notin S$. Similarly, $A = \{0, 1, 2\}$ is a BCK-subalgebra which is not a BCK-ideal since $32 \in A$ but $3 \notin A$. Thus in BCC-algebras BCC-ideals, BCK-ideals and BCK-subalgebras are independent concepts.

Proposition 2.7. *Let \mathbf{G} be a BCC-algebra. Then a BCC-subalgebra \mathbf{A} of \mathbf{G} is a BCC-ideal iff $x \in A, yz \notin A$ imply $yx \cdot z \notin A$.*

Proof. If a BCC-subalgebra A is a BCC-ideal, then $x \in A, yz \notin A$ imply $yx \cdot z \notin A$. If not, then $yx \cdot z \in A, x \in A$ imply $yz \in A$, which is a contradiction.

Conversely, let A be a BCC-subalgebra in which $x \in A, yz \notin A$ imply $yx \cdot z \notin A$. Then obviously $0 \in A$. Moreover, $x \in A, yx \cdot z \in A$ gives $yz \in A$, because for $yz \notin A$ we have (by assumption) $yx \cdot z \notin A$. Hence A is a BCC-ideal. \square

Putting $z = 0$ in the above Proposition we obtain

Proposition 2.8. *Let \mathbf{G} be a BCC-algebra. Then a BCC-subalgebra \mathbf{A} of \mathbf{G} is a BCK-ideal iff $x \in A, y \notin A$ imply $yx \notin A$.*

Proposition 2.9. *Let A be a BCK-ideal of a BCC-algebra \mathbf{G} . If B is a BCK-ideal of A , then it is a BCK-ideal of \mathbf{G} .*

Proof. Since B is a BCK-ideal of A , then $0 \in B$. Let $y, xy \in B$ for some $x \in G$. Then $y, xy \in A$ and $x \in A$ because $B \subset A$ and A is a BCK-ideal of \mathbf{G} . Thus $x \in A$ and $xy, y \in B$ imply $x \in B$. This completes the proof. \square

Corollary 2.10. *Let A be a BCC-ideal of a BCC-algebra \mathbf{G} . If B is a BCK-ideal of A , then B is a BCK-ideal of \mathbf{G} .*

Remark 2.11. If a BCC-ideal A is a BCK-subalgebra of \mathbf{G} , then any of its sub-BCK-ideals is a BCC-ideal, but in general it is not a BCC-ideal of \mathbf{G} .

Remark 2.12. On any BCC-algebra $(G, \cdot, 0)$ one can define (cf. [2]) the so-called *natural order* by putting

$$x \leq y \text{ iff } xy = 0.$$

As in the case of BCK-algebras, this order is partial and 0 is its smallest element. Thus any BCC-algebra may be viewed as a groupoid $(G, \cdot, 0)$ with the natural order satisfying conditions $xy \cdot zy \leq xz, 0 \leq x, x0 = x, x \leq y \leq x$ imply $x = y$ (cf. Theorem 2 from [2]). But, in general, BCC-algebras with the same partial order are not isomorphic as groupoids (cf. [2]).

Remark 2.13. The above ideals are ideals in the sense of ordered structures. Indeed, if A is a BCC-ideal (or a BCK-ideal), then $y \in A$ and $x \leq y$ imply $x \in A$.

3. CONGRUENCES

In this section we describe congruences on BCC-algebras. We start with the following

Theorem 3.1. *If A is a BCC-ideal of a BCC-algebra \mathbf{G} , then the relation \sim defined by (7) is a congruence on \mathbf{G} .*

Proof. It is clear that this relation is reflexive and symmetric. It is also transitive, because $x \sim y$ and $y \sim z$ imply $xy, yx, yz, zy \in A$ and $(xz \cdot yz) \cdot xy = 0 \in A$, which by Lemma 2.2 gives $xz \in A$. Similarly $(zx \cdot yx) \cdot zy = 0 \in A$ gives $zx \in A$. Thus $x \sim z$ and \sim is an equivalence relation.

If $x \sim u$ and $y \sim v$, then $(xy \cdot uy) \cdot xu = 0 \in A$ and $xu \in A$, which by Lemma 2.2 gives $xy \cdot uy \in A$. Similarly $uy \cdot xy \in A$. Hence $xy \sim uy$. On the other hand $(uy \cdot vy) \cdot uv = 0 \in A$ and $vy \in A$ imply $uy \cdot uv \in A$. In the same manner from $(uv \cdot yv) \cdot uy = 0 \in A$ and $yv \in A$ we obtain $uv \cdot uy \in A$. Thus $uy \sim uv$. Since \sim is transitive, then $xy \sim uv$, which proves that \sim is a congruence. \square

Lemma 3.2. *If \sim is a congruence on a BCC-algebra \mathbf{G} , then*

$$C_0 = \{x \in G: x \sim 0\}$$

is a BCC-ideal.

Proof. Obviously $0 \in C_0 = \{x \in G: x \sim 0\}$. If $xy \cdot z, y \in C_0$, then $xy \cdot z \sim 0$ and $y \sim 0$. But $x \sim x$ and $z \sim z$ imply $xy \cdot z \sim x0 \cdot z = xz$. Thus $xz \sim 0$, which completes the proof. \square

Since $C_0 = A$ for any congruence defined by (7), then as a consequence of the above results we obtain

Corollary 3.3. *Any BCC-ideal is determined by some congruence.*

Corollary 3.4. *The lattice of all congruences of a BCC-algebra is complete. The least congruence is defined by the BCC-ideal $\{0\}$, the greatest by $A = G$.*

Let \sim be a congruence relation on \mathbf{G} and let $C_x = \{y \in G: y \sim x\}$. Then the family $\{C_x: x \in G\}$ gives a partition of G which is denoted by G/\sim . For $x, y \in G$ we define $C_x * C_y = C_{xy}$. Since \sim has the substitution property, the operation $*$ is well-defined. As is easily seen, $(G/\sim, *, C_0)$ satisfies all axioms of a BCC-algebra except (5). This axiom is not satisfied also in the case of BCK-algebras (cf. [5] and

[8]). If (5) holds for all classes $C_x \in G/\sim$, i.e. if $(G/\sim, *, C_0)$ is a BCC-algebra, then the congruence \sim is called *regular*.

For a congruence defined by (7) we put $G/\sim = G/A$ and $C_0 = A$.

Theorem 3.5. *A congruence is regular iff it is defined by some BCC-ideal.*

Proof. Let ϱ_A be a congruence defined by a BCC-ideal A . Then $A_0 = A$ and $A_x * A_y = A_0 = A_y * A_x$ imply $xy, yx \in A$, which shows that $x \sim y$ and $A_x = A_y$. Hence a congruence defined by a BCC-ideal is regular.

Now let ϱ be an arbitrary regular congruence. If $x\varrho y$, then $xy\varrho 0$ and $yx\varrho 0$ since ϱ is reflexive and has the substitution property. Therefore $C_{xy} = C_0 = C_{yx}$, $xy, yx \in C_0$ and $A = C_0$ is a BCC-ideal (by Lemma 3.2). Hence $\varrho \leq \varrho_A$.

Conversely, if $x\varrho_A y$, then $xy, yx \in A = C_0$ and $C_x * C_y = C_0 = C_y * C_x$, which implies $C_x = C_y$ because ϱ_A is regular. Thus $x\varrho y$ and $\varrho_A \leq \varrho$. Hence $\varrho = \varrho_A$. The proof is complete. \square

Corollary 3.6. *All congruences of a finite BCC-algebra are regular.*

If \mathbf{G}/\mathbf{A} is a BCC-algebra, then the canonical mapping $f: \mathbf{G} \mapsto \mathbf{G}/\mathbf{A}$ defined by $f(x) = A_x$ is an epimorphism. Since the kernel $\ker f = f^{-1}(0)$ of any BCC-homomorphism is a BCC-ideal, then in the same manner as in [5] we can prove the following results:

Theorem 3.7. *If f is an epimorphism from a BCC-algebra \mathbf{G} onto a BCC-algebra \mathbf{H} , then the quotient BCC-algebra $\mathbf{G}/\ker(f)$ is isomorphic to \mathbf{H} .*

Theorem 3.8. *Let $\mathbf{X}, \mathbf{Y}, \mathbf{Z}$ be BCC-algebras, let $h: \mathbf{X} \mapsto \mathbf{Y}$ be an epimorphism, and let $g: \mathbf{X} \mapsto \mathbf{Z}$ be a homomorphism. If $\ker(h) \subset \ker(g)$, then there exists a unique homomorphism $f: \mathbf{X} \mapsto \mathbf{Z}$ such that $f \circ h = g$.*

Corollary 3.9. *Let ϱ be a regular congruence on a BCC-algebra \mathbf{X} defined by a BCC-ideal A , and let h be a canonical mapping from \mathbf{X} onto $\mathbf{Y} = \mathbf{X}/\mathbf{A}$. If $A_0 \subset \ker(g)$, then there exists a unique homomorphism $f: \mathbf{X} \mapsto \mathbf{Z}$ such that $f \circ h = g$.*

Corollary 3.10. *Let h be a homomorphism from a BCC-algebra \mathbf{G} onto a BCC-algebra \mathbf{H} . Then the inverse image of a BCC-ideal, a BCC-subalgebra and a BCK-subalgebra of \mathbf{H} is a BCC-ideal, a BCC-subalgebra and a BCK-subalgebra of \mathbf{G} , respectively.*

The theory of universal algebras yields immediately

Theorem 3.11. *The composition $\varrho \circ \sigma$ of two congruences on a BCC-algebra \mathbf{G} is a congruence on \mathbf{G} iff these congruences commute, i.e. iff $\varrho \circ \sigma = \sigma \circ \varrho$.*

Corollary 3.12. *Let A and B be BCC-ideals. If congruences ϱ_A and ϱ_B commute, then*

$$\bigcup_{a \in A} B_a = \bigcup_{b \in B} A_b$$

is a BCC-ideal.

Proof. Let $\varrho_A \circ \varrho_B = \varrho$. Then by Lemma 3.2

$$\begin{aligned} \bigcup_{a \in A} B_a &= \{x \in B : x \varrho_B a \text{ for some } a \in A\} \\ &= \{x \in G : x \varrho_B a \text{ and } a \varrho_A 0\} = \{x \in G : x \varrho 0\} = C_0 \end{aligned}$$

is a BCC-ideal. Since $\varrho_A \circ \varrho_B = \varrho_B \circ \varrho_A$ then

$$\bigcup_{a \in A} B_a = \bigcup_{b \in B} A_b.$$

□

4. MAXIMAL IDEALS

A proper ideal is called *maximal* iff it is not properly contained in any proper ideal of the same type. A BCC-algebra without proper BCC-ideals (BCK-ideals) is called *BCC-simple* (*BCK-simple*). Obviously any BCK-simple BCC-algebra is BCC-simple. The converse is not true. A BCC-algebra \mathbf{G} given in our Example 2.1 is BCC-simple, but it is not BCK-simple because it has two maximal BCK-ideals $A = \{0, 1\}$ and $B = \{0, 2\}$.

A BCC-simple BCC-algebra has only two regular congruences.

Theorem 4.1. *Let A be a proper BCC-ideal of a BCC-algebra \mathbf{G} . Then A is a maximal BCC-ideal of \mathbf{G} iff \mathbf{G}/A is a BCC-simple BCC-algebra.*

Proof. Let \mathbf{G}/A be a BCC-simple BCC-algebra. If A is not a maximal BCC-ideal, then there exists a proper BCC-ideal B such that $A \subset B \subset G$. Obviously B/A is properly contained in G/A and has at least two elements, because $x \in A_x \subset B/A$ for all $x \in B - A$. Obviously $A = A_0 \in B/A$. Moreover, if $A_y \in B/A$ and $A_{xy \cdot z} = (A_x * A_y) * A_z \in B/A$, then $y, xy \cdot z \in B$, which implies $xz \in B$. Therefore

$A_x * A_z \in B/A$. Thus B/A is a proper BCC-ideal of \mathbf{G}/\mathbf{A} , i.e. \mathbf{G}/\mathbf{A} is not simple. This contradiction proves that A is a maximal BCC-ideal.

Conversely, if A is a maximal BCC-ideal of \mathbf{G} and \mathbf{G}/\mathbf{A} is not BCC-simple, then there exists a proper BCC-ideal D of \mathbf{G}/\mathbf{A} . Then $\varphi^{-1}(D)$, where $\varphi(x) = A_x$ is the canonical homomorphism from \mathbf{G} onto \mathbf{G}/\mathbf{A} , is a proper BCC-ideal of \mathbf{G} . Moreover $A = A_0 \subset \varphi^{-1}(D)$ and $A \neq \varphi^{-1}(D)$, which contradicts our hypothesis. Hence \mathbf{G}/\mathbf{A} is simple. The proof is complete. \square

Theorem 4.2. *Any BCC-algebra may be viewed as a maximal BCC-ideal of some BCC-algebra.*

Proof. Corollary 3 in [2] (cf. also Construction 5 in [3]) implies that if S is a (proper) BCC-algebra and $e \notin S$, then $G = S \cup \{e\}$ with the multiplication defined by

$$x * y = \begin{cases} xy & \text{for } x, y \in S, \\ 0 & \text{for } y = e, \\ e & \text{for } x = e, y \neq e \end{cases}$$

is a (proper) BCC-algebra and e is the greatest element of G . It is not difficult to verify that S is a maximal BCC-ideal of \mathbf{G} . \square

Corollary 4.3. *If a BCC-algebra \mathbf{G} has an element e such that $xy = e$ iff $x = e$, $y \neq e$, then $G - \{e\}$ is the maximal BCC-ideal of \mathbf{G} .*

Proof. Assume $xy \cdot z \neq e$ for some $y \neq e$. Then $xz \neq e$. If not, then $xz = e$, by the assumption, implies $x = e$, $z \neq e$. Hence $xy \cdot z = ey \cdot z = ez = e$, which is impossible. \square

Corollary 4.4. *If a BCC-algebra \mathbf{G} has an element e such that $G \setminus \{e\}$ is a BCC-ideal (BCK-ideal), then $ey = e$ for all $y \neq e$ and e is the maximal element of \mathbf{G} .*

References

- [1] *W. A. Dudek:* The number of subalgebras of finite BCC-algebras. Bull. Inst. Math. Academia Sinica 20 (1992), 129–136.
- [2] *W. A. Dudek:* On proper BCC-algebras. Bull. Inst. Math. Academia Sinica 20 (1992), 137–150.
- [3] *W. A. Dudek:* On constructions of BCC-algebras. Selected Papers on BCK-and BCI-algebras, 1 (1992), 93–96.
- [4] *K. Iseki and S. Tanaka:* Ideal theory of BCK-algebras. Math. Japonica 21 (1976), 351–366.
- [5] *K. Iseki and S. Tanaka:* An introduction to the theory of BCK-algebras. Math. Japonica 23 (1978), 1–26.

- [6] *Y. Komori*: The variety generated by BCC-algebras is finitely based. Reports Fac. Sci. Shizuoka Univ. *17* (1983), 13–16.
- [7] *Y. Komori*: The class of BCC-algebras is not a variety. Math. Japonica *29* (1984), 391–394.
- [8] *A. Wroński*: BCK-algebras do not form a variety. Math. Japonica *28* (1983), 211–213.

Authors' addresses: Wiesław A. Dudek Institute of Mathematics, Wrocław University of Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław Poland, e-mail: dudek@im.pwr.wroc.pl; Xiaohong Zhang, Department of Mathematics, Hanzhong Teachers College, Hanzhong, Shaanxi Province, Peoples Republic of China.