

L. Nochefranca; Kar-Ping Shum

Pseudo-symmetric ideals of semigroup and their radicals

*Czechoslovak Mathematical Journal*, Vol. 48 (1998), No. 4, 727–735

Persistent URL: <http://dml.cz/dmlcz/127450>

## Terms of use:

© Institute of Mathematics AS CR, 1998

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

PSEUDO-SYMMETRIC IDEALS OF SEMIGROUP  
AND THEIR RADICALS

L. NOCHEFRANCA, Quezon City, and K. P. SHUM, Hong Kong

(Received December 5, 1995)

## 1. INTRODUCTION

Pseudo-symmetric ideals and pseudo-symmetric semigroups were studied for the first time by A. Anjaneyulu in 1980 ([2]). The aim of this paper is to establish some further properties of such ideals and semigroups. Also, the radicals of a pseudo-symmetric ideal of an arbitrary semigroup will be characterized. In fact, we will see that the concept of “pseudo-symmetry” plays an important role in the study of radicals of semigroups.

Let  $S$  be a semigroup and  $I$  an ideal of  $S$ . Then we have the following well known definitions:

- (i)  $I$  is prime  $\iff$  for all  $x, y \in S$ ,  $xSy \subseteq I$  implies  $x \in I$  or  $y \in I$ ;
- (ii)  $I$  is completely prime  $\iff$  for all  $x, y \in S$ ,  $xy \in I$  implies  $x \in I$  or  $y \in I$ ;
- (iii)  $I$  is semiprime  $\iff$  for all  $x \in S$ ,  $xSx \subseteq I$  implies  $x \in I$ ;
- (iv)  $I$  is completely semiprime  $\iff$  for all  $x \in S$ ,  $x^2 \in I$  implies  $x \in I$ .

In addition, if  $S$  has a zero element  $0$ , then

- (v)  $I$  is nilpotent  $\iff I^n = 0$  for some integer  $n > 0$ ;
- (vi)  $I$  is nil  $\iff x$  is nilpotent for all  $x \in I$ ;
- (vii)  $I$  is locally nilpotent  $\iff$  the subsemigroup generated by any finite number of elements of  $I$  is nilpotent.

The following radicals occurred in J. Bosák [3]. Let  $S$  be a semigroup and  $A$  an ideal of  $S$ . Then we define:

---

The second author would like to thank the University of the Philippines for the hospitality extended to him during the period of his visit to the Department of Mathematics, January-March 1995. His research is partially supported by a CUHK direct grant # 2060126

- (i)  $R_A(S)$ : the union of all ideals of  $S$  nilpotent with respect to  $A$ , i.e.,  $R_A(S) = \bigcup_{i \in \Lambda} I_i$ , where  $I_i$  is an ideal of  $S$  for each  $i$  and  $I_i^{n_i} \subseteq A$  for some  $n_i \geq 1$ ;
- (ii)  $M_A(S)$ : the intersection of all prime ideals of  $S$  containing  $A$ ;
- (iii)  $L_A(S)$ : the union of all ideals of  $S$  locally nilpotent with respect to  $A$ ;
- (iv)  $R_A^*(S)$ : the union of all ideals of  $S$  nil with respect to  $A$ ;
- (v)  $N_A(S)$ : the set of all elements of  $S$  nilpotent with respect to  $A$ ;
- (vi)  $C_A(S)$ : the intersection of all completely prime ideals of  $S$  containing  $A$ .

The following beautiful result was also given in J. Bosák [3].

**Lemma 1.1.** ([3] Theorem 2) *Let  $A$  be an ideal of a semigroup  $S$ . Then*

- (i)  $A \subseteq R_A(S) \subseteq M_A(S) \subseteq L_A(S) \subseteq R_A^*(S) \subseteq N_A(S) \subseteq C_A(S) \subseteq S$ ;
- (ii) *there exists a periodic semigroup  $U$  with a zero element  $0$  such that*

$$0 \subset R_0(U) \subset M_0(S) \subset L_0(S) \subset R_0^*(S) \subset N_0(S) \subset C_0(S) \subset S,$$

where “ $\subset$ ” means the proper inclusion.

**Remark 1.2.** Let  $A$  be an ideal. Then, by Lemma 1.1, we can see that all the above radicals, except  $N_A(S)$  are ideals of  $S$ .

**Remark 1.3.** It is clear that

$$R_A(S) = \{x \in S \mid \langle x \rangle^n \subseteq A \text{ for some integer } n > 0\},$$

where  $\langle x \rangle$  means the principal ideal generated by  $x$ .

**Definition 1.4.** Let  $S$  be a semigroup and  $T$  a non-empty subset of  $S$ . We call  $T$  a pseudo-symmetric subset of  $S$  if for all  $x, y \in S$ ,  $xy \in T$  implies  $xSy \subseteq T$ . An ideal  $A$  of  $S$  is called a pseudo-symmetric ideal if  $A$  is also a pseudo-symmetric subset. A semigroup  $S$  is said to be pseudo-symmetric if every ideal of  $S$  is pseudo-symmetric.

**Remark 1.5.** All normal, quasi-commutative, left zero, right zero semigroups and bands are pseudo-symmetric semigroups (see [2] and [7]).

The following example shows that a pseudo-symmetric subset of  $S$  need not be a subsemigroup of  $S$ .

**Example 1.6.** Let  $S = \{0, a, b, c\}$  with the following Cayley table:

	0	a	b	c
0	0	0	0	0
a	0	0	0	0
b	0	0	a	a
c	0	0	0	a

Then  $T = \{0, c\}$  is clearly a pseudo-symmetric subset but not a subsemigroup of  $S$ .

In this paper, we will prove that  $C_A(S) = R_A(S)$  if  $A$  is a pseudo-symmetric ideal of the semigroup  $S$ . We will also give a characterization for the radical  $N_A(S)$  to be an ideal of  $S$ . Some results in [1] are generalized.

## 2. PSEUDO-SYMMETRIC IDEALS

We first discuss the relationships among prime, completely prime and pseudo-symmetric ideals.

**Proposition 2.1.** *Let  $S$  be a semigroup. Then the following statements hold:*

- (i) *Every completely prime ideal is both prime and pseudo-symmetric.*
- (ii) *Let  $A$  be a pseudo-symmetric ideal of  $S$ . Then  $A$  is prime  $\iff A$  is completely prime.*
- (iii) *Let  $A$  be a prime ideal of  $S$ . Then  $A$  is pseudo-symmetric  $\iff A$  is completely prime.*


*Proof.* (i) This statement is easy to observe. We hence omit the proof.

(ii) This result follows from Lemma 1 in [2].

(iii) ( $\implies$ ) Let  $A$  be a pseudo-symmetric ideal of  $S$ . If  $xy \in A$  for some  $x, y \in S$ , then  $xSy \subseteq A$ . Since  $A$  is prime, we have  $x \in A$  or  $y \in A$ . This shows that  $A$  is completely prime.

( $\impliedby$ ) This part follows immediately from (i). □

In general, we have the following diagram:



**Example 2.2.** Let  $S = \{0, a, \dots, a^{n-1}\}$  be a semigroup with  $a^n = 0$ . Then  $S$  clearly is a commutative semigroup and  $\{0\}$  is a pseudo-symmetric ideal which is neither prime nor completely prime. This shows that  $\boxed{2}$  and  $\boxed{4}$  are valid in the above diagram. The following example shows that  $\boxed{1}$  and  $\boxed{3}$  hold in the above diagram.

**Example 2.3.** Let  $S = \{0, e, f, a, b\}$  be a set with the following Cayley table:

	0	e	f	a	b
0	0	0	0	0	0
e	0	e	0	0	b
f	0	0	f	a	0
a	0	a	0	0	f
b	0	0	b	e	0

Then  $S$  is a 0-simple semigroup. Clearly,  $\{0\}$  is a prime ideal but not a pseudo-symmetric ideal of  $S$  because  $ef = 0$  but  $ebf = b \neq 0$ . It is obvious that  $\{0\}$  is not completely prime.

A semigroup  $S$  is called a left (right) duo semigroup if every left (right) ideal of  $S$  is a two sided ideal. We call  $S$  a duo semigroup if  $S$  is both a left and a right duo semigroup.

We now describe the relationship between the one-side duo semigroup and the pseudo-symmetric semigroup.

**Proposition 2.4.** *A left (right) duo semigroup is pseudo-symmetric.*

*Proof.* Let  $S$  be a left duo semigroup. Clearly, for any  $x \in S$ ,  $x \cup Sx$  is a left ideal of  $S$  containing  $x$ . Then we have  $xS \subseteq x \cup Sx$  for all  $x \in S$  since  $S$  is a left duo semigroup. Now, let  $A$  be an arbitrary ideal of  $S$  with  $xy \in A$  for some  $x, y \in S$ . Then we have  $xSy \subseteq (x \cup Sx)y \subseteq xy \cup Sxy \subseteq A$ . This shows that  $A$  is a pseudo-symmetric ideal and hence  $S$  is a pseudo-symmetric semigroup.

Similarly, we can prove that if  $S$  is a right duo semigroup then  $S$  is a pseudo-symmetric semigroup. □

The following example illustrates that a pseudo-symmetric semigroup need not be a left (right) duo semigroup.

**Example 2.5.** Let  $S = \{0, a, b, c\}$  be a set with the following Cayley table:

	0	a	b	c
0	0	0	0	0
a	0	0	a	0
b	0	0	b	0
c	0	a	a	c

Then it is clear that

- (i)  $S$  contains five ideals, namely,  $I_1 = \{0\}$ ,  $I_2 = \{0, a\}$ ,  $I_3 = \{0, a, c\}$ ,  $I_4 = \{0, a, b\}$  and  $I_5 = S$ . These ideals are all pseudo-symmetric;
- (ii)  $S$  is neither a left nor a right duo semigroup since  $bS = \{0, b\}$  and  $Sc = \{0, c\}$  are not ideals of  $S$ .

### 3. RADICALS

**Lemma 3.1.** *Let  $S$  be a semigroup and  $A$  an ideal of  $S$ . Then*

- (i) *for any  $x \in S$ ,  $x \in N_A(S) \iff x^2 \in N_A(S)$ ;*
- (ii) *for any  $x, y \in S$ ,  $xy \in N_A(S) \iff yx \in N_A(S)$ .*

*Proof.* (i) If  $x \in N_A(S)$  then there exists  $n > 0$  such that  $x^n \in A$ . Since  $A$  is an ideal, we have  $x^{2n} = (x^2)^n \in A$ . This leads to  $x^2 \in N_A(S)$ .

Conversely, if  $x^2 \in N_A(S)$ , then there exists  $n > 0$  such that  $(x^2)^n = x^{2n} \in A$ . Therefore,  $x \in N_A(S)$ .

(ii) Let  $xy \in N_A(S)$ . Then we have  $(xy)^n \in A$  for some  $n > 0$ . We can assume that  $n \geq 2$ . Since  $x(yx)^{n-1}y = (xy)^n \in A$ ,  $(yx)^{n+1} \in A$ . Therefore,  $yx \in N_A(S)$ . The converse can be proved similarly.  $\square$

The following theorem is the main theorem of this paper. It shows that the concept “pseudo-symmetry” is essential for the radical  $N_A(S)$  to become an ideal of  $S$ .

**Theorem 3.2.** *Let  $A$  be an ideal of a semigroup  $S$ . Then the following statements are equivalent:*

- (i)  $N_A(S)$  is an ideal of  $S$ .
- (ii)  $N_A(S)$  is a pseudo-symmetric ideal of  $S$ .
- (iii)  $N_A(S)$  is a pseudo-symmetric subset of  $S$ .
- (iv)  $N_A(S)$  is a one-side ideal of  $S$ .
- (v)  $R_A^*(S) = N_A(S) = C_A(S)$ .

*Proof.* (ii)  $\implies$  (iii) and (v)  $\implies$  (i) are clear.

(i)  $\implies$  (ii). Assume that  $N_A(S)$  is an ideal of  $S$ . Then, by Lemma 3.1 (ii), for any  $x, y \in S$  we know that  $xy \in N_A(S)$  implies  $yx \in N_A(S)$ . Suppose that  $yx \in N_A(S)$  for some  $x, y \in S$ . Since  $N_A(S)$  is an ideal of  $S$  we have  $(yx)s \in N_A(S)$  for any  $s \in S$ . Thus, by using Lemma 3.1(ii) again, we have  $xsy \in N_A(S)$ . This shows that  $xSy \subseteq N_A(S)$ . Consequently,  $N_A(S)$  is a pseudo-symmetric ideal of  $S$ .

(iii)  $\implies$  (iv). Let  $N_A(S)$  be a pseudo-symmetric subset of  $S$ . Then, by Lemma 3.1, for all  $x \in N_A(S)$  and  $s \in S$ , we have

$$\begin{aligned} x \in N_A(S) &\implies x^2 \in N_A(S) \implies xsx \in N_A(S) \\ &\implies x(xs) \in N_A(S) \implies xs(xs) = (xs)^2 \in N_A(S) \implies xs \in N_A(S). \end{aligned}$$

This shows that  $N_A(S)$  is a right ideal of  $S$ . The fact that  $N_A(S)$  is a left ideal can be proved similarly.

(iv)  $\implies$  (v). Without loss of generality, we may assume that  $N_A(S)$  is a left ideal of  $S$ . Then for all  $x \in N_A(S)$  and all  $s \in S$  we have  $sx \in N_A(S)$ . It follows from

Lemma 3.1(ii) that  $xs \in N_A(S)$ . This shows that  $N_A(S)$  is an ideal of  $S$ . Therefore,  $R_A^*(S) = N_A(S)$ . Also, it is obvious that  $N_A(S)$  is a completely semiprime ideal of  $S$ . Thus, it follows from Theorem II. 3.7 in [6] that  $N_A(S)$  is the intersection of some completely prime ideals of  $S$  and so  $N_A(S) \supseteq C_A(S)$ . By using Lemma 1.1, we obtain  $N_A(S) = C_A(S)$ .  $\square$

**Theorem 3.3.** *Let  $A$  be an ideal of a semigroup  $S$ . Then the following statements are equivalent:*

- (i)  $N_A(S) = A$ .
- (ii)  $A$  is completely semiprime.
- (iii)  $A = R_A(S) = M_A(S) = L_A(S) = R_A^*(S) = N_A(S) = C_A(S)$ .

**Proof.** (i)  $\implies$  (ii). Assume that  $N_A(S) = A$  and  $x^2 \in A$ . Thus  $x^2 \in N_A(S)$  and so  $x \in N_A(S) = A$  by Lemma 3.1. This shows that  $A$  is completely semiprime.

(ii)  $\implies$  (i). Let  $A$  be completely semiprime. We only need to prove that  $N_A(S) \subseteq A$ . For any  $x \in N_A(S)$  there exists  $n > 0$  such that  $x^n \in A$ . This leads to  $(x^{\frac{n}{2}})^2$  or  $(x^{\frac{n+1}{2}})^2 \in A$ . As a consequence, we have  $x^{\frac{n}{2}}$  or  $x^{\frac{n+1}{2}} \in A$ . Notice that  $\frac{n}{2}, \frac{n+1}{2} < n$ , so by induction on  $n$ , we can eventually obtain that  $x \in A$ . This shows that  $N_A(S) = A$ .

(iii)  $\implies$  (i). Clear.

(i)  $\implies$  (iii). By Theorem 3.2 and Lemma 1.1 we can easily obtain the required result.  $\square$

The following theorem gives a condition for the radicals described by J. Bosák in [3] to be equal.

**Theorem 3.4.** *Let  $S$  be an arbitrary semigroup and  $A$  a pseudo-symmetric ideal of  $S$ . Then the following equalities hold:*

$$R_A(S) = M_A(S) = L_A(S) = R_A^*(S) = N_A(S) = C_A(S).$$

**Proof.** In view of Lemma 1.1 and Theorem 3.2, we only need to prove that  $N_A(S) \subseteq R_A(S)$ . For this purpose, we let  $x \in N_A(S)$  and  $s \in S$ . Then we have  $x^n \in A$  for some  $n \geq 1$ . If  $n = 1$  then clearly  $x \in A \subseteq R_A(S)$ . If  $n > 1$ , then we let  $S^1$  be the semigroup adjoint with an identity 1. Since the ideal  $A$  is pseudo-symmetric, we have  $\langle x \rangle x^{n-1} = (S^1 x S^1) x^{n-1} \subseteq S^1 A \subseteq A$ . By using induction on  $n$ , we can easily obtain that  $\langle x \rangle^{n-1} x \subseteq A$ . Thus,

$$\langle x \rangle^n = \langle x \rangle^{n-1} (S^1 x S^1) = (\langle x \rangle^{n-1} S^1 x) S^1 \subseteq (\langle x \rangle^{n-1} x) S^1 \subseteq A.$$

Therefore,  $x \in R_A(S)$ . The proof is completed.  $\square$

We would like to point out here that the converse of Theorem 3.4 is not true. This can be illustrated by the following example:

**Example 3.5.** Let  $S = \{0, a, b, c\}$  be a set with the following Cayley table:

	0	a	b	c
0	0	0	0	0
a	0	0	a	0
b	0	0	b	0
c	0	a	0	c

Then  $S$  is a semigroup. Moreover, we have the following:

- (i)  $S$  contains five ideals, namely,  $I_1 = \{0\}$ ,  $I_2 = \{0, a\}$ ,  $I_3 = \{0, a, b\}$ ,  $I_4 = \{0, a, c\}$  and  $I_5 = S$ .
- (ii) All the above ideals except  $\{0\}$  are pseudo-symmetric ideals.
- (iii)  $I_1$  and  $I_2$  are not prime ideals, but  $I_3$ ,  $I_4$  and  $I_5$  are all completely prime ideals.
- (iv)  $R_0(S) = I_2 = C_0(S)$ .

By Proposition 2.4, we can immediately deduce the following result obtained by A. Anjaneyulu in [1].

**Corollary 3.6.** ([1] Proposition 1.3) *For any ideal  $A$  in a left (right) duo semigroup  $S$ , we have  $R_A(S) = M_A(S) = N_A(S)$ .*

Finally, we discuss the one-sided primary ideal given in [1].

**Definition 3.7.** ([1]) Call an ideal  $A$  of a semigroup  $S$  left (right) primary provided that the following conditions hold:

- (i) If  $X, Y$  are ideals of  $S$  such that  $XY \subseteq A$  and  $Y \not\subseteq A$  ( $X \not\subseteq A$ ) then  $X \subseteq M_A(S)$  ( $Y \subseteq M_A(S)$ );
- (ii)  $M_A(S)$  is a prime ideal of  $S$ .

The following theorem gives a characterization for a pseudo-symmetric ideal to be a one-sided primary ideal.

**Theorem 3.8.** *Let  $S$  be a semigroup and  $A$  a pseudo-symmetric ideal of  $S$ . Then  $A$  is left primary if and only if the following condition holds:*

- (\*) *for all  $x, y \in S$ ,  $xy \in A$  and  $y \notin A$  imply  $x^n \in A$  for some  $n > 0$ .*

**P r o o f.**  $\implies$ ) Suppose that  $A$  is a left primary ideal of  $S$  and  $xy \in A$  with  $y \notin A$ . Then, since  $A$  is pseudo-symmetric and  $xy \in A$ , we have  $\langle x \rangle \langle y \rangle = S^1(xS^1S^1y)S^1 \subseteq$



$S^1AS^1 \subseteq A$ . Thus, by Definition 3.7, we have  $\langle x \rangle \subseteq M_A(S)$ . By Theorem 3.4 we have  $M_A(S) = N_A(S)$ . This implies that  $x \in N_A(S)$  and so  $x^n \in A$  for some  $n \geq 1$ . Hence, (\*) holds.

$\Leftarrow$ ) Suppose that (\*) holds. Then we have the following situations:

- (i)  $X$  and  $Y$  are ideals of  $S$  with  $XY \subseteq A$  but  $Y \not\subseteq A$ . Then there exists an element  $y \in Y$  but  $y \notin A$  such that for all  $x \in X$ ,  $xy \in XY \subseteq A$ . By (\*) and Theorem 3.4, we immediately obtain that  $x \in M_A(S)$  for all  $x \in X$ . This implies that  $X \subseteq M_A(S)$ .
- (ii) Assume that  $xy \in M_A(S)$ . Then we have  $xy \in N_A(S)$  and hence we can find a smallest positive integer  $n$  such that  $(xy)^n \in A$ . If  $n = 1$  then  $xy \in A$ . By (\*) we have  $x^k \in A$  for some integer  $k > 0$  or  $y \in A$ . This means that  $x \in M_A(S)$  or  $y \in M_A(S)$ . Now, we assume that  $n > 1$ . We have the following cases:
  - Case (i). If  $y(xy)^{n-1} \notin A$ , then by (\*) and  $x(y(xy)^{n-1}) = (xy)^n \in A$  we have  $x^n \in A$  for some  $n > 0$ . This implies that  $x \in N_A(S) = M_A(S)$ .
  - Case (ii). If  $y(xy)^{n-1} \in A$  then since  $(xy)^{n-1} \notin A$ , we have  $y \in N_A(S) = M_A(S)$ .

Hence, in all cases we must have  $x \in M_A(S)$  or  $y \in M_A(S)$ . This shows that  $M_A(S)$  is completely prime, and so  $M_A(S)$  is prime.  $\square$

By Proposition 2.4, we re-deduce the following result in [1]:

**Corollary 3.9.** ([1] Theorem 2.4) *If  $S$  is a one-side duo-semigroup then an ideal  $A$  of  $S$  is left primary if and only if (\*) holds.*

**Remark 3.10.** It is well known that every ideal of a one-side duo semigroup has a primary decomposition. Unfortunately, we can not extend this result to pseudo-symmetric semigroups. (See [1] Example 2.2)

**Acknowledgement.** The authors would like to thank the referee for his valuable comments which helped to improve the original version of this paper substantially. We also thank Chen Yuqun for his constructive suggestions given to this paper.

#### References

- [1] A. Anjaneyulu: Primary ideals in semigroups. *Semigroup Forum* 20 (1980), 129–144.
- [2] A. Anjaneyulu: On primary semigroups. *Czechoslovak Math. J.* 30 (105) (1980), 382–386.
- [3] J. Bosák: On radicals of semigroups. *Mat. Časop.* 18 (1968), 204–212.
- [4] J. M. Howie: *An Introduction to Semigroup Theory*. Academic Press, 1976.
- [5] H. Lai: Commutative semi-primary semigroups. *Czechoslovak Math. J.* 25 (100) (1975), 1–3.
- [6] M. Pertriš: *Introduction to Semigroups*. Merill, Ohio, USA, 1973.

- [7] *B. PONDĚLÍČEK*: On weakly commutative semigroups. *Czechoslovak Math. J.* *25 (100)* (1975), 20–23.
- [8] *M. SATYANARAYANA*: Commutative primary semigroups. *Czechoslovak Math. J.* *22 (97)* (1972), 509–516.
- [9] *M. SATYANARAYANA*: Structure and ideal theory of commutative semigroups. *Czechoslovak Math. J.* *28 (103)* (1978), 171–180.
- [10] *S. SCHWARZ*: Prime ideals and maximal ideals in semigroups. *Czechoslovak Math. J.* *29 (94)* (1969), 72–79.
- [11] *O. ZARISKI and P. SAMUEL*: *Commutative Algebra*, vol 1. Von Nostrand, Princeton, 1958.

*Authors' addresses:* L. N o c h e f r a n c a, Department of Mathematics, University of the Philippines, Quezon City, The Philippines; K. P. S h u m, Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong.