

Judita Lihová

On posets with isomorphic interval posets

Czechoslovak Mathematical Journal, Vol. 49 (1999), No. 1, 67–80

Persistent URL: <http://dml.cz/dmlcz/127468>

Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON POSETS WITH ISOMORPHIC INTERVAL POSETS

JUDITA LIHOVÁ, Košice

(Received April 4, 1996)

Let $\mathbb{A} = (A, \leq)$ be a partially ordered set, $\text{Int } \mathbb{A}$ the system of all (nonempty) intervals of \mathbb{A} , partially ordered by the set-theoretical inclusion \subseteq . We are interested in partially ordered sets $\mathbb{B} = (B, \leq)$ with $\text{Int } \mathbb{B}$ isomorphic to $\text{Int } \mathbb{A}$. We are going to show that they correspond to couples of binary relations on A satisfying some conditions. If \mathbb{A} is a directed partially ordered set, the only \mathbb{B} with $\text{Int } \mathbb{B}$ isomorphic to $\text{Int } \mathbb{A}$ are $\mathbb{A}_1^\delta \times \mathbb{A}_2$ corresponding to direct decompositions $\mathbb{A}_1 \times \mathbb{A}_2$ of \mathbb{A} (\mathbb{A}_1^δ denotes the dual of \mathbb{A}_1). The present results include those presented in the paper [11] by V. Slavík. Systems of intervals, particularly of lattices, have been investigated by many authors, cf. [1]–[11].

1.

By an interval of a partially ordered set $\mathbb{A} = (A, \leq)$ a set $\langle a, b \rangle = \{x \in A : a \leq x \leq b\}$ with $a, b \in A, a \leq b$ is meant. If $a = b$, we use the notation $\langle a \rangle$ instead of $\langle a, a \rangle$. The system of all intervals of \mathbb{A} is denoted by $\text{Int } \mathbb{A}$. Consider the set-theoretical inclusion on $\text{Int } \mathbb{A}$. The following lemma is easy to verify:

- 1.1. Lemma.** a) $\langle a, b \rangle = \inf\{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle\}$ if and only if $\langle a, b \rangle = \langle a_1, b_1 \rangle \cap \langle a_2, b_2 \rangle$;
 b) $\langle a, b \rangle = \sup\{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle\}$ if and only if $a = \inf\{a_1, a_2\}, b = \sup\{b_1, b_2\}$.

Let U, V be binary relations on A . Consider the following conditions:

- (P1) $U, V \subseteq \{(x, y) \in A \times A : x \not\parallel y\}$;
 (P2) $x, y \in A, x \leq y \implies$ there exists a unique couple of elements $p, q \in \langle x, y \rangle$ satisfying $pVxUqVyUp$;
 (P3) $u \leq x, y, xVuUy \implies u = \inf\{x, y\}, v = \sup\{x, y\}$ exists and $yVvUx$ holds;
 (P3') $v \geq x, y, yVvUx \implies v = \sup\{x, y\}, u = \inf\{x, y\}$ exists and $xVuUy$ holds;

- (P4) $a = a_1 U a_2 U \dots U a_n = a'$, $a = a'_1 V a'_2 V \dots V a'_m = a'$ ($n, m \in N$) $\implies a = a'$;
(P5) for every $a, a' \in A$ there exist $n, m \in N$, $a_1, \dots, a_n, a'_1, \dots, a'_m \in A$ satisfying
 $a = a_1 U a_2 U \dots U a_n = a'_1 V a'_2 V \dots V a'_m = a'$.

We are going to prove the following theorem:

1.2. Theorem. *Let \mathbb{A} be a connected partially ordered set. Then there exists a mapping Φ of the system of all couples of binary relations U, V on A satisfying the conditions (P1)–(P3') onto the system of all isomorphism classes of partially ordered sets \mathbb{B} with $\text{Int } \mathbb{B}$ isomorphic to $\text{Int } \mathbb{A}$. If a couple (U, V) satisfies (P1)–(P5), then the class $\Phi((U, V))$ consists of all partially ordered sets isomorphic to $\mathbb{A}_1^\delta \times \mathbb{A}_2$ for a direct decomposition $\mathbb{A}_1 \times \mathbb{A}_2$ of \mathbb{A} . Conversely, the class of all partially ordered sets isomorphic to $\mathbb{A}_1^\delta \times \mathbb{A}_2$ for a direct decomposition $\mathbb{A}_1 \times \mathbb{A}_2$ of \mathbb{A} is $\Phi((U, V))$ for a couple (U, V) satisfying (P1)–(P5).*

Let us remark that the connectivity of \mathbb{A} is not a limiting assumption. Namely, if \mathbb{P} is any partially ordered set, P can be decomposed into maximal connected subsets P_i ($i \in I$) and the system $\text{Int } \mathbb{P}$ is the cardinal sum of the interval posets $\text{Int } P_i$ of these subsets. Now a partially ordered set \mathbb{Q} satisfies the condition $\text{Int } \mathbb{Q} \cong \text{Int } \mathbb{P}$ if and only if \mathbb{Q} is the cardinal sum of some \mathbb{Q}_i ($i \in I$) with $\text{Int } \mathbb{Q}_i \cong \text{Int } P_i$.

Further let us notice that if partially ordered sets \mathbb{A}, \mathbb{B} have isomorphic interval posets, then they are of the same cardinality; so we may assume, without loss of generality, that \mathbb{A}, \mathbb{B} have the same underlying set.

2.

Let $\mathbb{A} = (A, \leq)$ be a connected partially ordered set, U, V binary relations on A satisfying (P1)–(P3'). First we will show some properties of U, V following from the conditions (P1)–(P3').

The following is obtained immediately, using (P2).

2.1. Lemma. *The relations U, V are reflexive.*

2.2. Lemma. *The relations U, V are symmetric.*

Proof. Let xUy . By (P1) x, y are comparable. Suppose, e.g., that $x \leq y$. We have $x \leq x, y, xVxUy$ and since $y = \sup\{x, y\}$, using (P3) we obtain yUx . To prove yUx for $x \geq y$, we use (P3'). □

2.3. Lemma. *If $x, y \in A$ and one of these elements covers the other, then $(x, y) \in U \cup V$.*

This follows immediately from (P2).

2.4. Lemma. *If $(x, y) \in U \cap V$, then $x = y$.*

Proof. Let $(x, y) \in U \cap V$. Without loss of generality we can suppose $x \leq y$. Then both $xUyVy$ and $xUxVy$ hold, so $x = y$ by (P2). \square

2.5. Lemma. *If $x \leq y \leq z$, then $xUyUz$ implies xUz and $xVyVz$ implies xVz .*

Proof. We are going to prove, e.g., the part concerning U . Hence let $x \leq y \leq z$, $xUyUz$. (P2) ensures the existence of an element $p \in \langle x, z \rangle$ with $zUpVx$. Now $x \leq p, y, pVxUy$, so that $\sup\{p, y\} = v$ exists and satisfies $yVvUp$ by (P3). Evidently $v \leq z$. We have $y \leq v, z, vVyUz$, so in view of (P3) we obtain $y = \inf\{v, z\} = v$. But then $x = \inf\{p, y\} = p$ and consequently xUz . \square

2.6. Lemma. *Let $x, y \in A$, $x \leq y$, p, q be as in (P2). If $a \in \langle x, y \rangle$, there exists a unique quadruple of elements $p_1 \in \langle x, p \rangle$, $q_1 \in \langle x, q \rangle$, $p_2 \in \langle p, y \rangle$, $q_2 \in \langle q, y \rangle$ satisfying $aUp_1VxUq_1VaUq_2VyUp_2Va$, p_1VpUp_2 , q_1UqVq_2 .*

Proof. Let $a \in \langle x, y \rangle$. Then $x \leq a$ implies the existence of $p_1, q_1 \in \langle x, a \rangle$ satisfying $p_1VxUq_1VaUp_1$ and $a \leq y$ implies that $p_2VaUq_2VyUp_2$ for some $p_2, q_2 \in \langle a, y \rangle$, by (P2). Using again (P2) we obtain that there exist $p' \in \langle p_1, p_2 \rangle$, $q' \in \langle q_1, q_2 \rangle$ such that $p_1Vp'Up_2$, $q_1Uq'Vq_2$. But then 2.5 yields $p'VxUq'VyUp'$. The uniqueness of p, q in (P2) implies $p' = p, q' = q$. The uniqueness of p_1, q_1, p_2, q_2 follows from (P3) and (P3'). Namely, $p_1 = \inf\{p, a\}$, $q_1 = \inf\{a, q\}$, $p_2 = \sup\{p, a\}$, $q_2 = \sup\{a, q\}$. \square

2.7. Lemma. *If $x \leq a \leq y$, then xUy implies $xUaUy$ and xVy implies $xVaVy$.*

Proof. Let $x \leq a \leq y$, xUy . Using the notation as in 2.6, we have $p = x$, $q = y$, $p_1 = x$, $q_1 = a$, $p_2 = a$, $q_2 = y$. By 2.6 we have pUp_2Uy , hence $xUaUy$. The part concerning V can be shown analogously. \square

2.8. Lemma. *Let $x, y \in A$, $x \leq y$, p, q be as in (P2). Then for each $a \in \langle x, y \rangle$, $\inf\{p, a\}$, $\inf\{a, q\}$ exist and they satisfy $pV\inf\{p, a\}UaV\inf\{a, q\}Uq$. The mapping $\alpha: a \mapsto (\inf\{p, a\}, \inf\{a, q\})$ is an isomorphism of $\langle x, y \rangle$ onto $\langle x, p \rangle \times \langle x, q \rangle$.*

Proof. Let $a \in \langle x, y \rangle$, p_1, q_1 be as in 2.6. Then $p_1 = \inf\{p, a\}$, $q_1 = \inf\{a, q\}$ by (P3). Further, 2.6 ensures that pVp_1UaVq_1Uq holds. Now using (P3') and 2.6 we obtain $a = \sup\{p_1, q_1\}$. Let $p'_1 \in \langle x, p \rangle$, $q'_1 \in \langle x, q \rangle$. Since $x \leq p'_1, q'_1$ and $p'_1VxUq'_1$ holds, by 2.7, the condition (P3) yields that $\sup\{p'_1, q'_1\} = a'$ exists and we have $q'_1Va'Up'_1$. But then $p'_1 = \inf\{p, a'\}$, $q'_1 = \inf\{a', q\}$, so that $\alpha(a') = (p'_1, q'_1)$. We have proved that α is onto.

Let $a, a' \in \langle x, y \rangle$, $a \leq a'$. Then evidently $(\inf\{p, a\}, \inf\{a, q\}) \leq (\inf\{p, a'\}, \inf\{a', q\})$. Hence α preserves the order.

Finally, let $a, a' \in \langle x, y \rangle$, $(\inf\{p, a\}, \inf\{a, q\}) \leq (\inf\{p, a'\}, \inf\{a', q\})$. Then $a = \sup\{\inf\{p, a\}, \inf\{a, q\}\} \leq \sup\{\inf\{p, a'\}, \inf\{a', q\}\} = a'$, completing the proof. \square

2.9. Lemma. *Let $x, y \in A$, $x \leq y$, p, q be as in (P2). If $x \leq a \leq a' \leq y$ and $aUa'(aVa')$, then $\inf\{p, a\} = \inf\{p, a'\}(\inf\{a, q\} = \inf\{a', q\})$.*

Proof. Suppose that $x \leq a \leq a' \leq y$ and, e.g., aUa' . Using 2.8 we get $\inf\{p, a\} \leq \inf\{p, a'\}$, $pV\inf\{p, a\}UaUa'$. Now $\inf\{p, a'\} \in \langle \inf\{p, a\}, a' \rangle$, so that $\inf\{p, a\}U\inf\{p, a'\}$. But simultaneously $\inf\{p, a\}V\inf\{p, a'\}$ by 2.7. Hence $\inf\{p, a\} = \inf\{p, a'\}$ by 2.4. \square

Now we are going to introduce a “new” order on A , corresponding to a couple of U, V satisfying (P1)–(P3').

2.10. Definition. For $x, y \in A$ set $x \leq_1 y$ if there exists $u \in A$, $u \leq x, y$, satisfying $xVuUy$.

2.11. Lemma. *The above defined relation \leq_1 is a partial order.*

Proof. The reflexivity of U, V ensures that $x \leq_1 x$ for each $x \in A$. Let $x \leq_1 y, y \leq_1 x$. Then there exist u_1, u_2 such that $u_1 \leq x, y, xVu_1Uy, u_2 \leq y, x, yVu_2Ux$. Using (P3) we obtain $u_1 = \inf\{x, y\} = u_2$. Hence $(u_1, x) \in U \cap V$ and also $(u_1, y) \in U \cap V$ and consequently $x = u_1 = y$ by 2.4. Let $x \leq_1 y, y \leq_1 z$. Then there exist $u_1, u_2 \in A$ satisfying $u_1 \leq x, y, xVu_1Uy, u_2 \leq y, z, yVu_2Uz$. Using (P3') we obtain that $\inf\{u_1, u_2\} = u$ exists and u_1VuUu_2 holds. But then $u \leq x, z$ and $xVuUz$ by 2.5, so that $x \leq_1 z$. \square

The aim is to prove that $\text{Int}(A, \leq) \cong \text{Int}(A, \leq_1)$. Let $x, y \in A, x \leq y, p, q$ be as in (P2). Then evidently $p \leq_1 q$. Set $f(\langle x, y \rangle) = \langle p, q \rangle_1$, where $\langle p, q \rangle_1 = \{t \in A: p \leq_1 t \leq_1 q\}$. Recall that $\langle x, y \rangle$ is isomorphic to $\langle x, p \rangle \times \langle x, q \rangle$. Now we have:

2.12. Lemma. *The mapping α defined in 2.8 is an isomorphism of $\langle p, q \rangle_1$ onto $\langle x, p \rangle^\delta \times \langle x, q \rangle$.*

Proof. Evidently $a \in \langle x, y \rangle$ if and only if $a \in \langle p, q \rangle_1$ and α is onto. Further let us suppose that $a, a' \in \langle p, q \rangle_1, a \leq_1 a'$. We have to prove $\inf\{p, a\} \geq \inf\{p, a'\}$, $\inf\{a, q\} \leq \inf\{a', q\}$. Take $p_1 = \inf\{p, a\}, q'_1 = \inf\{a', q\}$ and $u \leq a, a'$ satisfying $aVuUa'$. In view of (P3'), $r = \inf\{p_1, u\}, s = \inf\{u, q'_1\}$ exist such that $p_1VrUuVsUq'_1$. But then $pVrUa', aVsUq$, so that $r = \inf\{p, a'\}, s = \inf\{a, q\}$

and we have $r \leq p_1, s \leq q'_1$. Conversely let $a, a' \in \langle p, q \rangle_1, p_1 \geq p'_1, q_1 \leq q'_1$, where $p_1 = \inf\{p, a\}, q_1 = \inf\{a, q\}, p'_1 = \inf\{p, a'\}, q'_1 = \inf\{a', q\}$. Since $x \leq p'_1, q_1$ and $p'_1 V x U q_1, \sup\{p'_1, q_1\} = t$ exists and $q_1 V t U p'_1$. Obviously $t \leq a, a'$. Moreover, $a V q_1$ yields $a V t$ and $p'_1 U a'$ implies $t U a'$. Thus $a \leq_1 a'$. The proof is complete. \square

2.13. Lemma. *The mapping f assigning to $\langle x, y \rangle$ the interval $\langle p, q \rangle_1$ is an isomorphism of $\text{Int}(A, \leq)$ onto $\text{Int}(A, \leq_1)$.*

Proof. Let $r \leq_1 s$. Then there exists $u \leq r, s$ such that $r V u U s$. By (P3), $v = \sup\{r, s\}$ exists and $s V v U r$ holds. Evidently $f(\langle u, v \rangle) = \langle r, s \rangle_1$. The mapping f is onto.

Now let $\langle x, y \rangle \subseteq \langle x_1, y_1 \rangle, f(\langle x, y \rangle) = \langle p, q \rangle_1, f(\langle x_1, y_1 \rangle) = \langle p_1, q_1 \rangle_1$. Take $\inf\{p_1, p\} = p'_1, \inf\{q, q_1\} = q'_1$. We have $p'_1 \leq p_1, p, p_1 V p'_1 U p$, so $p_1 \leq_1 p$. Analogously $q'_1 \leq q, q_1, q V q'_1 U q_1$ ensures $q \leq_1 q_1$. Hence $\langle p, q \rangle_1 \subseteq \langle p_1, q_1 \rangle_1$.

Next suppose that $f(\langle x, y \rangle) = \langle p, q \rangle_1 \subseteq \langle p_1, q_1 \rangle_1 = f(\langle x_1, y_1 \rangle)$. We have to show $\langle x, y \rangle \subseteq \langle x_1, y_1 \rangle$. Let $u \leq p_1, p, p_1 V u U p$ and $v \leq q, q_1, q V v U q_1$. Since $p \geq u, x, x V p U u$ and $q \geq x, v, v V q U x$, there exist $a \leq u, x, b \leq x, v$ satisfying $u V a U x V b U v$, by (P3'). Finally consider $c = \inf\{a, b\}$, whose existence follows from (P3'). We have $p_1 V u V a V c U b U v U q_1$, hence $c = \inf\{p_1, q_1\} = x_1$ by (P3) and (P2). Now obviously $x_1 \leq x$. The relation $y \leq y_1$ can be proved analogously. \square

Summarizing, we have:

2.14. Theorem. *Let $\mathbb{A} = (A, \leq)$ be a connected partially ordered set, U, V binary relations on A satisfying (P1)–(P3'). If \leq_1 is the relation on A defined as in 2.10 with the aid of U, V , then (A, \leq_1) is a partially ordered set with $\text{Int}(A, \leq_1)$ isomorphic to $\text{Int}(A, \leq)$.*

It is easy to see that the couples $U_1 = \{(x, y) \in A \times A: x \not\parallel y\}, V_1 = \{(x, x): x \in A\}$ and $U_2 = \{(x, x): x \in A\}, V_2 = \{(x, y) \in A \times A: x \not\parallel y\}$ satisfy the conditions (P1)–(P3'). The corresponding orders \leq_1, \leq_2 are \leq and \leq^δ , respectively. Some partially ordered sets $\mathbb{A} = (A, \leq)$ have no other orders \leq_1 besides \leq and \leq^δ , satisfying $\text{Int}(A, \leq_1) \cong \text{Int}(A, \leq)$. This is the case e.g. for \mathbb{A} in Fig. 1. On the other hand, it is easy to see that the partially ordered sets in Fig. 2 and Fig. 3 have isomorphic interval systems, but they are neither isomorphic nor dually isomorphic. In fact, the first is the direct product of two copies of \mathbb{A} in Fig. 1, while the other is isomorphic to $\mathbb{A}^\delta \times \mathbb{A}$.

Further assume that U, V satisfy also the conditions (P4), (P5). Define binary relations $\overline{U}, \overline{V}$ on A as follows:




Fig. 1




Fig. 2




Fig. 3

2.15. Definition. For $x, y \in A$ set $x\overline{U}y$ ($x\overline{V}y$) if there exists a finite sequence x_1, x_2, \dots, x_n of elements of A such that $x_1 = x$, $x_n = y$ and every two adjoining elements are in the relation $U(V)$.

The following statement is evident.

2.16. Lemma. *The relations \overline{U} , \overline{V} are equivalence relations.*

Consider the decompositions A/\overline{U} , A/\overline{V} . Denote by $[a]\overline{U}$, $[a]\overline{V}$ the equivalence classes containing the element a .

2.17. Definition. Set $[a]\overline{U} \leq [b]\overline{U}$ ($[a]\overline{V} \leq [b]\overline{V}$) if and only if there exist $a_1 \in [a]\overline{U}$, $b_1 \in [b]\overline{U}$ ($a_1 \in [a]\overline{V}$, $b_1 \in [b]\overline{V}$) satisfying $a_1 \leq b_1$.

2.18. Lemma. *For any $a, b \in A$ the following conditions are equivalent:*

- (1) $[a]\overline{U} \leq [b]\overline{U}$;
- (2) for each $a_1 \in [a]\overline{U}$ there exists $b_1 \in [b]\overline{U}$ with $a_1 \leq b_1$;
- (3) for each $b_1 \in [b]\overline{U}$ there exists $a_1 \in [a]\overline{U}$ with $a_1 \leq b_1$.

Proof. The implications (2) \implies (1), (3) \implies (1) are evident. We are going to prove (1) \implies (2). The proof of (1) \implies (3) would be analogous. So let $[a]\overline{U} \leq [b]\overline{U}$. We can suppose that $a \leq b$. Take any $a_1 \in [a]\overline{U}$. Then there exist x_1, \dots, x_n such that $a = x_1$, $a_1 = x_n$, $x_1 \leq x_2$, $x_2 \geq x_3, \dots, x_{n-1} \geq x_n$, $x_1 U x_2 U \dots U x_n$. Using the conditions (P2), (P3) we can construct elements y_1, y_2, \dots, y_n such that $y_1 \in \langle x_1, b \rangle$, $x_1 V y_1 U b$, $y_2 \geq y_1, x_2$, $x_2 V y_2 U y_1$, $y_3 \in \langle x_3, y_2 \rangle$, $x_3 V y_3 U y_2$, \dots , $y_n \in \langle x_n, y_{n-1} \rangle$, $x_n V y_n U y_{n-1}$. We have $a_1 \leq y_n$, $y_n \in [b]\overline{U}$. \square

Obviously the same holds for \overline{V} .

2.19. Lemma. *The above defined relation \leq on A/\overline{U} is a partial order.*

Proof. The reflexivity is trivial. Further let $[a]\overline{U} \leq [b]\overline{U}$, $[b]\overline{U} \leq [a]\overline{U}$. Then there exist $a_1, a_2 \in [a]\overline{U}$ satisfying $a_1 \leq b \leq a_2$. Take $z \in \langle a_1, a_2 \rangle$ such that $a_1 U z V a_2$. We have $z\overline{U}a_2$ and simultaneously $z V a_2$. Using (P4) we obtain $z = a_2$ and consequently $a_1 \leq b \leq z$, which implies $a_1 U b$ by 2.7. Hence $[b]\overline{U} = [a_1]\overline{U} = [a]\overline{U}$. Finally, let $[a]\overline{U} \leq [b]\overline{U}$, $[b]\overline{U} \leq [c]\overline{U}$. Then there exist $a_1 \in [a]\overline{U}$, $c_1 \in [c]\overline{U}$ such that $a_1 \leq b \leq c_1$ and this implies $[a]\overline{U} \leq [c]\overline{U}$. \square

Evidently the same holds for \bar{V} . The symbol \mathbb{A}/\bar{U} (\mathbb{A}/\bar{V}) will be used for A/\bar{U} (A/\bar{V}) with the order \leq as above.

2.20. Theorem. *Let $\mathbb{A} = (A, \leq)$ be a connected partially ordered set, U, V binary relations on A satisfying (P1)–(P5). If \leq_1 is as in 2.10, then \mathbb{A} is isomorphic to $\mathbb{A}/\bar{U} \times \mathbb{A}/\bar{V}$, while $\mathbb{A}_1 = (A, \leq_1)$ is isomorphic to $(\mathbb{A}/\bar{U})^\delta \times \mathbb{A}/\bar{V}$.*

Proof. Define $\alpha: A \rightarrow A/\bar{U} \times A/\bar{V}$ by $\alpha(a) = ([a]\bar{U}, [a]\bar{V})$. α is onto: Take $([a_1]\bar{U}, [a_2]\bar{V}) \in A/\bar{U} \times A/\bar{V}$. By (P5) there exists $x \in A$ satisfying $a_1\bar{U}x\bar{V}a_2$. Then $\alpha(x) = ([a_1]\bar{U}, [a_2]\bar{V})$.

The implication $a \leq b \implies \alpha(a) \leq \alpha(b)$ is evident. Conversely, let $\alpha(a) \leq \alpha(b)$. Then $[a]\bar{U} \leq [b]\bar{U}$, $[a]\bar{V} \leq [b]\bar{V}$ and consequently $a \leq b_1, b_2$ for some $b_1 \in [b]\bar{U}$, $b_2 \in [b]\bar{V}$. Take $b'_1 \in \langle a, b_1 \rangle$, $b'_2 \in \langle a, b_2 \rangle$ such that aVb'_1Ub_1 , aUb'_2Vb_2 . The condition (P3) yields the existence of $t \geq b'_1, b'_2$ with $b'_2VtUb'_1$. Now $t\bar{U}b$, $t\bar{V}b$, hence $t = b$ by (P4). We have $b \geq a$.

Suppose $a \leq_1 b$. Then there exists $u \leq a, b$ satisfying $aVuUb$ and this implies that $[a]\bar{U} \geq [u]\bar{U} = [b]\bar{U}$, $[a]\bar{V} = [u]\bar{V} \leq [b]\bar{V}$.

Finally, let $[a]\bar{U} \geq [b]\bar{U}$, $[a]\bar{V} \leq [b]\bar{V}$. We have to show $a \leq_1 b$. The assumptions yield the existence of $a_1 \in [a]\bar{U}$, $a_2 \in [a]\bar{V}$ with $a_2 \leq b \leq a_1$. Take $c \in \langle a_2, a_1 \rangle$ satisfying a_2VcUa_1 . Then $c = a$ by (P4). In view of 2.8 $u = \inf\{a, b\}$ exists and $aVuUb$. The proof is complete. \square

3.

Let $\mathbb{A} = (A, \leq)$ be a connected partially ordered set, $\mathbb{A}' = (A, \leq')$ another partially ordered set with the same underlying set and let f be an isomorphism of $\text{Int } \mathbb{A}$ onto $\text{Int } \mathbb{A}'$. The aim is to prove that \mathbb{A}' can be obtained in the way described in the preceding section. Define $f': A \rightarrow A$ by

$$f'(a) = b \iff f(\langle a \rangle) = \langle b \rangle' = \langle b \rangle.$$

($\langle x, y \rangle'$ will mean the set $\{t \in A: x \leq' t \leq' y\}$). Evidently f' is a bijective mapping of A onto A . Consider the following binary relations on A : $U = \{(x, y) \in A \times A: x \leq y \text{ and } f'(x) \leq' f'(y)\} \cup \{(x, y) \in A \times A: x \geq y \text{ and } f'(x) \geq' f'(y)\}$, $V = \{(x, y) \in A \times A: x \leq y \text{ and } f'(x) \geq' f'(y)\} \cup \{(x, y) \in A \times A: x \geq y \text{ and } f'(x) \leq' f'(y)\}$. Evidently U, V satisfy the condition (P1).

3.1. Lemma. *Let $x, y \in A$, $x \leq y$, $f(\langle x, y \rangle) = \langle r, s \rangle'$. Then $r = \inf\{f'(x), f'(y)\}$, $s = \sup\{f'(x), f'(y)\}$ in \mathbb{A}' .*

Proof. Since $\langle x, y \rangle = \sup\{\langle x \rangle, \langle y \rangle\}$, we have $\langle r, s \rangle' = \sup\{f(\langle x \rangle), f(\langle y \rangle)\}$. But $f(\langle x \rangle) = \langle f'(x) \rangle$, $f(\langle y \rangle) = \langle f'(y) \rangle$ so that $r = \inf\{f'(x), f'(y)\}$, $s = \sup\{f'(x), f'(y)\}$ in \mathbb{A}' by 1.1. \square

Taking into account that f^{-1} is also an isomorphism and $(f^{-1})' = (f')^{-1}$, we obtain:

3.2. Lemma. *If $x, y \in A$, $x \leq y$, $f(\langle x, y \rangle) = \langle r, s \rangle'$, $r = f'(p)$, $s = f'(q)$, then $x = \inf\{p, q\}$, $y = \sup\{p, q\}$ in \mathbb{A} .*

3.3. Lemma. *The above defined U, V fulfil (P2).*

Proof. Let $x, y \in A$, $x \leq y$. The previous lemma guarantees the existence of such p, q as we need. Now let $p_1, q_1 \in \langle x, y \rangle$ also satisfy $p_1 V x U q_1 V y U p_1$. The relations $p_1 V x$, $x \leq p_1$ imply $f'(x) \geq' f'(p_1)$ while $y U p_1$, $p_1 \leq y$ imply $f'(p_1) \leq' f'(y)$. Hence $f'(p_1) \leq' r$ by 3.1. Analogously $f'(q_1) \geq' s$. On the other hand $\langle p_1 \rangle, \langle q_1 \rangle \subseteq \langle x, y \rangle$ yields $\langle f'(p_1) \rangle, \langle f'(q_1) \rangle \subseteq \langle r, s \rangle'$ and consequently $r \leq' f'(p_1)$, $f'(q_1) \leq' s$. So we have $f'(p_1) = r = f'(p)$, $f'(q_1) = s = f'(q)$, which implies $p_1 = p$, $q_1 = q$. \square

3.4. Lemma. *Let $x, y \in A$, $x \leq y$, $x U y$ ($x V y$). Then for each $t \in \langle x, y \rangle$ we have $x U t U y$ ($x V t V y$).*

Proof. We will prove, e.g., the part concerning U . Take any $t \in \langle x, y \rangle$. We have $\langle x, t \rangle \subseteq \langle x, y \rangle$, hence $f(\langle x, t \rangle) \subseteq f(\langle x, y \rangle)$. By the assumption $x U y$ we have $f'(x) \leq' f'(y)$ and using 3.1 we obtain $f(\langle x, y \rangle) = \langle f'(x), f'(y) \rangle'$. Let $f(\langle x, t \rangle) = \langle a, b \rangle'$. Then $\langle a, b \rangle' \subseteq \langle f'(x), f'(y) \rangle'$, which implies $f'(x) \leq' a$. On the other hand $a = \inf\{f'(x), f'(t)\}$ in \mathbb{A}' by 3.1, so that $a \leq' f'(t)$. Summarizing we obtain $f'(x) \leq' f'(t)$. We have $x U t$. The relation $t U y$ can be shown analogously. \square

3.5. Lemma. *The above defined U, V fulfil (P3) and (P3').*

Proof. We are going to show that (P3) holds. The condition (P3') can be verified analogously. Let $u \leq x, y$, $x V u U y$. Then $f'(x) \leq' f'(u) \leq' f'(y)$. Let $\langle f'(x), f'(y) \rangle' = f(\langle a, b \rangle)$. Using 3.1 and 3.2 we obtain $a = \inf\{x, y\}$, $b = \sup\{x, y\}$ in \mathbb{A} , $f'(x) = \inf\{f'(a), f'(b)\}$, $f'(y) = \sup\{f'(a), f'(b)\}$ in \mathbb{A}' . Since u is a lower bound of $\{x, y\}$, we infer $u \leq a$. Now $a \in \langle u, x \rangle \cap \langle u, y \rangle$, hence $u U a$ and simultaneously $u V a$ by 3.4. Since $u \leq a$, we have $f'(u) \leq' f'(a)$ and simultaneously $f'(u) \geq' f'(a)$. Then $f'(u) = f'(a)$ and consequently $u = a$. We have proved $u = \inf\{x, y\}$. It remains to show $y V b U x$. Since $b \geq x, y$, we have to verify $f'(x) \leq' f'(b) \leq' f'(y)$, but this is evident. \square

Summarizing, having an isomorphism $f: \text{Int } \mathbb{A} \rightarrow \text{Int } \mathbb{A}'$, we can construct binary relations U, V on A satisfying (P1)–(P3'). Further, using 2.14, we obtain a partially ordered set \mathbb{A}_1 such that $\text{Int } \mathbb{A}$ and $\text{Int } \mathbb{A}_1$ are isomorphic. The following theorem makes clear the relation between \mathbb{A}' and \mathbb{A}_1 .

3.6. Theorem. *Let $\mathbb{A} = (A, \leq)$ be a connected partially ordered set, $\mathbb{A}' = (A, \leq')$ any partially ordered set such that $\text{Int } \mathbb{A}'$ is isomorphic to $\text{Int } \mathbb{A}$. If U, V are defined with the aid of an isomorphism $f: \text{Int } \mathbb{A} \rightarrow \text{Int } \mathbb{A}'$ as above, then the partially ordered set \mathbb{A}_1 corresponding to U, V in the sense of 2.14 is isomorphic to \mathbb{A}' .*

Proof. We are going to show that the mapping f' belonging to the isomorphism $f: \text{Int } \mathbb{A} \rightarrow \text{Int } \mathbb{A}'$ is an isomorphism of \mathbb{A}_1 onto \mathbb{A}' . It is sufficient to prove that $x \leq_1 y$ if and only if $f'(x) \leq' f'(y)$. Let $x \leq_1 y$. Then there exist $u \leq x, y$ satisfying $xVuUy$. Using the definition of U, V we obtain $f'(x) \leq' f'(u) \leq' f'(y)$. Conversely, let $f'(x) \leq' f'(y)$. Considering $\langle a, b \rangle = f^{-1}(\langle f'(x), f'(y) \rangle')$ and using 3.2, 3.1, we obtain $a = \inf\{x, y\}$, $xVaUy$. Hence $x \leq_1 y$. The proof is complete. \square

Notice that if $(A, \leq), (A, \leq_1)$ are as in 2.14, then $\text{Int } (A, \leq), \text{Int } (A, \leq_1)$ are not only isomorphic, but even identical as systems of subsets of A . Moreover, every (A, \leq') satisfying that $\text{Int } (A, \leq')$ is identical with $\text{Int } (A, \leq)$ can be obtained in this way by 3.6.

To have 1.2 completely proved, we add:

3.7. Theorem. *Let $\mathbb{A} = \mathbb{C} \times \mathbb{D}$ be a connected partially ordered set, $\mathbb{A}' = \mathbb{C}^\delta \times \mathbb{D}$, let $f: \text{Int } \mathbb{A} \rightarrow \text{Int } \mathbb{A}'$ be defined by*

$$f(\langle c_1, d_1 \rangle, \langle c_2, d_2 \rangle) = \langle \langle c_2, d_1 \rangle, \langle c_1, d_2 \rangle \rangle'.$$

Then f is an isomorphism and if U, V are defined with the aid of f as at the beginning of this section, they satisfy the conditions (P4) and (P5).

Proof. The assertion that f is an isomorphism is evident. Obviously, f' is the identity mapping, so that $(c_1, d_1)U(c_2, d_2)$ means that $c_1 = c_2$ and simultaneously d_1, d_2 are comparable while $(c_1, d_1)V(c_2, d_2)$ means that c_1, c_2 are comparable and $d_1 = d_2$.

Now let

$$\begin{aligned} (c, d) &= (c_1, d_1)U(c_2, d_2)U \dots U(c_n, d_n) = (c', d'), \\ (c, d) &= (c'_1, d'_1)V(c'_2, d'_2)V \dots V(c'_m, d'_m) = (c', d'). \end{aligned}$$

Then $c = c_1 = c_2 = \dots = c'$ and $d = d'_1 = d'_2 = \dots = d'$, so that $(c, d) = (c', d')$.

Finally, taking any $(c, d), (c', d') \in C \times D$ and using the fact that \mathbb{A} is connected so that \mathbb{C}, \mathbb{D} are connected, too, we can find $c_1, \dots, c_n \in C$, $d_1, \dots, d_m \in D$ such that $c_1 = c, c_n = c', d_1 = d, d_m = d', c_i$ is comparable with c_{i+1} for each $i \in \{1, \dots, n-1\}$ and d_j is comparable with d_{j+1} for each $j \in \{1, \dots, m-1\}$. Then we have $(c, d) = (c_1, d_1)U(c_1, d_2)U \dots U(c_1, d_m) = (c_1, d')V(c_2, d')V \dots V(c_n, d') = (c', d')$. The proof is complete. \square

Notice that the mapping Φ in Theorem 1.2 is not one-to-one, in general. For example, if \mathbb{A} is a selfdual partially ordered set, then both $U_1 = \{(x, y) \in A \times A: x \not\ll y\}$, $V_1 = \{(x, x): x \in A\}$ and $U_2 = \{(x, x): x \in A\}$, $V_2 = \{(x, y) \in A \times A: x \not\ll y\}$ lead to the same isomorphism class of partially ordered sets. In this connection, a natural question arises: under what conditions two couples U_1, V_1 and U_2, V_2 of binary relations on A satisfying (P1)–(P3') give the same isomorphism class of partially ordered sets.

Having a bijection α of A onto A , binary relations U_1, V_1 on A satisfying (P1)–(P3') and the corresponding partial order \leq_1 (in the sense of 2.10), consider the following conditions:

- (C1) $\alpha(x) \leq \alpha(y) \implies$ there exists a unique couple of elements $p, q \in A$ satisfying $\alpha(x) \leq \alpha(p)$, $\alpha(q) \leq \alpha(y)$, $p \leq_1 x$, $y \leq_1 q$;
(C2) $p \leq_1 q \implies \inf\{\alpha(p), \alpha(q)\}$, $\sup\{\alpha(p), \alpha(q)\}$ exist and

$$x = \alpha^{-1}(\inf\{\alpha(p), \alpha(q)\}),$$

$$y = \alpha^{-1}(\sup\{\alpha(p), \alpha(q)\})$$

are the only elements of A satisfying $\alpha(x) \leq \alpha(p)$, $\alpha(q) \leq \alpha(y)$, $p \leq_1 x$, $y \leq_1 q$.

3.8. Theorem. *Let U_1, V_1 and U_2, V_2 be two couples of binary relations on A satisfying (P1)–(P3'), let \leq_1 and \leq_2 be the corresponding partial orders (in the sense of 2.10). If α is an isomorphism of (A, \leq_1) onto (A, \leq_2) , then*

- (1) α fulfils (C1), (C2) and
(2) $U_2 = \{(x, y) \in A \times A: x \leq y \text{ and } \alpha^{-1}(x) \leq_1 \alpha^{-1}(y)\} \cup \{(x, y) \in A \times A: x \geq y \text{ and } \alpha^{-1}(x) \geq_1 \alpha^{-1}(y)\}$, $V_2 = \{(x, y) \in A \times A: x \leq y \text{ and } \alpha^{-1}(x) \geq_1 \alpha^{-1}(y)\} \cup \{(x, y) \in A \times A: x \geq y \text{ and } \alpha^{-1}(x) \leq_1 \alpha^{-1}(y)\}$.

Proof. We start with (2). Obviously $U_2 = \{(x, y) \in A \times A: x \leq y \text{ and } x \leq_2 y\} \cup \{(x, y) \in A \times A: x \geq y \text{ and } x \geq_2 y\}$. Since $r \leq_2 s$ ($r, s \in A$) is equivalent to $\alpha^{-1}(r) \leq_1 \alpha^{-1}(s)$, we have what we need for U_2 . As to V_2 , we proceed analogously.

Now let $\alpha(x) \leq \alpha(y)$. Then there exists a unique couple of $\alpha(p), \alpha(q) \in \langle \alpha(x), \alpha(y) \rangle$ with $\alpha(p)V_2\alpha(x)U_2\alpha(q)V_2\alpha(y)U_2\alpha(p)$ by (P2). The latter is equivalent to $\alpha(p) \leq_2 \alpha(x)$, $\alpha(y) \leq_2 \alpha(q)$ and this holds if and only if $p \leq_1 x$, $x, y \leq_1 q$.

To prove (C2), take $p \leq_1 q$. Then $\alpha(p) \leq_2 \alpha(q)$ and there exists $\alpha(x) \leq \alpha(p)$, $\alpha(q)$ satisfying $\alpha(p)V_2\alpha(x)U_2\alpha(q)$. Using (P3) we obtain that $\alpha(x) = \inf\{\alpha(p), \alpha(q)\}$, $\alpha(y) = \sup\{\alpha(p), \alpha(q)\}$ exists and we have $\alpha(q)V_2\alpha(y)U_2\alpha(p)$. The latter means $\alpha(p) \leq_2 \alpha(y) \leq_2 \alpha(q)$, which is equivalent to $p \leq_1 y \leq_1 q$. The relation $p \leq_1 x \leq_1 q$ follows from $\alpha(p)V_2\alpha(x)U_2\alpha(q)$. Now having x_1, y_1 satisfying $\alpha(x_1) \leq \alpha(p)$, $\alpha(q) \leq \alpha(y_1)$, $p \leq_1 x_1$, $y_1 \leq_1 q$, it is easy to see that $\alpha(p)V_2\alpha(x_1)U_2\alpha(q)V_2\alpha(y_1)U_2\alpha(p)$, which yields $\alpha(x_1) = \inf\{\alpha(p), \alpha(q)\}$, $\alpha(y_1) = \sup\{\alpha(p), \alpha(q)\}$ by (P3) and (P3'). But then $\alpha(x_1) = \alpha(x)$, $\alpha(y_1) = \alpha(y)$ and consequently $x_1 = x$, $y_1 = y$.

The proof is complete. \square

Conversely, we have:

3.9. Theorem. *Let U_1, V_1 be a couple of binary relations on A satisfying (P1)–(P3') and let α be any bijection of A onto A satisfying (C1), (C2). Taking U_2, V_2 as in (2) of 3.8, they satisfy the conditions (P1)–(P3') and α is an isomorphism of (A, \leq_1) onto (A, \leq_2) (\leq_1 and \leq_2 are the partial orders corresponding to U_1, V_1 and U_2, V_2 , respectively, in the sense of 2.10).*

Proof. The relations U_2, V_2 satisfy (P1) trivially. To prove (P2), let $x', y' \in A$, $x' \leq y'$. Take $x, y \in A$ with $\alpha(x) = x'$, $\alpha(y) = y'$. The condition (C1) yields the existence of a unique couple of elements $p, q \in A$ satisfying $\alpha(x) \leq \alpha(p)$, $\alpha(q) \leq \alpha(y)$, $p \leq_1 x$, $y \leq_1 q$. Set $\alpha(p) = p'$, $\alpha(q) = q'$. Then p', q' are the only elements of the interval $\langle x', y' \rangle$ with $p'V_2x'U_2q'V_2y'U_2p'$.

We are going to prove (P3). The proof of (P3') would be analogous. Let $u' \leq p', q'$, $p'V_2u'U_2q'$. Take u, p, q with $\alpha(u) = u'$, $\alpha(p) = p'$, $\alpha(q) = q'$. Using the definition of U_2, V_2 we obtain $p \leq_1 u \leq_1 q$. Now (C2) ensures the existence of $\inf\{\alpha(p), \alpha(q)\}$ and $\sup\{\alpha(p), \alpha(q)\}$, together with $u' = \inf\{\alpha(p), \alpha(q)\}$. It remains to show that $q'V_2\sup\{p', q'\}U_2p'$, which is equivalent to $p \leq_1 \alpha^{-1}(\sup\{p', q'\}) \leq_1 q$. But this holds by (C2).

Finally, we have to prove that having any $x, y \in A$, $x \leq_1 y$ is equivalent to $\alpha(x) \leq_2 \alpha(y)$. Let $x \leq_1 y$. Then there exists $u \in \langle x, y \rangle_1$ such that $\alpha(u) = \inf\{\alpha(x), \alpha(y)\}$ by (C2). Now $\alpha(u) \leq \alpha(x)$ together with $u \geq_1 x$ yields $\alpha(u)V_2\alpha(x)$, while $\alpha(u) \leq \alpha(y)$, $u \leq_1 y$ implies $\alpha(u)U_2\alpha(y)$. Consequently $\alpha(x) \leq_2 \alpha(y)$. Conversely, let $\alpha(x) \leq_2 \alpha(y)$. Then there exists $\alpha(t) \leq \alpha(x), \alpha(y)$ with $\alpha(x)V_2\alpha(t)U_2\alpha(y)$. So we have $x \leq_1 t \leq_1 y$ and the proof is complete. \square

It is easy to see that if α is an automorphism or a dual automorphism of \mathbb{A} , then α satisfies the conditions (C1), (C2) for any U_1, V_1 fulfilling (P1)–(P3') and

the corresponding order \leq_1 . But a bijection α fulfilling (C1), (C2) need not be an isomorphism or a dual isomorphism, as the following example shows:

3.10. Example. Let \mathbb{A} be as in Fig. 4. Let $U_1 = \{(a, b) \in A \times A: a \not\parallel b\}$, $V_1 = \{(t, t): t \in A\}$, $U_2 = V_1$, $V_2 = U_1$, $U_3 = \{(u, y), (y, u), (x, v), (v, x)\} \cup \{(t, t): t \in A\}$, $V_3 = \{(x, u), (u, x), (y, v), (v, y)\} \cup \{(t, t): t \in A\}$, $U_4 = V_3$, $V_4 = U_3$. It is easy to see that all couples satisfying (P1)–(P3') are those of $U_1, V_1, U_2, V_2, U_3, V_3$ and U_4, V_4 . But each of them yields the same isomorphism class of partially ordered sets with the interval system isomorphic to $\text{Int } \mathbb{A}$. E.g., the mapping α such that $\alpha(u) = v$, $\alpha(v) = u$, $\alpha(x) = x$, $\alpha(y) = y$ mediates the transition from U_1, V_1 to U_2, V_2 , while Ψ defined by $\Psi(u) = y$, $\Psi(y) = v$, $\Psi(v) = x$, $\Psi(x) = u$ is an intermediary between U_1, V_1 and U_4, V_4 .




Fig. 4

The following question remains open:

Let $\mathbb{A}_1 \times \mathbb{A}_2$ be a direct decomposition of \mathbb{A} . Consider the class of all partially ordered sets isomorphic to $\mathbb{A}_1^\delta \times \mathbb{A}_2$. Does every pre-image (U, V) of this class under Φ satisfy (P4), (P5)? In particular, if the class $\Phi((U, V))$ consists of all partially ordered sets isomorphic to \mathbb{A} (or \mathbb{A}^δ), does (U, V) satisfy (P4), (P5)?

4.

In this section we will apply the foregoing results to the case of a directed partially ordered set.

4.1. Lemma. Let $\mathbb{A} = (A, \leq)$ be a directed partially ordered set, U, V binary relations on A satisfying (P1)–(P3'). Then U, V satisfy also (P4) and (P5).

Proof. Let $a = a_1 U a_2 U \dots U a_n = a'$, $a = a'_1 V a'_2 V \dots V a'_m = a'$. Take a lower bound x and an upper bound y of the set $\{a_1, \dots, a_n, a'_1, \dots, a'_m\}$ and elements p, q as in (P2). Using 2.9 we get $\inf\{p, a\} = \inf\{p, a_1\} = \inf\{p, a_2\} = \dots = \inf\{p, a'\}$ and analogously $\inf\{a, q\} = \inf\{a', q\}$. But then $a = a'$ by 2.8.

Further let $a, a' \in A$. Take a lower bound x and an upper bound y of $\{a, a'\}$ and p, q as in (P2). Then $p_1 = \inf\{p, a\}$, $q'_1 = \inf\{a', q\}$ satisfy pVp_1Ua , $a'Vq'_1Uq$, $p_1VxUq'_1$. (P3) ensures the existence of $t = \sup\{p_1, q'_1\}$ with q'_1VtUp_1 . Hence $aUp_1UtVq'_1Va'$, completing the proof. \square

Using 1.2 we immediately get:

4.2. Theorem. *Let \mathbb{A} be a directed partially ordered set. If \mathbb{B} is a partially ordered set with $\text{Int } \mathbb{B}$ isomorphic to $\text{Int } \mathbb{A}$, then there exist partially ordered sets \mathbb{C} , \mathbb{D} such that \mathbb{A} is isomorphic to $\mathbb{C} \times \mathbb{D}$ and \mathbb{B} is isomorphic to $\mathbb{C}^\delta \times \mathbb{D}$.*

The converse is evident, so we have:

4.3. Corollary. *Let \mathbb{A} be a directed partially ordered set, \mathbb{B} any partially ordered set. The following conditions are equivalent:*

- (i) *$\text{Int } \mathbb{B}$ is isomorphic to $\text{Int } \mathbb{A}$,*
- (ii) *there exist partially ordered sets \mathbb{C} , \mathbb{D} such that \mathbb{A} is isomorphic to $\mathbb{C} \times \mathbb{D}$ and \mathbb{B} is isomorphic to $\mathbb{C}^\delta \times \mathbb{D}$.*

Since lattices are directed partially ordered sets, we obtain Theorem 1 of [11] as a consequence of 4.3. Let us notice that if \mathbb{A} is a lattice and \mathbb{B} is a partially ordered set with $\text{Int } \mathbb{B}$ isomorphic to $\text{Int } \mathbb{A}$, then \mathbb{B} is also a lattice, as 4.3 shows.




Fig. 5

Without the assumption that \mathbb{A} is directed, the assertion of 4.3 is false. To show this, consider \mathbb{A} as in Fig. 5. Let U and V be the relations marked out by the full and dashed lines, respectively. It is easy to see that U , V fulfil the conditions (P1)–(P3') and (P5), but (P4) is not satisfied (e.g. e_0Ve_1 and simultaneously $e_0Ud_0Ua_0Ub_0Uf_1Ue_1$ holds). Taking the corresponding \mathbb{A}_1 (in the sense of 2.10) as \mathbb{B} , depicted in Fig. 6, it fulfils (i) of 4.3, while it fails to satisfy (ii) of 4.3.




Fig. 6

References

- [1] V. I. Igošin: Selfduality of lattices of intervals of finite lattices. Inst. Matem. Sibir. Otdel. AN SSSR, Meždunarodnaja konferencija po algebre posvjaščennaja pamjati A. I. Maļceva. Tezisy dokladov po teoriji modelej i algebraičeskich sistem, Novosibirsk 1989, p. 48. (In Russian.)
- [2] V. I. Igošin: Lattices of intervals and lattices of convex sublattices of lattices. Uporjadočennyje množstva i rešetki. Saratov 6 (1990), 69–76. (In Russian.)
- [3] V. I. Igošin: Identities in interval lattices of lattices. Coll. Math. Soc. J. Bolyai 33 (Contributions to Lattice Theory), Szeged 1980 (1983), 491–501.
- [4] V. I. Igošin: On lattices with restriction on their intervals. Coll. Math. Soc. J. Bolyai 43 (Lectures in Universal Algebra), Szeged 1983 (1986), 209–216.
- [5] V. I. Igošin: Algebraic characteristic of lattices of intervals. Uspechi matem. nauk 40 (1985), 205–206. (In Russian.)
- [6] V. I. Igošin: Semimodularity in lattices of intervals. Math. Slovaca 38 (1988), 305–308.
- [7] J. Jakubík: Selfduality of the system of intervals of a partially ordered set. Czechoslov. Math. J. 41 (1991), 135–140.
- [8] J. Jakubík, J. Lihová: Systems of intervals of partially ordered sets. Math. Slovaca, to appear.
- [9] M. Kolibiar: Intervals, convex sublattices and subdirect representations of lattices. Universal Algebra and Applications, Banach Center Publications, Vol. 9, Warsaw 1982, 335–339.
- [10] J. Lihová: Posets having a selfdual interval poset. Czechoslovak Math. J. 44 (1994), 523–533.
- [11] V. Slavík: On lattices with isomorphic interval lattices. Czechoslovak Math. J. 35 (1985), 550–554.

Author's address: Prírodovedecká fakulta UPJŠ, 041 54 Košice, Jesenná 5, Slovakia.