

Emília Halušková

Direct limits of monounary algebras

Czechoslovak Mathematical Journal, Vol. 49 (1999), No. 3, 645–656

Persistent URL: <http://dml.cz/dmlcz/127516>

Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

DIRECT LIMITS OF MONOUNARY ALGEBRAS

EMÍLIA HALUŠKOVÁ, Košice¹

(Received December 10, 1996)

1. INTRODUCTION

The direct limit construction is a well-known method for building up algebras from families of algebras, e.g. [2], §21.

In this paper we investigate direct limits of monounary algebras.

Several examples of direct limit classes of monounary algebras will be given. We will describe all monounary algebras A which satisfy the following condition:

(C) If an algebra B can be obtained as a direct limit of algebras which are isomorphic to A , then B is isomorphic to A .

Further, we will show that every direct limit class of monounary algebras contains at least one algebra A which satisfies the condition (C).

2. PRELIMINARIES

As usual, by a monounary algebra we understand an algebra with a single unary operation; cf. e.g. [8], [9]. The notion of homomorphism is essentially applied in the construction of direct limits. Homomorphisms and endomorphisms of monounary algebras were thoroughly studied in [4], [7]–[9].

The class of all monounary algebras will be denoted by \mathcal{U} . We will use the symbol f for the operation in algebras of \mathcal{U} .

Let A be a monounary algebra.

The algebra A is said to be connected, if for each $x, y \in A$ there are positive integers m, n with $f^m(x) = f^n(y)$. A maximal connected subalgebra of A is said to be a component of A .

The class of all connected monounary algebras will be denoted by the symbol \mathcal{U}^c .

¹ Supported by grant GA SAV 1230/96.

Let \mathbb{N} be the set of all positive integers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

Let us denote by C_0 the connected monounary algebra which has \aleph_0 elements and a bijective operation. If $k \in \mathcal{N}$, then the connected monounary algebra which has k elements and f is a bijective operation will be denoted by C_k . We will say that an algebra A is a cycle of length k , if A is isomorphic to C_k . The class of all connected monounary algebras having a cycle of length k as its subalgebra will be denoted by \mathcal{U}_k^c . For the notation of the class of all connected monounary algebras without a cycle we will use the symbol \mathcal{U}_0^c , more precisely we put

$$\mathcal{U}_0^c = \mathcal{U}^c - \bigcup_{k \in \mathbb{N}} \mathcal{U}_k^c.$$

Let $A \in \mathcal{U}$. We will say that A has a cycle, if there exists $k \in \mathbb{N}$ such that a cycle of length k is a subalgebra of A .

These definitions immediately imply the following three lemmas:

Lemma 1. *Let $k \in \mathbb{N}$. If $x \in C_k$, then $f^k(x) = x$ and $|\{x, f(x), \dots, f^{k-1}(x)\}| = k$.*

Lemma 2. *Let $A \in \mathcal{U}$ and $i, j \in \mathbb{N}$. Let $u \in A$. If $f^i(u) = u$ and $v = f^j(u)$, then $f^i(v) = v$.*

Lemma 3. *Let $A \in \mathcal{U}$. If there exist $n \in \mathbb{N}$ and $x \in A$ such that $f^n(x) = x$, then A has a cycle.*

Lemma 4. *Let A, B be monounary algebras, φ a homomorphism from A into B and $k \in \mathbb{N}$. If A has a cycle C of length k , then there exists $l \in \mathbb{N}$ such that $\varphi(C)$ is a cycle of length l and l divides k .*

Proof. Let $x \in C$. Then $f^k(\varphi(x)) = \varphi(f^k(x)) = \varphi(x)$. Therefore there exists $l \in \mathbb{N}$ such that l divides k and $\{\varphi(x), f(\varphi(x)), \dots, f^{l-1}(\varphi(x))\}$ is a cycle of length l . Further, $\varphi(C) = \{\varphi(x), f(\varphi(x)), \dots, f^{k-1}(\varphi(x))\}$. \square

We recall the notion of the direct limit: in fact, we apply it to the case of monounary algebras.

Let $\langle P, \leq \rangle$ be a directed partially ordered set, $P \neq \emptyset$. For each $p \in P$ let A_p be a monounary algebra and assume that if $p, q \in P$, $p \neq q$, then $A_p \cap A_q = \emptyset$. Suppose that for each pair of elements p and q in P with $p < q$, a homomorphism φ_{pq} of A_p into A_q is defined such that $p < q < s$ implies that

$$\varphi_{ps} = \varphi_{pq} \circ \varphi_{qs}.$$

For each $p \in P$ let φ_{pp} be the identity on A_p . Then $\{A_p\}_{p \in P}$ is said to be a direct family of monounary algebras.

Let p and q be elements of P and let $x \in A_p$, $y \in A_q$. We put $x \equiv y$ if there exists $s \in P$ with $p \leq s$, $q \leq s$ such that $\varphi_{ps}(x) = \varphi_{qs}(y)$. For each $z \in \bigcup_{p \in P} A_p$ put $\bar{z} = \{t \in \bigcup_{p \in P} A_p : z \equiv t\}$. Denote $\bar{A} = \{\bar{z} : z \in \bigcup_{p \in P} A_p\}$.

If z_1, z_2 are elements of $\bigcup_{p \in P} A_p$ such that $\bar{z}_1 = \bar{z}_2$, then clearly $\overline{f(z_1)} = \overline{f(z_2)}$. Hence if we put $f(\bar{z}_1) = \overline{f(z_1)}$, then the operation f on \bar{A} is correctly defined and with respect to this operation \bar{A} is a monounary algebra. It is said to be the direct limit of the direct family $\{A_p\}_{p \in P}$. We express this situation by writing

$$(1) \quad \{A_p\}_{p \in P} \longrightarrow \bar{A}.$$

The definition of the direct limit yields the following four assertions.

Lemma 5. *Let (1) hold. Let $p \in P$ and φ_p be the mapping of A_p into \bar{A} such that $\varphi_p(x) = \bar{x}$ for every $x \in A_p$. Then φ_p is a homomorphism of A_p into \bar{A} .*

Lemma 6. *Let $m \in \mathbb{N}$ and let (1) be valid. If $|A_p| \leq m$ for every $p \in P$, then $|\bar{A}| \leq m$.*

Lemma 7. *Let (1) be valid. If the operation of A_p is injective for every $p \in P$, then the operation of \bar{A} is injective.*

Lemma 8. *Let (1) be valid and let $p \in P$. If $q \leq p$ for all $q \in P$, then $\bar{A} \cong A_p$.*

Lemma 9. *Let A be an algebra and let (1) be valid. If $A_p \cong A$ for all $p \in P$ and φ_{pq} is an isomorphism between A_p and A_q for all $p, q \in P$, $p \leq q$, then $\bar{A} \cong A$.*

It is obvious that Lemmas 5, 6, 8, 9 are not specific for monounary algebras, they are valid for direct limits of arbitrary type of algebraic systems.

Example. Suppose that P is the set of all finite subsets of the interval $(0, 1)$. Let $\leq = \subseteq$. For $p \in P$, $p = \{p_1, \dots, p_n\}$, where $n \in \mathbb{N}$, put $A_p = \{(0, p), (p_1, p), \dots, (p_n, p)\}$. Further, put $f((0, p)) = f((p_i, p)) = (0, p)$ for $i = 1, \dots, n$. If $p \subseteq q$, then let $\varphi_{pq}((z, p)) = (z, q)$ for every $z \in \{0, p_1, \dots, p_n\}$. The family $\{A_p\}_{p \in P}$ is direct and its direct limit is isomorphic to the algebra $(\langle 0, 1 \rangle, f)$, where $f(x) = 0$ for each $x \in \langle 0, 1 \rangle$.

This example shows that Lemma 6 cannot be generalized to the case when an infinite cardinal number will be put instead of m .

Lemma 10. *Let (1) be valid. The direct family $\{A_p\}_{p \in P}$ contains an algebra with a cycle if and only if \overline{A} has a cycle. More precisely, \overline{A} contains a cycle of length k , where k is the length of the shortest cycle in algebras of $\{A_p\}_{p \in P}$.*

Proof. Assume that $\{A_p\}_{p \in P}$ contains an algebra with a cycle. Let l be the length of the shortest cycle in algebras of $\{A_p\}_{p \in P}$. Every algebra of $\{A_p\}_{p \in P}$ can be homomorphically embedded into \overline{A} according to Lemma 5. This implies that \overline{A} is an algebra with a cycle. Moreover, \overline{A} has at least one cycle with length less or equal to l .

Now let \overline{A} have a cycle of length n , $n \in \mathbb{N}$. Assume that $p \in P$ and $x \in A_p$ are such that $f^n(\overline{x}) = \overline{x}$. We have $f^n(x) \in \overline{x}$ because $f^n(\overline{x}) = \overline{f^n(x)}$. This means that there exists $q \in P$ such that $\varphi_{pq}(x) = \varphi_{pq}(f^n(x))$. We obtain $\varphi_{pq}(x) = f^n(\varphi_{pq}(x))$. Thus A_q has a cycle with length less or equal to n . \square

Lemma 11. *Suppose that (1) is valid. Let \overline{A} have no cycle and let the direct family $\{A_p\}_{p \in P}$ contain an algebra with a subalgebra isomorphic to C_0 . Then \overline{A} has a subalgebra isomorphic to C_0 .*

Proof. Let $p \in P$ be such that C is a subalgebra of A_p isomorphic to C_0 . Consider the homomorphism φ_p from Lemma 5. Then $\varphi_p(C)$ is a homomorphic image of C and $\varphi_p(C)$ is a subalgebra of \overline{A} . The algebra \overline{A} has no cycle by the assumption and thus $\varphi_p(C) \cong C_0$. \square

3. DIRECT LIMIT CLASSES

The operator $\underline{\mathbf{L}}$ on classes of algebras was introduced in the textbook [2], §23. By this definition, if \mathcal{K} is a class of algebras, then $\underline{\mathbf{L}}(\mathcal{K})$ is the class of all direct limits of algebras of \mathcal{K} .

Let \mathcal{K} be a class of algebras. We denote by $[\mathcal{K}]$ the class of all isomorphic copies of algebras of \mathcal{K} . Further, we denote by $\underline{\mathbf{L}}'(\mathcal{K})$ the class of all isomorphic copies of direct limits of algebras of \mathcal{K} , i.e., $\underline{\mathbf{L}}'(\mathcal{K}) = [\underline{\mathbf{L}}(\mathcal{K})]$.

We will use $\underline{\mathbf{L}}'\mathcal{K}$ instead of $\underline{\mathbf{L}}'(\mathcal{K})$. For an algebra A we will use $[A]$ instead of $\{[A]\}$.

Lemma 12. *Let \mathcal{K} be a class of algebras. Then $\mathcal{K} \subseteq \underline{\mathbf{L}}'\mathcal{K}$.*

Proof. It follows from Lemma 9. \square

Lemma 13. Let $\mathcal{K}_1, \mathcal{K}_2$ be classes of algebras. If $\mathcal{K}_1 \subseteq \mathcal{K}_2$, then $\underline{\mathbf{L}}'[\mathcal{K}_1] \subseteq \underline{\mathbf{L}}'[\mathcal{K}_2]$.

Proof. Let $A \in \underline{\mathbf{L}}'[\mathcal{K}_1]$. Then there exists a direct family $\{A_p\}_{p \in P}$ such that $A_p \in [\mathcal{K}_1]$ for every $p \in P$ and $\{A_p\}_{p \in P} \longrightarrow A$. Since $A_p \in [\mathcal{K}_2]$ for every $p \in P$, we have $A \in \underline{\mathbf{L}}'[\mathcal{K}_2]$. \square

Definition. Let \mathcal{K} be a class of algebras. If $\underline{\mathbf{L}}'[\mathcal{K}] = [\mathcal{K}]$ is satisfied, then we will say that \mathcal{K} is a direct limit class.

The next lemma we will often use without any notice.

Lemma 14. A class \mathcal{K} is a direct limit class if and only if the following condition is valid:

whenever (1) holds and $A_p \in [\mathcal{K}]$ for each $p \in P$, then $\overline{A} \in [\mathcal{K}]$.

Proof. It follows from definitions and Lemma 12. \square

Lemma 15. a) Let J be a nonempty set and for $j \in J$ let \mathcal{K}_j be a direct limit class. Then $\bigcap_{j \in J} \mathcal{K}_j$ is a direct limit class.

b) If \mathcal{K}_1 and \mathcal{K}_2 are direct limit classes, then $\mathcal{K}_1 \cup \mathcal{K}_2$ is a direct limit class.

Proof. The assertion a) follows from definitions.

b) Suppose that (1) is valid and $A_p \in [\mathcal{K}_1 \cup \mathcal{K}_2]$ for all $p \in P$. Denote $Q = \{q \in P : A_q \in [\mathcal{K}_2]\}$.

Let there exist $p \in P$ such that for every $q \in Q$ the relation $p \not\leq q$ holds. Put $R = \{r \in P : p \leq r\}$. The set R is directed. Further, if $r \in R$, then $A_r \in [\mathcal{K}_1]$. If $s \in P$, then we can choose $s' \in P$ such that $s \leq s', p \leq s'$. We have $s' \in R$. This means that R is cofinal with P . Thus $\{A_r\}_{r \in R} \longrightarrow \overline{A}$ and $\overline{A} \in [\mathcal{K}_1]$.

Now for every $p \in P$ let there exists $q \in Q$ such that $p \leq q$. Then Q is cofinal with P and $\{A_q\}_{q \in Q} \longrightarrow \overline{A}$. Since \mathcal{K}_2 is a direct limit class, we obtain $\overline{A} \in [\mathcal{K}_2]$. \square

Direct limit classes of cyclically ordered groups have been dealt with by J. Jakubík and G. Pringerová, [3].

Example. Let O_ω be a monounary algebra such that \mathbb{N}_0 is the underlying set of O_ω and $f(x) = 0$ for all $x \in \mathbb{N}_0$. Let $k \in \mathbb{N}$. Let $O_k = \{0, 1, \dots, k\}$ and $f(x) = 0$ for all $x \in \{0, 1, \dots, k\}$. Put $\mathcal{K}_k = \{C_1, O_1, \dots, O_k\}$.

Assume that (1) is valid and $A_p \in [\mathcal{K}_k]$ for all $p \in P$. Let $o_p = f(x)$ for every $p \in P$ and $x \in A_p$. We have $|\overline{A}| \leq k + 1$ according to Lemma 6. Suppose that $p, q \in P$. Then there is $s \in P$ such that $p, q \leq s$. Since $\varphi_{ps}(o_p) = o_s = \varphi_{qs}(o_q)$, we obtain $\overline{o_p} = \overline{o_q}$. Further, $f(\overline{x}) = \overline{f(x)} = \overline{o_p} = \overline{o_q} = \overline{f(y)} = f(\overline{y})$ for every $x \in A_p$ and $y \in A_q$. We conclude $\overline{A} \in [\mathcal{K}_k]$ and \mathcal{K}_k is a direct limit class.

Consider $\mathcal{K} = \bigcup_{k \in \mathbb{N}} \mathcal{K}_k$. Then $\mathcal{K} = \{C_1\} \cup \{O_i, i \in \mathbb{N}\}$. For every $i \in \mathbb{N}$ let E_i be the trivial monounary algebra on the set $\{i\}$ and let $A_i = O_i \times E_i$. Let $\varphi_{i,i+1}$ be an embedding of A_i into A_{i+1} . Then $\{A_i\}_{i \in \mathbb{N}}$ is a direct family which has the direct limit isomorphic to O_ω . Since $O_\omega \notin [\mathcal{K}]$, we have \mathcal{K} is not a direct limit class.

This example shows that the union of direct limit classes need not be a direct limit class. The following lemma and Proposition 4 give some sufficient conditions which yield that the union of direct limit classes is a direct limit class.

Lemma 16. *Let $\mathcal{K}_k \subseteq \mathcal{U}_k^c$ be a direct limit class for all $k \in \mathbb{N}_0$. Then $\bigcup_{k \in \mathbb{N}_0} \mathcal{K}_k$ is a direct limit class.*

Proof. Let $\mathcal{K} = \bigcup_{k \in \mathbb{N}_0} \mathcal{K}_k$. Suppose that $\mathcal{K} \neq \emptyset$. Let (1) be valid, $A_p \in [\mathcal{K}]$ for all $p \in P$.

Assume that $\{A_p\}_{p \in P}$ contains an algebra with a cycle. Let i be the length of the shortest cycle in the algebras of $\{A_p\}_{p \in P}$. Put $Q = \{q \in P : A_q \in \mathcal{U}_i^c\}$. A homomorphic image of a cycle of length i can be only a cycle of length less or equal than i , in our case thus a cycle of length i . If $q \in Q$ and $p \in P$ are such that $q \leq p$, then $p \in Q$. Thus Q is directed and cofinal with P . We have $\{A_q\}_{q \in Q} \longrightarrow \overline{A}$ and $\overline{A} \in \mathcal{U}_i^c$ by Lemma 10. Because \mathcal{K}_i is a direct limit class, we have $\overline{A} \in [\mathcal{K}_i]$ and $\overline{A} \in [\mathcal{K}]$.

Now assume that $\{A_p\}_{p \in P}$ contains no algebra with a cycle. Then $A_p \in [\mathcal{K}_0]$ for every $p \in P$. Since \mathcal{K}_0 is a direct limit class, we obtain $\overline{A} \in [\mathcal{K}_0] \subseteq [\mathcal{K}]$. \square

Proposition 1. *The classes \mathcal{U} , \mathcal{U}^c , \mathcal{U}_k^c and $\{C_k\}$ are direct limit classes for every $k \in \mathbb{N}_0$.*

Proof. It is obvious that \mathcal{U} is a direct limit class.

Let (1) be valid.

a) We will prove that \mathcal{U}^c is a direct limit class. Suppose that $A_p \in \mathcal{U}^c$ for all $p \in P$. Let $\overline{x}, \overline{y} \in \overline{A}$. There exist $p \in P$ and $x_1 \in \overline{x}$, $y_1 \in \overline{y}$ such that $x_1, y_1 \in A_p$. We can find $m, n \in \mathbb{N}_0$ such that $f^m(x_1) = f^n(y_1)$ by the connectivity of A_p . This means that $f^m(\overline{x}) = \overline{f^m(x_1)} = \overline{f^n(y_1)} = f^n(\overline{y})$. We obtain $\overline{A} \in \mathcal{U}^c$ and \mathcal{U}^c is a direct limit class by Lemma 14 .

b) Suppose that $k \neq 0$. Let $A_p \in \mathcal{U}_k^c$ for all $p \in P$. We have $\overline{A} \in \mathcal{U}^c$ according to a) and \overline{A} has a cycle of length k by Lemma 10. So, \mathcal{U}_k^c is a direct limit class.

Now assume that $A_p \in \mathcal{U}_0^c$ for all $p \in P$. We have $\overline{A} \in \mathcal{U}_0^c$ according to a) and Lemma 10. Therefore \mathcal{U}_0^c is a direct limit class.

c) Let $k \neq 0$ and let $A_p \cong C_k$ for all $p \in P$. The algebra \overline{A} is connected by a) and \overline{A} contains a cycle of length k according to Lemma 10. The operation f of \overline{A} is

injective according to Lemma 7. We have $\overline{A} \cong C_k$. Conclude $\{C_k\}$ is a direct limit class.

Let $A_p \cong C_0$ for all $p \in P$. The algebra \overline{A} is connected by a), \overline{A} has no cycle by Lemma 10 and \overline{A} possesses a subalgebra isomorphic to C_0 by Lemma 11. In view of Lemma 7 we have $\overline{A} \cong C_0$. \square

Let A be a monounary algebra. Let A satisfy the following condition: If $C \subseteq A$ and C is a cycle of A , then $C \cong C_1$. Then A is called a cycle-free algebra. Cycle-free algebras have been dealt with by G. Bordalo [1].

Proposition 2. *The class of all cycle-free monounary algebras is a direct limit class.*

Proof. It follows from Lemma 10. \square

4. ALGEBRAS OF TYPE τ

Let A be a monounary algebra and let $\{A_j\}_{j \in J}$ be a component partition of A . We will say that A is of type τ if the following two conditions are valid:

1. If $j \in J$, then there exists $k \in \mathbb{N}$ such that $A_j \cong C_k$;
2. if $i, j \in J$, $i \neq j$ and $k, l \in \mathbb{N}$ are such that $A_i \cong C_k$, $A_j \cong C_l$, then k does not divide l .

Denote by \mathcal{T} the class of all algebras of type τ .

We will prove that \mathcal{T} is a direct limit class, and some special subclasses of \mathcal{T} are direct limit classes.

The definition of algebras of type τ yields that $C_k \in \mathcal{T}$ for every $k \in \mathbb{N}$. Further, if A is of type τ and C_1 is a subalgebra of A , then $A \cong C_1$. Further, if $A \in \mathcal{T}$ and B is a subalgebra of A , then $B \in \mathcal{T}$.

Lemma 17. *If $A \in \mathcal{T}$, then the set $\{A\}$ is a direct limit class.*

Proof. Suppose that (1) is valid and $A_p \cong A$ for each $p \in P$. Let $p, q \in P$. The algebra A is of type τ and thus φ_{pq} is an isomorphism between A_p and A_q in view of Lemma 4. This implies $\overline{A} \cong A$ according to Lemma 9. \square

Lemma 18. *Let (1) be valid and let $k \in \mathbb{N}$. If \overline{A} contains a cycle of length k , then there exists $p \in P$ such that A_q contains a cycle of length k for each $q \in P$ with $p \leq q$.*

P r o o f. We prove this assertion indirectly. Suppose that for each $p \in P$ there exists $q \in P, p \leq q$ such that A_q does not contain a cycle of length k .

We will show that for every $\bar{x} \in \bar{A}$ either $f^k(\bar{x}) \neq \bar{x}$ or

$$|\{\bar{x}, f(\bar{x}), \dots, f^{k-1}(\bar{x})\}| < k.$$

Then \bar{A} does not contain a cycle of length k by virtue of Lemma 1.

Assume that $\bar{x} \in \bar{A}$ and $f^k(\bar{x}) = \bar{x}$. Let $p \in P$ be such that $x \in A_p$. In view of the relation $\overline{f^k(x)} = \bar{x}$, there exists $q \in P, p \leq q$ such that $\varphi_{pq}(x) = \varphi_{pq}(f^k(x))$. We obtain $f^k(\varphi_{pq}(x)) = \varphi_{pq}(x)$. Thus A_q has a cycle of length m , where $m \leq k$.

Let $m < k$. The equality

$$\{\bar{x}, f(\bar{x}), \dots, f^{k-1}(\bar{x})\} = \{\overline{\varphi_{pq}(x)}, f(\overline{\varphi_{pq}(x)}), \dots, f^{k-1}(\overline{\varphi_{pq}(x)})\}$$

is valid. Therefore $|\{\bar{x}, f(\bar{x}), \dots, f^{k-1}(\bar{x})\}| < k$.

Let $m = k$. Choose $s \in P, q \leq s$ such that A_s does not contain a cycle of length k . Then the element $\varphi_{qs}(\varphi_{pq}(x))$ belongs to a cycle of A_s which has length $n, n < k$. Analogously as in the previous case we obtain $|\{\bar{x}, f(\bar{x}), \dots, f^{k-1}(\bar{x})\}| \leq n < k$. \square

Proposition 3. *The class \mathcal{T} is a direct limit class.*

P r o o f. Let (1) be valid and $A_p \in \mathcal{T}$ for all $p \in P$. According to Lemma 7 and Lemma 11, every component of \bar{A} is isomorphic to C_k for some $k \in \mathbb{N}$.

Assume that \bar{B}, \bar{C} are components of \bar{A} such that $\bar{B} \cong C_k, \bar{C} \cong C_l, k, l \in \mathbb{N}$. In view of Lemma 18 there exist $p, r \in P$ such that for each $q \in P, p \leq q$ the algebra A_q contains a cycle of length k and for each $s \in P, r \leq s$ the algebra A_s contains a cycle of length l . Choose $t \in P$ such that $r \leq t$ and $p \leq t$. We obtain that A_t has cycles of lengths k, l . The algebra A_t is of type τ . We get that if $k \neq l$, then k does not divide l . If $k = l$, then $\bar{B} = \bar{C}$. \square

Proposition 4. *Let $\mathcal{K} \subseteq \mathcal{T}$ and $n \in \mathbb{N}$. If every element of \mathcal{K} has less than n components, then \mathcal{K} is a direct limit class.*

P r o o f. Let (1) be valid and $A_p \in [\mathcal{K}]$ for all $p \in P$. We have $\bar{A} \in \mathcal{T}$ by the previous theorem.

Let $\{\bar{B}_i\}_{i \in I}$ be a component partition of \bar{A} . Put

$$m = \begin{cases} |I| & \text{if } I \text{ is finite,} \\ n & \text{otherwise.} \end{cases}$$

Let $i(1), \dots, i(m)$ be different elements of I .

Assume that $j \in \{1, \dots, m\}$ and $k(j) \in \mathbb{N}$ is such that $\overline{B}_{i(j)} \cong C_{k(j)}$. We use Lemma 18 and choose $p(j) \in P$ which has the following property: if $q \in P$ is such that $p(j) \leq q$, then the algebra A_q has a cycle of length $k(j)$.

Now let $s \in P$ be such that $p(1) \leq s, \dots, p(m) \leq s$. The algebra A_s contains cycles of lengths $k(1), \dots, k(m)$. Numbers $k(1), \dots, k(m)$ are different and thus \overline{A} is a subalgebra of A_s and $m < n$.

Assume that the algebra A_s has a component B such that $B \not\cong C_{k(j)}$ for $j = 1, \dots, m$. Then B is a cycle of length k and $k(j)$ does not divide k for $j = 1, \dots, m$. So, the algebra A_s cannot be homomorphically embedded into \overline{A} according to Lemma 4, which is a contradiction with Lemma 5. Thus $\overline{A} \cong A_s$ and $\overline{A} \in [\mathcal{K}]$. □

5. ONE-ELEMENT DIRECT LIMIT CLASSES

In this section we will describe all monounary algebras A such that $\mathbf{L}'[A] = [A]$; in this case we will speak about one-element direct limit class.

Let A be a monounary algebra.

The notion of degree $s(x)$ of an element $x \in A$ was introduced by M. Novotný [9] as follows. Let us denote by $A^{(\infty)}$ the set of all elements $x \in A$ such that there exists a sequence $\{x_n\}_{n \in \mathbb{N}_0}$ of elements belonging to A with the property $x_0 = x$ and $f(x_n) = x_{n-1}$ for each $n \in \mathbb{N}$. Further, we put

$$A^{(0)} = \{x \in A : f^{-1}(x) = \emptyset\}.$$

Now we define a set $A^{(\lambda)} \subseteq A$ for each ordinal λ by induction. Let λ be an ordinal, $\lambda \neq 0$. Assume that we have defined $A^{(\alpha)}$ for each ordinal $\alpha < \lambda$. Then we put

$$A^{(\lambda)} = \left\{ x \in A - \bigcup_{\alpha < \lambda} A^{(\alpha)} : f^{-1}(x) \subseteq \bigcup_{\alpha < \lambda} A^{(\alpha)} \right\}.$$

The sets $A^{(\lambda)}$ are pairwise disjoint. For each $x \in A$, either $x \in A^{(\infty)}$ or there is an ordinal λ with $x \in A^{(\lambda)}$. In the former case we put $s(x) = \infty$, in the latter we set $s(x) = \lambda$. We put $\lambda < \infty$ for each ordinal λ .

Let B be a subalgebra of A . Assume that there exists a homomorphism φ of A onto B such that $\varphi(b) = b$ for each $b \in B$. Then B is said to be a retract of A and φ is called a retract mapping corresponding to B .

Retracts of monounary algebras were thoroughly studied by D. Jakubíková-Studenovská [5], [6]. In view of [5], Theorem 1.3 we have

Lemma 19. Let $A \in \mathcal{U}$ and let B be a subalgebra of A . Then B is a retract of A if and only if the following conditions are satisfied:

- (a) If $y \in A$ is such that $f(y) \in B$, then there is $z \in B$ such that $s(y) \leq s(z)$ and $f(y) = f(z)$.
- (b) For any component D of A with $D \cap B = \emptyset$, the following conditions are satisfied:
 - (b1) If D contains a cycle with d elements, then there is a component D' of A with $D' \cap B \neq \emptyset$ and there is $n \in \mathbb{N}$ such that n divides d and D' has a cycle with n elements.
 - (b2) If D contains no cycle and $x \in D$, then there is $y \in B$ such that $s(f^k(x)) \leq s(f^k(y))$ for every $k \in \mathbb{N}_0$.

Lemma 20. Let $A \in \mathcal{U}$. If A contains a cycle, then there exists a retract T of A such that $T \in \mathcal{T}$.

Proof. Follows from the previous statement. □

Lemma 21. Let $A \in \mathcal{U}$ and let B be a retract of A . Then $B \in \underline{\mathbf{L}}'[A]$.

Proof. Let φ be a retract mapping corresponding to B . Let P be the set of all positive integers with the natural linear order. Assume that for each $p \in P$ there is an isomorphism ψ_p of A onto A_p . Put $\varphi_{pq}(\psi_p(a)) = \psi_q(\varphi(a))$ for all $a \in A$ and $p, q \in P$ such that $p < q$. Then $\{A_p\}_{p \in P}$ is a direct family and the direct limit of this family is an algebra isomorphic to B .

Corollary 5. Let \mathcal{K} be a direct limit class. Let $A \in \mathcal{K}$. If B is a retract of A , then $B \in [\mathcal{K}]$.

Proof. We have $\underline{\mathbf{L}}'[A] \subseteq \underline{\mathbf{L}}'[\mathcal{K}] = [\mathcal{K}]$. Thus Lemma 21 yields this assertion. □

Corollary 6. Let \mathcal{K} be a direct limit class of monounary algebras. If \mathcal{K} contains an algebra with a cycle, then \mathcal{K} contains an algebra of type τ .

Proof. The class $[\mathcal{K}]$ possesses an algebra of type τ according to Lemma 20 and Corollary 1. So, the claim follows from $[\mathcal{T}] = \mathcal{T}$. □

Lemma 22. Let $A \in \mathcal{U}$. Then there exists an algebra $B \in \underline{\mathbf{L}}'[A]$ such that each component of B is isomorphic to C_k for some $k \in \mathbb{N}_0$.

P r o o f. Let P be the set of all positive integers with the natural linear order. Assume that for each $p \in P$ there is an isomorphism ψ_p of A onto A_p . If $a \in A$, we will write $\psi_p(a) = a_p$. If $p \in P$, then we define $\varphi_{p,p+1}$ by setting

$$\varphi_{p,p+1}(a_p) = f(a_{p+1})$$

for each $a \in A$. So we have defined a direct family of monounary algebras. Let $\{A_p\}_{p \in P} \longrightarrow \overline{A}$. We will show that the operation f on \overline{A} is an injective and surjective mapping. Then the proof will be ready.

Assume that $u, v \in \overline{A}$ and $f(u) = f(v)$. Choose $p, q \in P$ and $a, b \in A$ such that $a_p \in u$, $b_q \in v$. Then $\overline{f(a_p)} = \overline{f(b_q)}$ and therefore there exists $s \in P$ such that $p \leq s$, $q \leq s$ and $\varphi_{ps}(f(a_p)) = \varphi_{qs}(f(b_q))$. Thus $f^{s+1-p}(a_s) = f^{s-p}(f(a_s)) = f^{s-q}(f(b_s)) = f^{s+1-q}(b_s)$. This yields $f^{s+1-p}(a) = f^{s+1-q}(b)$ because ψ_s is an isomorphism. We get $\varphi_{p,s+1}(a_p) = f^{s+1-p}(a_{s+1}) = f^{s+1-q}(b_{s+1}) = \varphi_{q,s+1}(b_q)$. This means $u = v$.

Further, let $p \in P$ and $a \in A$. Then $\overline{a_p} = \overline{\varphi_{p,p+1}(a_p)} = \overline{f(a_{p+1})} = f(\overline{a_{p+1}})$. □

Corollary 7. *Let \mathcal{K} be a direct limit class of monounary algebras. If $\mathcal{K} \neq \emptyset$, then there exists an algebra $B \in \mathcal{K}$ such that each component of B is isomorphic to C_k for some $k \in \mathbb{N}_0$.*

P r o o f. Lemma 22 yields this assertion. □

Theorem 1. *Let $A \in \mathcal{U}$. The following conditions are equivalent:*

- (i) $\{A\}$ is a direct limit class,
- (ii) $\underline{\mathbf{L}}'[A] = [A]$,
- (iii) either $A \cong C_0$ or $A \in \mathcal{T}$.

P r o o f. The equivalence (i) and (ii) follows from the definition. Now let (iii) hold. The set $\{C_0\}$ is a direct limit class in view of Proposition 1. If A is an algebra of type τ , then $\{A\}$ is a direct limit class by Lemma 17.

Conversely, assume that $A \not\cong C_0$ and A is not of type τ . Let \mathcal{K} be a direct limit class and $A \in \mathcal{K}$.

If A has a cycle, then \mathcal{K} contains an algebra of type τ by Corollary 2. Thus \mathcal{K} has more than one element.

If A has no cycle, then \mathcal{K} contains an algebra B such that each component of B is isomorphic to C_k for some $k \in \mathbb{N}_0$ according to Corollary 3. If $A \not\cong B$, then \mathcal{K} possesses more than one element. If $A \cong B$, then A is not connected and each component of A is isomorphic to C_0 . Thus there exists a retract C of A such that $C \cong C_0$ and $C \in [\mathcal{K}]$ by Corollary 1. Therefore \mathcal{K} has more than one element. □

Theorem 2. *Let \mathcal{K} be a direct limit class of monounary algebras. Then there exists an algebra $A \in \mathcal{K}$ such that $\{A\}$ is a direct limit class.*

P r o o f. We will prove that $\mathcal{K} \cap (\mathcal{T} \cup [C_0]) \neq \emptyset$.

If \mathcal{K} contains an algebra with a cycle, then $\mathcal{K} \cap \mathcal{T} \neq \emptyset$ according to Corollary 2.

Let \mathcal{K} contain no algebra with a cycle. Then Corollary 3 implies that \mathcal{K} contains an algebra B which has all components isomorphic to C_0 . Thus B has a retract C isomorphic to C_0 . We have $C \in [\mathcal{K}]$ by Corollary 1 and $C \in [C_0]$. Conclude $\mathcal{K} \cap [C_0] \neq \emptyset$. □

Acknowledgement. The author is indebted to the referee for his valuable remarks and suggestions.

References

- [1] *G. Bordalo*: A duality between unary algebras and their subuniverse lattices. Portugal Math. *46* (1989), 431–439.
- [2] *G. Grätzer*: Universal Algebra. Princeton, 1968.
- [3] *J. Jakubík, G. Pringerová*: Direct limits of cyclically ordered groups. Czechoslovak Math. J. *44* (1994), 231–250.
- [4] *D. Jakubíková-Studenovská*: Systems of unary algebras with common endomorphisms I, II. Czechoslovak Math. J. *29* (1979), 406–420, 421–429.
- [5] *D. Jakubíková-Studenovská*: Retract irreducibility of connected monounary algebras I, II. Czechoslovak Math. J. *46* (1996), 291–307; *47* (1997), 113–126.
- [6] *D. Jakubíková-Studenovská*: Two types of retract irreducibility of connected monounary algebras. Mathematica Bohemica *121* (1996), 143–150.
- [7] *O. Kopeček, M. Novotný*: On some invariants of unary algebras. Czechoslovak Math. J. *24* (1974), 219–246.
- [8] *E. Nelson*: Homomorphisms of monounary algebras. Pacific J. Math. *99* (1982(2)), 427–429.
- [9] *M. Novotný*: Über Abbildungen von Mengen. Pacific J. Math. *13* (1963), 1359–1369.

Author's address: Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia, e-mail: ehaluska@mail.saske.sk.