

Ariel Fernández; Miguel Florencio; J. Oliveros
Barrelledness of generalized sums of normed spaces

Czechoslovak Mathematical Journal, Vol. 50 (2000), No. 3, 459–465

Persistent URL: <http://dml.cz/dmlcz/127585>

Terms of use:

© Institute of Mathematics AS CR, 2000

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

BARRELLEDNESS OF GENERALIZED SUMS OF NORMED SPACES

A. FERNÁNDEZ, M. FLORENCIO and J. OLIVEROS, Sevilla

(Received January 13, 1997)

Abstract. Let $(E_i)_{i \in I}$ be a family of normed spaces and λ a space of scalar generalized sequences. The λ -sum of the family $(E_i)_{i \in I}$ of spaces is

$$\lambda\{(E_i)_{i \in I}\} := \{(x_i)_{i \in I}, x_i \in E_i, \text{ and } (\|x_i\|)_{i \in I} \in \lambda\}.$$

Starting from the topology on λ and the norm topology on each E_i , a natural topology on $\lambda\{(E_i)_{i \in I}\}$ can be defined. We give conditions for $\lambda\{(E_i)_{i \in I}\}$ to be quasi-barrelled, barrelled or locally complete.

Keywords: barrelled spaces, generalized sequences

MSC 2000: 46A08, 46A45, 46E40

1. INTRODUCTION AND PRELIMINARY RESULTS

The barrelledness, and related topics, of spaces of vector-valued sequences and functions have been studied in several papers [1]–[6], [8] and [10]. In particular, Florencio, Paúl and Sáez, extending the work of Lurje [10] where the barrelledness of $\ell^p\{E_n\}$ had been studied, characterized the barrelledness of the λ -sum of a sequence of normed spaces in [6]. More recently, Kakol and Roelcke in [8] have studied the barrelledness of ℓ^p -direct sums of a family of seminormed spaces for $1 \leq p \leq \infty$. Drewnowski, Florencio and Paúl have studied the barrelledness of bounded vector functions defined on a set with certain restrictions on its cardinal in [5].

In this paper we continue this investigations using techniques similar to those used in [1]–[6], to obtain more general results in the setting of the locally convex sum of a family of normed spaces.

This research has been partially supported by La Consejería de Educación y Ciencia de la Junta de Andalucía and the DGICYT project no. PB94-1460.

Let us recall at this point some definitions and notation. In what follows we will consider a fixed index set I and a space λ of scalar families, or generalized sequences, on I , i.e., a linear subspace of the space of all real or complex functions defined on I .

We say that λ is solid (see [9, §30]) if whenever it contains $\beta = (\beta_i)_i$ it also contains all families $\alpha = (\alpha_i)_i$ with $|\alpha_i| \leq |\beta_i|$ for all $i \in I$. The Köthe-dual λ^\times of the space λ is defined as for sequences spaces, i.e., λ^\times consists of all generalized sequences $(\eta_i)_i$ such that $\sum |\alpha_i \eta_i| < \infty$ for every $(\alpha_i)_i$ in λ . We always consider on λ a normal locally convex Hausdorff topology in the sense of Rosier (see [12]). Such a topology can be given by a system Q of seminorms with the following properties:

- (a) If $\alpha \leq \beta$ (i.e. $|\alpha_i| \leq |\beta_i|$ for all $i \in I$), then $q(\alpha) \leq q(\beta)$ for all $q \in Q$.
- (b) For every η in the Köthe-dual space λ^\times there exists a seminorm $q \in Q$ such that $|\langle \gamma, \eta \rangle| \leq q(\gamma)$ for all $\gamma \in \lambda$.

If $(E_i)_{i \in I}$ is a family of real or complex normed spaces, we define the λ -sum of $(E_i)_{i \in I}$ as

$$\lambda\{(E_i)_{i \in I}\} := \{(x_i)_{i \in I} : x_i \in E_i \text{ and } (\|x_i\|)_{i \in I} \in \lambda\}.$$

To ensure that $\lambda\{(E_i)\}$ is a linear space we must assume that λ be solid. Starting from the topology of λ and the norm topology on each E_i , we consider the locally convex topology on $\lambda\{(E_i)\}$ determined by the seminorms:

$$\sigma_q : x = (x_i)_i \in \lambda\{(E_i)\} \longrightarrow \sigma_q(x) := q((\|x_i\|)_i) \in \mathbb{R},$$

as q runs through Q .

In this paper we study the barrelledness of $\lambda\{(E_i)\}$. Recall that a locally convex space E is barrelled if and only if it is quasi-barrelled (every $\beta(E', E)$ -bounded set in the dual of E is equicontinuous) and has the Banach-Mackey property (every $\sigma(E', E)$ -bounded set in its dual is $\beta(E', E)$ -bounded). We refer the reader to the monographs [7], [9] and [11] for the terminology in local convexity and barrelledness used here.

We start by lifting the property of quasi-barrelledness from the space λ to the space $\lambda\{(E_i)\}$.

Following a way similar to the proofs of [4, Theorem 1] or [6, Theorem 4] we can show the following

Theorem 1. *If λ is quasi-barrelled, then $\lambda\{(E_i)\}$ is quasi-barrelled.*

The next step will be to analyze when the space $\lambda\{(E_i)\}$ has the Banach-Mackey property. We will do this in two cases.

First, in Section 2, when the space λ is defined on an index set I that has nonmeasurable cardinal. Recall that a set I has nonmeasurable cardinal if there exists no

countable additive measure $\mu: \mathcal{P}(I) \rightarrow \{0, 1\}$ such that $\mu(I) = 1$ and $\mu(\{i\}) = 0$ for all $i \in I$ [11, Def. 6.2.21]. As we shall see, this concept will be strongly connected to the assumption that the space $\lambda\{(E_i)\}$ is not barrelled.

Secondly, in Section 3, we will deal with a space λ which has the property of convergence of sections without any restrictions on the cardinal of I .

Before to do this we need to prove a result about the local completeness of the space $\lambda\{(E_i)\}$. Its proof is standard but we include it for the sake of convenience.

Theorem 2. *If λ is locally complete and every E_i is a Banach space, then $\lambda\{(E_i)\}$ is locally complete.*

P r o o f. Taking into account [11, Prop. 5.1.6 and Prop. 3.2.3] we shall show that if B is a closed disc in $\lambda\{(E_i)\}$ and $(x^{(n)})_n$ is a sequence of elements of B , then the series $\sum_{n=1}^{\infty} 2^{-n}x^{(n)}$ converges to an element of B .

Let $M = \{(\|z_i\|)_i: z = (z_i)_i \in B\}$. Note that M is bounded in λ . Since λ is locally complete it follows that $D = \overline{acx}(M)$ is a Banach disc. Now $((\|x_i^{(n)}\|)_i)_n \subset D$ so that the series $\sum_{n=1}^{\infty} 2^{-n}(\|x_i^{(n)}\|)_i$ converges in λ to an element, say $\alpha = (\alpha_i)_i$.

Coordinatewise we have that $\sum_{n=1}^{\infty} 2^{-n}\|x_i^{(n)}\| = \alpha_i$ for every $i \in I$.

From the boundedness of $(x_i^{(n)})_n$ and the completeness of each E_i it follows that $\sum_{n=1}^{\infty} 2^{-n}x_i^{(n)}$ converges to an element $x_i \in E_i$. Observe that $x = (x_i)_i$ is an element of $\lambda\{(E_i)\}$ since $\|x_i\| \leq \alpha_i$ for every $i \in I$.

We complete the proof by proving that x is the sum of the series $\sum_{n=1}^{\infty} 2^{-n}x^{(n)}$ and $x \in B$. Given an arbitrary seminorm q on λ we have

$$\begin{aligned} \sigma_q\left(x - \sum_{n=1}^k 2^{-n}x^{(n)}\right) &= q\left(\left(\|x_i - \sum_{n=1}^k 2^{-n}x_i^{(n)}\|_i\right)\right) \\ &\leq q\left(\sum_{n \geq k+1} 2^{-n}(\|x_i^{(n)}\|)_i\right) \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

since $\sum_{n=1}^{\infty} 2^{-n}(\|x_i\|)_i$ is convergent in λ . Finally, note that $x \in B$ because B is a closed disc. □

2. WHEN I HAS NONMEASURABLE CARDINAL

We start this section by studying when the space $\lambda\{(E_i)\}$ has the Banach-Mackey property. In the next theorem we will use the following notation about projections. If J is a subset of I and x is an element of $\lambda\{(E_i)\}$, then $P_J(x)$ is the generalized sequence that agrees with x on J and is null on $I \setminus J$.

Theorem 3. *If I has nonmeasurable cardinal, the space λ is locally complete and the normed spaces $(E_i)_i$ are all barrelled, then the space $\lambda\{(E_i)\}$ has the Banach-Mackey property.*

Proof. Suppose, on the contrary, that there exists G in the dual space $\lambda\{(E_i)\}'$ that is $\sigma(\lambda\{(E_i)\}', \lambda\{(E_i)\})$ -bounded but is not a bounded set in $\beta(\lambda\{(E_i)\}', \lambda\{(E_i)\})$. Then there exists a bounded set A in $\lambda\{(E_i)\}$ such that

$$\sup\{|\langle a, u \rangle| : u \in G, a \in A\} = +\infty.$$

The set $B := \bigcup_{J \subseteq I} P_J(A)$ is bounded because A is bounded and the set of projections $\{P_J : J \subseteq I\}$ is equicontinuous.

From the sets G and B let us consider the filter given on I by

$$\mathcal{F} = \{J \subseteq I : G^\circ \text{ absorbs } P_{I \setminus J}(B)\},$$

and let \mathcal{U} be the ultrafilter generated by the filter \mathcal{F} . Denote by μ the standard finitely additive measure on $\mathcal{P}(I)$ associated with the ultrafilter \mathcal{U} . Since I has nonmeasurable cardinal, we have that μ is noncountable additive, hence there exists a sequence $(J_k)_k$ of pairwise disjoint subsets of I with $\mu(J_k) = 0$ for all k and $\mu\left(\bigcup_{k=1}^{\infty} J_k\right) = 1$.

If we put $I_n = \bigcup_{k \geq n} J_k$ for all $n = 1, 2, \dots$, we have a decreasing sequence of subsets of I with empty intersection such that $\mu(I_n) = 1$ for all $n = 1, 2, \dots$ because

$$\begin{aligned} 1 &= \mu\left(\bigcup_{k=1}^{\infty} J_k\right) = \mu\left(J_1 \cup \dots \cup J_{n-1} \cup \bigcup_{k=n}^{\infty} J_k\right) \\ &= \mu(J_1) + \dots + \mu(J_{n-1}) + \mu\left(\bigcup_{k=n}^{\infty} J_k\right) \\ &= \mu(I_n). \end{aligned}$$

It follows that each I_n is in \mathcal{U} and therefore G° does not absorb $P_{I_n}(B)$, so there exist $z^{(n)} \in B$, supported in I_n ($P_{I_n}(z^{(n)}) = z^{(n)}$), and $u^{(n)} \in G$ such that

$$(1) \quad |\langle z^{(n)}, u^{(n)} \rangle| > n, \quad \text{for all } n = 1, 2, \dots$$

From the bounded sequence $(z^{(n)})_n$ we consider the set

$$D = \left\{ \sum_{n=1}^{\infty} \alpha_n z^{(n)} : (\alpha_n)_n \text{ in the unit ball of } \ell^1 \right\}.$$

Let us observe that for all $(\alpha_n)_n$ in ℓ^1 (the space of absolutely summable sequences) the series $\sum_{n=1}^{\infty} \alpha_n z^{(n)}$ converges in $\lambda\{(\widehat{E}_i)_{i \in I}\}$ to an element which we will denote by $z = (z_i)_{i \in I}$. This follows by applying Theorem 2 above to the space λ and the Banach spaces $(\widehat{E}_i)_i$. Moreover, as we shall see in a moment, each z_i is in E_i , so $\sum_{n=1}^{\infty} \alpha_n z^{(n)}$ really converges in $\lambda\{(E_i)\}$. Indeed, since $(I_n)_n$ is a decreasing sequence of subsets of I with empty intersection, for each $i \in I$ there are two possibilities:

- 1) If $i \notin I_1$, then $P_{\{i\}}(z^{(n)}) = 0$ for all $n = 1, 2, \dots$, so $z_i = 0$.
- 2) There exists a natural number p_i such that $i \in I_{p_i}$ but $i \notin I_k$ for all $k > p_i$. In this case we have

$$\begin{aligned} z_i &= P_{\{i\}}(z) = P_{\{i\}}\left(\sum_{n=1}^{\infty} \alpha_n z^{(n)}\right) = \sum_{n=1}^{\infty} \alpha_n P_{\{i\}}(z^{(n)}) \\ &= \sum_{n=1}^{p_i} \alpha_n P_{\{i\}}(z^{(n)}) \in E_i. \end{aligned}$$

Now, it follows from [1, Prop. p. 74] that D is a Banach disc in $\lambda\{(E_i)\}$. As barrels absorb every Banach disc [7, 8.3.3], we have that there exists a number $\rho > 0$ such that $D \subset \rho G^\circ$. On the other hand, from (1), $z^{(n)} \notin nG^\circ$ for all $n \geq 1$. This contradiction completes the proof of the theorem. \square

Our main result is the next

Theorem 4. *If λ is a locally complete and barrelled space of generalized sequences on a set I which has nonmeasurable cardinal, then $\lambda\{(E_i)\}$ is barrelled if and only if every E_i is barrelled.*

Proof. The direct implication follows from Theorems 1 and 3, and the inverse one follows from the fact that every E_i is complemented in $\lambda\{(E_i)\}$. \square

Remark 1. This theorem implies, as important particular cases, the barrelledness of spaces $\ell_I^p\{(E_i)_{i \in I}\}$ with $1 \leq p \leq \infty$, where I is a nonmeasurable set. Compare our Theorem 4 with the results of Kakol and Roelcke in [8] and Drewnowski, Florencio and Paúl in [5].

3. SPACES λ WITH THE PROPERTY OF CONVERGENCE OF SECTIONS

Let us introduce some more notation in order to establish the barrelledness of $\lambda\{(E_i)\}$ in the setting of a space λ with the property of convergence of sections. The sections of an element of λ are defined to be its projections over finite subsets of I . This property allows us to represent the dual of $\lambda\{(E_i)\}$ as the space $\lambda^\times\{(E'_i)_{i \in I}\}$. With similar arguments to those used in [6, Theorem 1] we can prove the equality

$$(2) \quad \lambda\{(E_i)_{i \in I}\}' = \lambda^\times\{(E'_i)_{i \in I}\} \\ = \left\{ (u_i)_{i \in I}, u_i \in E'_i, \sum_{i \in I} |\langle x_i, u_i \rangle| < \infty, \text{ for all } (x_i)_{i \in I} \in \lambda\{(E_i)\} \right\}.$$

Lemma 5. *If the space λ has the property of convergence of sections and $(E_i)_i$ is a family of normed spaces such that every E'_i is $\sigma(E'_i, E_i)$ -sequentially complete, then $\lambda\{(E_i)\}'$ is $\sigma(\lambda\{(E_i)\}', \lambda\{(E_i)\})$ -sequentially complete. In particular, the space $\lambda\{(E_i)\}$ has the Banach-Mackey property.*

Proof. According to (2) let $(u^{(n)})_n$ be a $\sigma(\lambda\{(E_i)\}', \lambda\{(E_i)\})$ -Cauchy sequence in $\lambda\{(E_i)\}'$. By using the natural inclusion of E_i in $\lambda\{(E_i)\}$ we see that for every $i \in I$ the sequence $(u_i^{(n)})_n$ is $\sigma(E'_i, E_i)$ -Cauchy in E'_i . Since every E'_i is $\sigma(E'_i, E_i)$ -sequentially complete, $(u_i^{(n)})_n$ is actually convergent to an element $u_i \in E'_i$. Put $u = (u_i)_{i \in I}$. We complete the proof by proving that u is in $\lambda\{(E_i)\}'$ and that $(u^{(n)})_n$ converges to u in the $\sigma(\lambda\{(E_i)\}', \lambda\{(E_i)\})$ -topology.

If $x = (x_i)_i$ is in $\lambda\{(E_i)\}$, then $\alpha^{(n)} = (\langle u_i^{(n)}, x_i \rangle)_i$ is in ℓ_I^1 for all $n \geq 1$. Furthermore, $(\alpha^{(n)})_n$ is a Cauchy sequence in $\sigma(\ell_I^1, \ell_I^\infty)$ since $(u^{(n)})_n$ is a Cauchy sequence in the topology $\sigma(\lambda\{(E_i)\}', \lambda\{(E_i)\})$ and $\lambda\{(E_i)\}$ is solid. Now the Schur lemma [7, §10.5 Cor. 4] ensures that $(\alpha^{(n)})_n$ is norm convergent to an element $\alpha = (\alpha_i)_i$ of ℓ_I^1 .

Coordinatewise we have that

$$\alpha_i = \lim_n \langle x_i, u_i^{(n)} \rangle = \langle x_i, u_i \rangle$$

for every $i \in I$. This yields that $(\langle x_i, u_i \rangle)_i$ is in ℓ_I^1 , since $u \in \lambda\{(E_i)\}'$.

Now

$$|\langle x, u^{(n)} - u \rangle| \leq \sum_{i \in I} |\langle x_i, u_i^{(n)} \rangle - \langle x_i, u_i \rangle| \\ = \sum_{i \in I} |\alpha_i^{(n)} - \alpha_i|$$

and taking into account that $\alpha^{(n)} \rightarrow \alpha$ in $(\ell_I^1, \|\cdot\|_1)$, it follows that $u = \lim_n u^{(n)}$ and the proof is complete. □

Theorem 6. *If λ has the property of convergence of sections and is barrelled, then $\lambda\{(E_i)\}$ is barrelled if and only if each E_i is barrelled.*

Proof. If every E_i is barrelled, then it follows from [9, §23.6 (4)] that E'_i is $\sigma(E'_i, E_i)$ -sequentially complete. The barrelledness of $\lambda\{(E_i)\}$ follows from Theorem 1 and Lemma 5. The inverse implication follows from the fact that every E_i is complemented in $\lambda\{(E_i)\}$. \square

Remark 2. As spaces ℓ^p_I ($1 \leq p < \infty$) and c_{0I} have the property of convergence of sections, the spaces $\ell^p_I\{(E_i)_{i \in I}\}$ and $c_{0I}\{(E_i)_{i \in I}\}$ are barrelled if each E_i is barrelled.

References

- [1] *J. C. Díaz, M. Florencio, P. J. Paúl:* A uniform boundedness theorem for $L^\infty(\mu, E)$. Arch. Math. (Basel) 60 (1993), 73–78.
- [2] *S. Díaz, A. Fernández, M. Florencio, P. J. Paúl:* An abstract Banach-Steinhaus theorem and applications to function spaces. Results Math. 23 (1993), 242–250.
- [3] *L. Drewnowski, M. Florencio, P. J. Paúl:* Barrelled subspaces of spaces with subseries decompositions or Boolean rings of projections. Glasgow Math. J. 36 (1994), 57–69.
- [4] *L. Drewnowski, M. Florencio, P. J. Paúl:* Barrelled function spaces. Progress in Functional Analysis (K.D. Bierstedt et al., eds.). North-Holland Math. Studies, Elsevier/North-Holland, Amsterdam, Oxford, New York and Tokyo, 1992, pp. 191–199.
- [5] *L. Drewnowski, M. Florencio, P. J. Paúl:* On the barrelledness of spaces of bounded vector functions. Arch. Math. (Basel) 63 (1994), 449–458.
- [6] *M. Florencio, P. J. Paúl, C. Sáez:* Barrelledness in λ -sums of normed spaces. Simon Stevin 63 (1989), 209–217.
- [7] *H. Jarchow:* Locally Convex Spaces. B.G. Teubner. Stuttgart, 1981.
- [8] *J. Kakol, W. Roelcke:* On the barrelledness of ℓ^p -direct sums of seminormed spaces for $1 \leq p \leq \infty$. Arch. Math. (Basel) 62 (1994), 331–334.
- [9] *G. Köthe:* Topological Vector Spaces I. Springer-Verlag, Berlin, Heidelberg and New York, 1969.
- [10] *P. Lurje:* Tonnelierheit in lokalkonvexen Vektorgruppen. Manuscripta Math. 14 (1974), 107–121.
- [11] *P. Pérez Carreras, J. Bonet:* Barrelled Locally Convex Spaces. North-Holland Math. Studies, North-Holland, Amsterdam, New York, Oxford and Tokyo, 1987.
- [12] *R. C. Rosier:* Dual spaces of certain vector sequence spaces. Pacific J. Math. 46 (1973), 487–501.

Authors' address: Dpto. Matemática Aplicada II, Escuela Superior Ingenieros, Camino de los Descubrimientos, s/n, 41092-Sevilla, Spain, e-mail: anfercar@matinc.us.es.