

Fernando Etayo; Mario Fioravanti; Ujué R. Trías

The dimension function of holomorphic spaces of a real submanifold of an almost complex manifold

Czechoslovak Mathematical Journal, Vol. 51 (2001), No. 1, 139–141

Persistent URL: <http://dml.cz/dmlcz/127633>

Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE DIMENSION FUNCTION OF HOLOMORPHIC SPACES OF A
REAL SUBMANIFOLD OF AN ALMOST COMPLEX MANIFOLD

FERNANDO ETAYO, MARIO FIORAVANTI and UJUÉ R. TRÍAS, Santander

(Received December 2, 1997)

Abstract. Let M be a real submanifold of an almost complex manifold $(\overline{M}, \overline{J})$ and let $H_x = T_x M \cap \overline{J}(T_x M)$ be the maximal holomorphic subspace, for each $x \in M$. We prove that $c: M \rightarrow \mathbb{N}$, $c(x) = \dim_{\mathbb{R}} H_x$ is upper-semicontinuous.

Keywords: holomorphic space, submanifold, almost complex

MSC 2000: primary 53C40, secondary 53C15

Let $(\overline{M}, \overline{J})$ be an almost complex manifold and let M be a real submanifold of \overline{M} . Then one can consider, $\forall x \in M$, the maximal holomorphic subspaces $H_x = T_x M \cap \overline{J}(T_x M)$ and define the dimension function $c: M \rightarrow \mathbb{N}$, $c(x) = \dim_{\mathbb{R}} H_x$. Submanifolds verifying that the function c is constant are called *generic submanifolds* (cf. [C]). As is well known, this condition is equivalent to the following: the maximal holomorphic subspaces H_x define a distribution over M . Many results have been found for generic submanifolds and particular cases, such as almost complex, totally real, Cauchy-Riemann, etc.

However, no global properties of this function have been proved. The purpose of this note is to prove that the function c is upper-semicontinuous, i.e., the sets $A_n = \{x \in M / c(x) < n\}$ are open subsets of M .

One can easily prove the following properties, for all $x \in M$: (a) $c(x)$ is an even number and (b) $2 \dim M - \dim \overline{M} \leq c(x) \leq \dim M$. In particular, if M is a real hypersurface of \overline{M} , then c is the constant function $c(x) = \dim M - 2$, $\forall x \in M$.

On the other hand, observe that $(M, \overline{J}|_M)$ is a complex submanifold of $(\overline{M}, \overline{J})$ iff $c(x) = \dim M$, $\forall x \in M$. (See [K-N, prop. IX.2.3].)

The work of the first author is partially supported by Spanish Grant DGICYT PB95-0124.

Let M be a real submanifold of an almost complex manifold $(\overline{M}, \overline{J})$. We shall use the following notation: $G_r(T_x M)$ (resp. $G_r(T_x \overline{M})$) is the Grassmann manifold of r -planes of $T_x M$ (resp. of $T_x \overline{M}$) and $\pi: G_r(M) \rightarrow M$ and $\overline{\pi}: \overline{G}_r(M) \rightarrow M$ denote the Grassmann bundles over M with fibres $\pi^{-1}(x) = G_r(T_x M)$ and $\overline{\pi}^{-1}(x) = G_r(T_x \overline{M})$, $\forall x \in M$. Observe that the almost complex structure \overline{J} on \overline{M} induces a continuous fibred automorphism $\overline{J}: \overline{G}_r(M) \rightarrow \overline{G}_r(M)$ which is involutive. i.e., $\overline{J} \circ \overline{J} = \text{id}$.

We can prove the following

Lemma. *Let M be a real submanifold of an almost complex manifold $(\overline{M}, \overline{J})$ and let $x \in M$ and $r \in \mathbb{N}$ be such that $c(x) < r \leq \dim M$. Then $G_r(T_x M) \cap \overline{J}(G_r(T_x M)) = \emptyset$.*

Proof. As $r > c(x)$, then $\overline{J}(W_r) \neq W_r, \forall W_r \in G_r(T_x M)$. Let us assume that there exists $W_r \in G_r(T_x M) \cap \overline{J}(G_r(T_x M))$. Then $W_r = \overline{J}(W'_r)$ with $W'_r \in G_r(T_x M)$, and one can easily prove that $W_r + W'_r$ is a \overline{J} -invariant subspace of $T_x M$ with $\dim(W_r + W'_r) \geq r > c(x)$, which is not possible, thus proving the lemma. \square

Then we have:

Theorem. *If M is a real submanifold of an almost complex manifold $(\overline{M}, \overline{J})$, then the function $c: M \rightarrow \mathbb{N}$, $c(x) = \dim_{\mathbb{R}} H_x$ is upper-semicontinuous.*

Proof. We shall prove that the sets $A_n = \{x \in M / c(x) < n\}$ are open subsets of M showing that $\forall x \in M$, there exists a neighborhood U of x in M such that $\forall y \in U$, $c(y) \leq c(x)$.

If $c(x) = \dim M$, the result is trivial. Let us assume that $c(x) < \dim M$. We shall prove that for every $r \in \mathbb{N}$ such that $c(x) < r \leq \dim M$, there exists a neighborhood U_r of x in M , verifying $c(y) < r, \forall y \in U_r$. Then the solution will be the neighborhood $U = \bigcap \{U_r, c(x) < r \leq \dim M\}$.

Let us consider $x \in M$ and $r \in \mathbb{N}$ such that $c(x) < r \leq \dim M$. Then, by the lemma, one has $G_r(T_x M) \cap \overline{J}(G_r(T_x M)) = \emptyset$. As $G_r(T_x M)$ is compact and $\overline{G}_r(M)$ is a Hausdorff space, there exist two neighborhoods \overline{V} of $G_r(T_x M)$ and \overline{W} of $\overline{J}(G_r(T_x M))$ such that $\overline{V} \cap \overline{W} = \emptyset$. Then $\overline{N} = \overline{V} \cap \overline{J}(\overline{W})$ is a neighborhood of $G_r(T_x M)$ verifying $\overline{N} \cap \overline{J}(\overline{N}) = \emptyset$.

Let us consider $N = \overline{N} \cap G_r(M)$. Then N is a neighborhood of $G_r(T_x M)$ in $G_r(M)$. One can consider that N is included in $\pi^{-1}(U')$, where U' is a trivialization neighborhood of x , and then there exists a diffeomorphism

$$\varphi: \pi^{-1}(U') \rightarrow U' \times G_r(\mathbb{R}^{\dim M})$$

given by $\varphi(W_r) = (\pi(W_r), \alpha(W_r))$.

For each $W_r \in G_r(T_x M)$ there exists a basic neighborhood $\varphi^{-1}(A^{W_r} \times B^{W_r})$, where A^{W_r} (resp. B^{W_r}) is a neighborhood of $\pi(W_r)$ (resp. $\alpha(W_r)$) and $\varphi^{-1}(A^{W_r} \times B^{W_r}) \subset N$, thus defining an open covering of $G_r(T_x M)$. Taking into account that $G_r(T_x M)$ is compact, one obtains a finite family $\{\varphi^{-1}(A^{W_r^i} \times B^{W_r^i}), 1 \leq i \leq k\}$ covering $G_r(T_x M)$.

Finally, let $U_r = \bigcap \{A^{W_r^i}, 1 \leq i \leq k\}$. Then $\pi^{-1}(U_r) \subset N$ and $\pi^{-1}(U_r) \cap \overline{J}(\pi^{-1}(U_r)) = \emptyset$, thus proving that $c(y) < r, \forall y \in U_r$. \square

Remarks. (1) Let M be a real compact submanifold of the complex euclidean space \mathbb{C}^m . As is well known (cf. [W, p. 11]), M cannot be immersed into \mathbb{C}^m as a holomorphic submanifold. Therefore, there exists a point $x \in M$ such that $c(x) < \dim M$. The above theorem shows that $A_{\dim M}$ is a non-empty open subset of M .

(2) The theorem is also true if one replaces the almost complex manifold $(\overline{M}, \overline{J})$ by an almost product manifold $(\overline{M}, \overline{J})$, i.e., if $\overline{J} \circ \overline{J} = \text{id}$.

References

- [C] *B. Y. Chen: Geometry of Submanifolds and its Applications.* Sci. Univ. Tokyo, 1981.
- [K-N] *S. Kobayashi and K. Nomizu: Foundations of Differential Geometry, II.* Interscience, New York, 1969.
- [W] *R. O. Wells, Jr.: Differential Analysis on Complex Manifolds.* Springer, New York, 1980.

Author's address: Departamento de Matemáticas, Estadística y Computación, Facultad de Ciencias, Universidad de Cantabria, 39071 Santander, Spain, e-mails: etayof@unican.es, fioravam@unican.es.