

María Silvina Riveros; A. de la Torre

On the best ranges for  $A_p^+$  and  $RH_r^+$

*Czechoslovak Mathematical Journal*, Vol. 51 (2001), No. 2, 285–301

Persistent URL: <http://dml.cz/dmlcz/127648>

## Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE BEST RANGES FOR  $A_p^+$  AND  $RH_r^+$ 

M. S. RIVEROS, Córdoba, and A. DE LA TORRE, Málaga

(Received May 28, 1998)

*Abstract.* In this paper we study the relationship between one-sided reverse Hölder classes  $RH_r^+$  and the  $A_p^+$  classes. We find the best possible range of  $RH_r^+$  to which an  $A_1^+$  weight belongs, in terms of the  $A_1^+$  constant. Conversely, we also find the best range of  $A_p^+$  to which a  $RH_\infty^+$  weight belongs, in terms of the  $RH_\infty^+$  constant. Similar problems for  $A_p^+$ ,  $1 < p < \infty$  and  $RH_r^+$ ,  $1 < r < \infty$  are solved using factorization.

*Keywords:* one-sided weights, one-sided reverse Hölder, factorization

*MSC 2000:* 42B25

## 1. INTRODUCTION

It is well known that there is a relationship between the  $A_p$  classes and the so called reverse Hölder classes  $RH_r$ . C. J. Neugebauer [8] studied the following problems:

- (1) For  $w \in A_p$ , find the precise range of  $r$ 's such that  $w \in RH_r$ , the precise range of  $q < p$  for which  $w \in A_q$ , and the precise range of  $s > 1$  such that  $w^s \in A_p$ .
- (2) Conversely, for a fixed  $w \in RH_r$ , find the precise range of  $p$ 's such that  $w \in A_p$ , and the precise range of  $q > r$  for which  $w \in RH_q$ .

For the one-sided Hardy-Littlewood maximal operator

$$M^+ f(x) = \sup_{h>0} \frac{1}{h} \int_x^{x+h} |f|,$$

the  $A_p^+$  classes were introduced by E. Sawyer [9]. He proved that  $M^+$  is bounded in  $L^p(w)$  ( $p > 1$ ) if, and only if, the weight satisfies  $A_p^+$ , i.e., there exists a constant  $C$

---

Research supported by D.G.I.Y.T. (PB94-1496), Junta de Andalucía and Universidad Nacional de Córdoba.

such that for any three points  $a < b < c$  we have

$$\int_a^b w \left( \int_b^c w^{1-p'} \right)^{p-1} \leq C(c-a)^p.$$

The smallest constant for which this is satisfied will be called the  $A_p^+$  constant of  $w$  and will be denoted by  $A_p^+(w)$ . For  $p = 1$  the weak type of the operator holds if, and only if, the weight  $w$  satisfies  $A_1^+$ , i.e., there exists  $C$  such that for any  $a$  and almost every  $b > a$ ,

$$\int_a^b w \leq C(b-a)w(b).$$

The smallest such constant will be called the  $A_1^+$  constant of  $w$  and will be denoted by  $A_1^+(w)$ . For later reference we point out that it is an easy consequence of Lebesgue's differentiation theorem that the constant in the definition of  $A_1^+$  is always greater than, or equal to, one.

These classes are of interest, not only because they control the boundedness of the one-sided Hardy-Littlewood maximal operator, but they are the right classes for the weighted estimates for one-sided singular integrals [1] and they also appear in PDE [4]. In contrast to the Muckenhoupt weights, the one-sided weights are not doubling, but they possess a one-sided doubling property, namely if  $w \in A_p^+$  then there exists  $C$  such that for any  $a \in \mathbb{R}$  and  $h > 0$ ,  $\int_a^{a+2h} w \leq C \int_{a+h}^{a+2h} w$ . The reverse Hölder property is not satisfied by these weights, either, but nevertheless, Martín-Reyes [5] proved that there is a weak substitute of this notion, that we will denote by  $RH_r^+$ , which is good enough to prove the “ $p - \varepsilon$ ” property. In [7] the class  $A_\infty^+$  was introduced and it was proved that  $A_\infty^+ = \bigcup_{p < \infty} A_p^+ = \bigcup_{1 < r} RH_r^+$ .

In this note we solve the problems of the Neugebauer paper in this context. In the proofs we will make essential use of the one-sided minimal operator introduced by Cruz-Uribe, Neugebauer and Olesen [3]. It is defined as  $m^+ f(x) = \inf_{c > x} \frac{1}{c-x} \int_x^c |f|$ . We will also use the fact that for any positive function  $g$ , the maximal operator  $M_g f(x) = \sup_{x \in I} \frac{1}{g(I)} \int_I |f| g dx$  is of weak type one-one with respect to the measure  $g dx$ . Note that for  $g = 1$  we have the classical Hardy-Littlewood maximal operator, which is denoted by  $Mf$ .

The paper is organized as follows: in Section 2 we give definitions and characterizations of  $RH_r^+$ ,  $1 < r < \infty$ . In Section 3 we prove two theorems of the best range for the extreme classes  $A_1^+$  and  $RH_\infty^+$ . In Section 4 we give a factorization theorem for weights in  $RH_r^+$ , and finally in Section 5 we extend the theorems of Section 3 to  $A_p^+$  and  $RH_r^+$ , using the factorization proved in Section 4. We shall see that the index range depends on the factorization of the weight.

We end this introduction with some notation: for a given interval  $I = (a, a + h)$  we denote by  $I^-$  the interval  $(a - h, a)$ ,  $I^+$  the interval  $(a + h, a + 2h)$ , and  $I^{++}$  the interval  $(a + 2h, a + 3h)$ . For any  $1 < p < \infty$ ,  $p'$  will be its conjugate exponent, if  $g$  is locally integrable and  $E$  is a measurable set,  $g(E)$  will stand for  $\int_E g$  and  $C$  will represent a constant that may change from time to time. Finally, we remark that we can change the orientation on the real line obtaining similar results for classes  $RH_r^-, A_p^-, 1 < r \leq \infty$  and  $1 \leq p \leq \infty$ .

## 2. DEFINITION, CHARACTERIZATION OF $RH_r^+$ FOR $1 < r < \infty$

We start this section with the definition of  $RH_r^+, 1 < r < \infty$ .

**Definition 2.1.** A weight  $w$  satisfies the one-sided reverse Hölder  $RH_r^+$  condition, if there exists  $C$  such that for any  $a < b$ ,

$$(2.2) \quad \int_a^b w^r \leq C (M(w\chi_{(a,b)})(b))^{(r-1)} \int_a^b w.$$

The smallest such constant will be called the  $RH_r^+$  constant of  $w$  and will be denoted by  $RH_r^+(w)$ .

**Definition 2.3.** A weight satisfies the one-sided reverse Hölder  $RH_\infty^+$  condition, if there exists  $C$  such that

$$(2.4) \quad w(x) \leq Cm^+w(x)$$

for almost all  $x \in \mathbb{R}$ .

The smallest such constant will be called the  $RH_\infty^+$  constant of  $w$  and will be denoted by  $RH_\infty^+(w)$ . It is clear that  $C \geq 1$ .

The following lemma gives several characterizations of  $RH_r^+$ . The constants are not necessarily the same.

**Lemma 2.5.** Let  $a < b < c < d, 1 < r < \infty$ , and let  $w \geq 0$  be locally integrable. Then the following statements are equivalent.

- (i)  $\int_a^b w^r \leq C (M(w\chi_{(a,b)})(b))^{(r-1)} \int_a^b w.$
- (ii)  $\frac{1}{b-a} \int_a^b w^r \leq C \left( \frac{1}{c-b} \int_b^c w \right)^r$  with  $b-a = 2(c-b).$
- (iii)  $\frac{1}{b-a} \int_a^b w^r \leq C \left( \frac{1}{d-c} \int_c^d w \right)^r$  with  $b-a = d-b = 2(d-c).$

- (iv)  $\frac{1}{b-a} \int_a^b w^r \leq C \left( \frac{1}{c-b} \int_b^c w \right)^r$  with  $b-a = c-b$ .
- (v)  $\frac{1}{b-a} \int_a^b w^r \leq C \left( \frac{1}{d-c} \int_c^d w \right)^r$  with  $b-a = d-c = \gamma(d-a)$ ,  $0 < \gamma \leq \frac{1}{2}$ .

Proof. To see  $i) \implies ii)$ , we fix  $a < b < c$ ,  $b-a = 2(c-b)$  and take any  $x \in (b, c)$ . Then

$$\int_a^b w^r \leq \int_a^x w^r \leq C (M(w\chi_{(a,x)})(x))^{r-1} \int_a^x w \leq C (M(w\chi_{(a,c)})(x))^{r-1} \int_a^c w.$$

Therefore  $(b, c) \subset \{x: (M(w\chi_{(a,c)})(x))^{r-1} \geq \frac{1}{C} \int_a^c w\}$ . The weak type  $(1, 1)$  of the Hardy-Littlewood maximal operator yields

$$(c-b) \left( \int_a^b w^r \right)^{\frac{1}{r-1}} \leq C \left( \int_a^c w \right)^{\frac{r}{r-1}},$$

which implies

$$\frac{1}{b-a} \int_a^b w^r \leq C \left( \frac{1}{c-b} \int_a^c w \right)^r \leq C \left( \frac{1}{c-b} \int_b^c w \right)^r;$$

the last inequality follows from the fact, proved in [7], that a weight satisfying  $i)$  satisfies  $A_p^+$  for some  $p$  and thus it satisfies the one-sided doubling condition.

We will prove now that  $ii) \implies i)$ . Let us fix  $a < b$  and define a sequence  $(x_k)$  as follows:  $x_0 = a$  and  $b-x_k = 2(b-x_{k+1})$ . In particular,  $x_{k+1}-x_k = 2(x_{k+2}-x_{k+1}) = (b-x_{k+1})$ . Using condition  $ii)$  for the points  $x_k, x_{k+1}, x_{k+2}$ , we have

$$\begin{aligned} \int_a^b w^r &= \sum_0^\infty \int_{x_k}^{x_{k+1}} w^r \leq C \sum_0^\infty (x_{k+1}-x_k)^{1-r} \left( \int_{x_{k+1}}^{x_{k+2}} w \right)^r \\ &\leq C \sum_0^\infty \int_{x_{k+1}}^{x_{k+2}} w \left( \frac{1}{b-x_{k+1}} \int_{x_{k+1}}^b w \right)^{r-1} \leq (M(w\chi_{(a,b)})(b))^{r-1} C \int_a^b w. \end{aligned}$$

To see  $ii) \implies iii)$  let  $a < b < c < d$  with  $b-a = d-b = 2(d-c)$ . Using that  $w$  satisfies the one-sided doubling condition, we have

$$\begin{aligned} \frac{1}{b-a} \int_a^b w^r &\leq C \left( \frac{1}{c-b} \int_b^c w \right)^r \leq C \left( \frac{d-c}{c-b} \frac{1}{d-c} \int_b^d w \right)^r \\ &\leq C \left( \frac{1}{d-c} \int_c^d w \right)^r. \end{aligned}$$

$iii) \implies iv)$  is immediate.

First of all we observe that iv) easily implies that the weight  $w$  satisfies the one-sided doubling condition. To see that iv)  $\implies$  v), let  $0 < \gamma \leq \frac{1}{2}$  and  $a < b < c < d$ ,  $b - a = d - c = \gamma(d - a)$ . Then if  $x$  is the midpoint between  $a$  and  $d$  we have

$$\frac{1}{b-a} \int_a^b w^r \leq \frac{1}{2\gamma} \frac{1}{x-a} \int_a^x w^r \leq \frac{C}{2\gamma} \left( \frac{1}{d-x} \int_x^d w \right)^r,$$

but it follows from the one-sided doubling condition that  $\int_x^d w \leq C_\gamma \int_c^d w$ .

Suppose v) holds, let  $a < b < c$ ,  $b-a = c-b = h$  and let us define for  $k = 0, 1, \dots, N$   $x_k = a + ksh$  and  $y_k = b + ksh$  where  $s = \frac{\gamma}{1-\gamma}$  and  $N$  is the first integer such that  $(N+1)s > 1$ . We observe that the choice of  $x_k, y_k$  has been made so that for any  $0 \leq k \leq (N-1)$  we have  $x_{k+1} - x_k = y_{k+1} - y_k = \gamma(y_{k+1} - x_k)$ . Applying v), using that  $r > 1$  and the fact that the intervals  $(y_k, y_{k+1})$  are disjoint, we have

$$\begin{aligned} \int_a^b w^r &\leq \sum_{k=0}^{N-1} \int_{x_k}^{x_{k+1}} w^r + \int_{b-sh}^b w^r \\ &\leq C(sh)^{1-r} \sum_{k=0}^{N-1} \left( \int_{y_k}^{y_{k+1}} w \right)^r + C(sh)^{1-r} \left( \int_{c-sh}^c w \right)^r \\ &\leq C_\gamma (c-a)^{1-r} \left( \int_b^c w \right)^r. \end{aligned}$$

So we have proved that v)  $\implies$  iv).

Finally, we will show that iv)  $\implies$  ii). Let  $a < b < c$  with  $b - a = 2(c - b)$ . Let  $x$  be the midpoint between  $a, b$ . Using the one-sided doubling property we have

$$\begin{aligned} \frac{1}{b-a} \int_a^b w^r &= \frac{1}{b-a} \left( \int_a^x w^r + \int_x^b w^r \right) \\ &= \frac{1}{2} \left( \frac{1}{x-a} \int_a^x w^r + \frac{1}{b-x} \int_x^b w^r \right) \\ &\leq \frac{C}{2} \left( \left( \frac{1}{b-x} \int_x^b w \right)^r + \left( \frac{1}{c-b} \int_b^c w \right)^r \right) \\ &\leq \frac{C}{2} \left( \left( \frac{1}{c-b} \int_x^c w \right)^r + \left( \frac{1}{c-b} \int_b^c w \right)^r \right) \\ &\leq C \left( \frac{1}{c-b} \int_b^c w \right)^r. \end{aligned}$$

□

**Remark.** The equivalence of i) and iv) was first proved in [3].

The following lemma tells us that in the definition of  $A_p^+$  we can take two intervals that are not contiguous. Note that in the case of  $RH_r^+$  we have seen this in the previous lemma.

**Lemma 2.6.** *A weight  $w$  belongs to  $A_p^+$ ,  $p > 1$  if, and only if, there exist  $0 < \gamma \leq \frac{1}{2}$  and a constant  $C_\gamma$  such that  $b - a = d - c = \gamma(d - a)$  for any  $a < b < c < d$ , then*

$$(2.7) \quad \int_a^b w \left( \int_c^d w^{1-p'} \right)^{p-1} \leq C_\gamma (b - a)^p.$$

*Proof.* If  $w \in A_p^+$ ,  $0 < \gamma \leq \frac{1}{2}$  and  $a < b < c < d$ ,  $b - a = d - c = \gamma(d - a)$  then

$$\int_a^b w \left( \int_c^d w^{1-p'} \right)^{p-1} \leq \int_a^c w \left( \int_c^d w^{1-p'} \right)^{p-1} \leq C(d - a)^p = C_\gamma (b - a)^p.$$

To prove that (2.7) implies  $A_p^+$  we will show that (2.7) implies that for  $\gamma$  and  $a, b, c, d$  as above we have

$$\frac{1}{b - a} \int_a^b w \exp \left( \frac{1}{d - c} \int_c^d -\log(w) \right) \leq C.$$

Indeed,

$$(2.8) \quad \begin{aligned} & \frac{1}{b - a} \int_a^b w \exp \left( \frac{1}{d - c} \int_c^d -\log(w) \right) \\ &= \frac{1}{b - a} \int_a^b \left[ w \exp \left( \frac{1}{d - c} \int_c^d \log(w)^{1-p'} \right) \right]^{p-1} \\ & \leq \frac{1}{b - a} \int_a^b w \left( \frac{1}{d - c} \int_c^d w^{1-p'} \right)^{p-1} \leq C. \end{aligned}$$

In the same way we prove that  $w^{1-p'}$  satisfies

$$(2.9) \quad \exp \left( \frac{1}{b - a} \int_a^b \log(w)^{p'-1} \right) \frac{1}{d - c} \int_c^d w^{1-p'} \leq C.$$

But, according to part *j*) of Theorem 1 in [7], (2.8) is equivalent to saying that  $w \in A_\infty^+$  while (2.9) means that  $w^{1-p'} \in A_\infty^-$ , and according to Theorem 2 in [7] these two conditions imply  $w \in A_1^+$ .  $\square$

**Remark 2.10.** We can easily see that  $w \in A_1^+$  if, and only if, there exists  $C > 0$  such that  $\frac{1}{h} \int_{a-h}^a w \leq Cw(a + h)$  for almost every  $a \in \mathbb{R}$  and  $h > 0$ .

### 3. THE EXTREME CASES: $A_1^+$ AND $RH_\infty^+$

**Theorem 3.1.** *Let  $w \in A_1^+$  with  $A_1^+$  constant  $C > 1$ . Then  $w \in RH_r^+$  for any  $1 < r < \frac{C}{C-1}$ , and this is the best possible range.*

*Proof.* Let us fix the interval  $I = (a, b)$ . We consider the truncation of  $w$  at height  $N$  defined by  $w_N = \min(w, N)$ , which also satisfies  $A_1^+$  with a constant  $C_N \leq C$ . We claim that if  $\lambda_I = M(w_N \chi_I)(b)$  and  $E_\lambda = \{x \in I : w_N(x) > \lambda\}$  then

$$(3.2) \quad \int_{E_\lambda} w_N \leq C_N \lambda |E_\lambda| \quad \forall \lambda \geq \lambda_I.$$

Indeed, if  $E_\lambda = I$  we do not even need the  $A_1^+$  condition, since

$$w_N(E_\lambda) = \int_a^b w_N \leq M(w_N \chi_I)(b)(b-a) = \lambda_I(b-a) \leq C_N \lambda |E_\lambda|.$$

If  $E_\lambda \neq I$  we fix  $\varepsilon > 0$  and an open set  $O$  such that  $E_\lambda \subset O \subset I$  and  $|O| \leq \varepsilon + |E_\lambda|$ . Let  $J_k = (c, d)$ , be one of the connected components of  $O$ . There are two cases:

- (1)  $a \leq c < d < b$ ,
- (2)  $a \leq c < d = b$ .

In the first case  $d \notin E_\lambda$  and then  $w_N(d) \leq \lambda$ . Now  $A_1^+$  gives  $\int_c^d w_N \leq C_N w_N(d)(d-c) \leq C_N \lambda(d-c)$ . The second case is handled as the case  $E_\lambda = I$ , since  $\int_c^b w_N \leq M(w_N \chi_I)(b)(b-c) \leq C \lambda(b-c)$ . In any case  $w_N(J_k) \leq C_N \lambda |J_k|$ . Adding up we get

$$w_N(E_\lambda) \leq w_N(O) \leq C_N \lambda |O| \leq C_N \lambda (\varepsilon + |E_\lambda|).$$

Since  $\varepsilon$  was arbitrary we are done. Now we proceed in the standard way, i.e., we fix  $s > -1$ , multiply both sides of (3.2) by  $\lambda^s$  and integrate from  $\lambda_I$  to infinity to obtain,

$$\frac{1}{s+1} \int_I (w_N^{s+2} - \lambda_I^{s+1} w_N) \leq \frac{C_N}{s+2} \int_I w_N^{s+2}.$$

Now if  $r = s+2 < \frac{C_N}{C_N-1}$  then  $\frac{1}{s+1} - \frac{C_N}{s+2} > 0$ , and we get

$$\int w_N^r \leq C_N \lambda_I^{r-1} \int_I w_N = C_N (M(w_N \chi_I)(b))^{r-1} \int_I w_N.$$

Now  $C_N \leq C$  implies  $\frac{C_N}{C_N-1} \geq \frac{C}{C-1}$ , and therefore if  $r \leq \frac{C}{C-1}$  then

$$\int_a^b w_N^r \leq C_N (M(w_N \chi_{(a,b)})(b))^{r-1} \int_a^b w_N \leq C (M(w \chi_{(a,b)})(b))^{r-1} \int_a^b w$$



and the monotone convergence theorem gives  $w \in RH_r^+$ . To see that this is the best possible range we consider the function

$$w(x) = x^{\frac{1}{c}-1} \chi_{(0,\infty)}(x).$$

It is clear that  $w$  does not satisfy  $RH_{\frac{C}{c-1}}^+$  because  $w^{\frac{C}{c-1}}(x) = \frac{1}{x}$  for  $x > 0$ . To see that it satisfies  $A_1^+$  with the constant  $C$ , we consider three cases:

- (1)  $a < b \leq 0$ ,
- (2)  $a \leq 0 < b$ ,
- (3)  $0 < a < b$ .

In the first case there is nothing to check. In the second case  $\frac{1}{b-a} \int_a^b w < \frac{1}{b} \int_0^b w(x) = \frac{C}{b} b^{\frac{1}{c}} = Cw(b)$ . Finally, if  $0 < a < b$ , then  $\int_a^b w = C(b^{\frac{1}{c}} - a^{\frac{1}{c}}) \leq C(b-a)w(b)$ .  $\square$

**Remark.** Note that if  $C = 1$ , then  $w(x) = M^-w(x)$ , and this implies that  $w$  is non-decreasing. This tells us that  $w \in RH_\infty^+$ .

**Theorem 3.3.** *If  $w$  satisfies  $RH_\infty^+$  with a constant  $C > 1$ , then  $w \in A_p^+$  for all  $p > C$ , and this is the best possible range.*

**Proof.** A truncation argument as in Theorem 3.1 allows us to suppose that  $w$  is bounded away from zero, i.e. there exists  $\beta > 0$  such that  $w(x) \geq \beta$  for all  $x$ . Let us fix  $I = (a, b)$  and consider  $\lambda_I = m^+(w \frac{1}{\chi_I})(a)$ . We claim that if  $\lambda < \lambda_I$  and  $E_\lambda = \{x \in I : w(x) < \lambda\}$ , then

$$(3.4) \quad \lambda|E_\lambda| \leq C \int_{E_\lambda} w.$$

As before, if  $E_\lambda = I$  then  $\lambda|E_\lambda| = \lambda(b-a) < \lambda_I(b-a) = \int_a^b w \leq w(E_\lambda)$ . If  $E_\lambda \neq I$  then we approximate it by an open set  $O = \bigcup J_k$  where  $E_\lambda \subset O \subset I$  and  $w(O) < \varepsilon + w(E_\lambda)$ . Let us fix  $J_k = (c, d)$ . There are two cases:

- (1)  $a < c$ ,
- (2)  $a = c$ .

In the first case  $c \notin E_\lambda$  and then  $\lambda(d-c) \leq w(c)(d-c) \leq Cm^+w(c)(d-c) \leq C \int_c^d w$ . In the second case  $\lambda(d-c) \leq \lambda_I(d-a) \leq \int_a^d w$ , and (3.4) follows. If we multiply both sides of (3.4) by  $\lambda^{-r}$  with  $r > 2$  and integrate we have

$$\int_0^{\lambda_I} \lambda^{1-r} \int \chi_{E_\lambda}(x) dx d\lambda \leq C \int_0^\infty \lambda^{-r} \int_{E_\lambda} w(x) dx d\lambda.$$

For the left hand side we obtain

$$\begin{aligned} \int_{\beta}^{\lambda_I} \lambda^{1-r} \int \chi_{E_\lambda}(x) dx d\lambda &= \frac{1}{2-r} \int_{\{x \in I: w(x) < \lambda_I\}} \lambda_I^{2-r} - w^{2-r} dx \\ &\geq \frac{1}{2-r} \int_I \lambda_I^{2-r} - w^{2-r} dx = \frac{1}{r-2} \int_I w^{2-r} - \frac{|I|}{r-2} \lambda_I^{2-r}, \end{aligned}$$

while the right hand side is equal to  $\frac{C}{r-1} \int_I w^{2-r}$ . Therefore

$$\frac{1}{r-2} \int_I w^{2-r} \leq \frac{C}{r-1} \int_I w^{2-r} + \frac{|I|}{r-2} \lambda_I^{2-r}.$$

If we choose  $r > 2$  such that  $C(r-2) < (r-1)$ , we obtain that there exists  $C$  such that

$$(3.5) \quad \frac{1}{|I|} \int_I w^{2-r} \leq C \left( m^+ \left( \frac{w}{\chi_I} \right) (a) \right)^{2-r}.$$

We now claim that (3.5) implies that  $w \in A_p^+$  with  $p = \frac{r-1}{r-2}$ . Let us fix  $a < b < c$  and choose  $x \in (a, b)$ . If we keep in mind that  $1 - p' = 2 - r$  we may write

$$\left( \frac{1}{c-a} \int_b^c w^{1-p'} \right)^{p-1} \leq \left( \frac{1}{c-x} \int_x^c w^{1-p'} \right)^{p-1} \leq C \left( m^+ \left( \frac{w}{\chi_{(x,c)}} \right) (x) \right)^{-1},$$

but

$$\left( m^+ \left( \frac{w}{\chi_{(x,c)}} \right) (x) \right)^{-1} = \left( \inf_{x < d < c} \frac{1}{d-x} \int_x^d w \right)^{-1} = \sup_{x < d < c} \frac{d-x}{\int_x^d w} = M_w \left( \frac{\chi_{(a,c)}}{w} \right) (x).$$

We have thus proved that if  $\lambda = \left( \frac{1}{c-a} \int_b^c w^{1-p'} \right)^{p-1}$  then

$$(a, b) \subset \left\{ x : CM_w \left( \frac{\chi_{(a,c)}}{w} \right) (x) > \lambda \right\},$$

and the weak type of  $M_w$  with respect to the measure  $w dx$  yields  $\int_a^b w \leq C(c-a)^p \left( \int_b^c w^{1-p'} \right)^{1-p}$  which is  $A_p^+$ . Finally, it can be checked that the function  $w(x)$  which is 0 for  $x < -1$ , identically one for  $x > 0$  and  $|x|^{C-1}$  between  $-1$  and  $0$ , satisfies  $RH_\infty^+$  with a constant  $C$ , but is not in  $A_C^+$ .  $\square$

**Remark.** Note that if  $C = 1$ , then  $w(x) = m^+ w(x)$ , and this implies that  $w$  is non-decreasing. This tells us that  $w \in A_1^+$ .

We have had several different characterizations of  $RH_r^+$ , one involved the maximal operator, but dealt with one interval, and the others involved two intervals but no operator. We can now prove that for  $RH_\infty^+$  the situation is the same, we can characterize  $RH_\infty^+$  using two intervals instead of the minimal operator.

**Corollary 3.6.** *We have  $w \in RH_\infty^+$  if, and only if, there exists  $C$  such that for any interval  $I$ ,*

$$(3.7) \quad \operatorname{ess\,sup}_I w \leq C \frac{1}{|I^+|} \int_{I^+} w.$$

*Proof.* It is immediate that (3.7) implies  $RH_\infty^+$ . Assume now that  $w \in RH_\infty^+$ . The preceding theorem tells us that  $w \in A_p$  for some  $p$ , and therefore it satisfies the one-sided doubling condition. Therefore if  $I = (a, b)$  is any interval,  $I^+ = (b, c)$  and  $x \in I$ , we have

$$w(x) \leq \frac{C}{c-x} \int_x^c w \leq \frac{C}{c-b} \int_b^c w,$$

which is (3.7). □

**Remark.** Note that with this definition, we have  $RH_\infty^+ \subset \bigcap_{r>1} RH_r^+$ .

#### 4. FACTORIZATION OF WEIGHTS IN $RH_r^+$ , $1 < r \leq \infty$

The theorems on the best range for weights in  $A_p^+$  ( $p > 1$ ) or in  $RH_r^+$ ,  $r < \infty$  will be stated in terms of factorizations of the given weight. Therefore this section will be devoted to proving a factorization of functions in  $RH_r^+$ . The bilateral case was studied in [2].

**Definition 4.1.** A function  $w$  is said to be essentially increasing if there exists  $C$  such that  $w(x) \leq Cw(y)$  for any  $x < y$ .

**Lemma 4.2.** *A function belongs to  $RH_\infty^+ \cap A_1^+$  if, and only if, it is essentially increasing.*

*Proof.* Assume that  $w \in RH_\infty^+ \cap A_1^+$  and  $x < y$ , then  $w(x) \leq C \frac{1}{y-x} \int_x^y w \leq Cw(y)$  and  $w$  is essentially increasing. Conversely, if  $w$  is essentially increasing then for any  $x$  and  $h > 0$  we have  $w(x) \leq \frac{C}{h} \int_x^{x+h} w$ , hence  $w \in RH_\infty^+$ . On the other hand,  $\frac{1}{h} \int_{x-h}^x w \leq Cw(x)$ , so  $w \in A_1^+$  □

**Lemma 4.3.** Let  $1 < r \leq \infty$  and  $1 \leq p < \infty$ .

- (1) If  $u$  is essentially increasing and  $v \in RH_r^+$  then  $w \in RH_r^+$ .
- (2) If  $u$  is essentially increasing and  $v \in A_p^+$  then  $w \in A_p^+$ .

*P r o o f.* This proof follows immediately from Definition 4.1. □

**Lemma 4.4.** Let  $1 < r \leq \infty$  and  $1 \leq p < \infty$ . We have  $w \in RH_r^+ \cap A_p^+$  if, and only if,  $w^r \in A_q^+$ , with  $q = r(p - 1) + 1$ .

*P r o o f.* Let  $C_1 = RH_r^+(w)$  and  $C_2 = A_p^+(w)$ ,  $w \in RH_r^+ \cap A_p^+$  and  $q = r(p - 1) + 1$ . Also note that  $1 - q' = 1 - \frac{r(p-1)+1}{r(p-1)} = \frac{1}{r(1-p)}$ ,

$$\begin{aligned} & \left( \frac{1}{|I^-|} \int_{I^-} w^r \right) \left( \frac{1}{|I^+|} \int_{I^+} w^{r(1-q')} \right)^{q-1} \\ & \leq C_1 \left( \frac{1}{|I|} \int_I w \right)^r \left( \frac{1}{|I^+|} \int_{I^+} w^{1-p'} \right)^{r(p-1)} \\ & \leq C_1 C_2^r, \end{aligned}$$

and by Lemma 2.6 we have that  $w^r \in A_q^+$ .

If  $w^r \in A_q^+$ , then by Hölder's inequality

$$\begin{aligned} & \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{p-1} \\ & \leq \left( \frac{1}{|I|} \int_I w^r \right)^{1/r} \left( \frac{1}{|I^+|} \int_{I^+} w^{-r/(q-1)} \right)^{(q-1)/r} \\ & \leq C^{1/r}, \end{aligned}$$

and we obtain in this way that  $w \in A_p^+$ . Now again by Hölder's inequality

$$1 = \frac{1}{|I^+|} \int_{I^+} w^{-1/p} w^{1/p} \leq \left( \frac{1}{|I^+|} \int_{I^+} w \right)^{1/p} \left( \frac{1}{|I^+|} \int_{I^+} w^{-p'/p} \right)^{1/p'},$$

so

$$\left( \frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{1-p} \leq \frac{1}{|I^+|} \int_{I^+} w,$$

and we get

$$\begin{aligned} \left( \frac{1}{|I|} \int_I w^r \right)^{1/r} & \leq C \left( \frac{1}{|I^+|} \int_{I^+} w^{-r/(q-1)} \right)^{-(q-1)/r} = C \left( \frac{1}{|I^+|} \int_{I^+} w^{-1/(p-1)} \right)^{1-p} \\ & \leq C \frac{1}{|I^+|} \int_{I^+} w, \end{aligned}$$

proving that  $w \in RH_r^+$ . □

**Factorization Theorem for weights in  $RH_r^+ \cap A_p^+$ .** A weight  $w$  satisfies  $w \in RH_r^+ \cap A_p^+$  with  $1 \leq p < \infty$ ,  $1 < r \leq \infty$  if, and only if, there exist weights  $w_0$  and  $w_1$  such that  $w_0 \in RH_r^+ \cap A_1^+$ ,  $w_1 \in RH_\infty^+ \cap A_p^+$  and  $w = w_0 w_1$ .

Observe that since  $\bigcup_{p < \infty} A_p^+ = \cap_{1 < r} RH_r^+$ , every weight in  $RH_r^+$  is in some  $A_p^+$ . See [7].

**P r o o f.** Let us first consider the cases  $p = 1$  or  $r = \infty$ .

If  $p = 1$  and  $r \leq \infty$ , we put  $w_1 = 1$  and  $w_0 = w$ , then obviously  $w_0 \in RH_r^+ \cap A_1^+$  and  $w_1 \in RH_\infty^+ \cap A_1^+$ .

If  $p \geq 1$  and  $r = \infty$ , we put  $w_0 = 1$  and  $w_1 = w$ , obtaining  $w_0 \in RH_\infty^+ \cap A_1^+$ ,  $w_1 \in RH_\infty^+ \cap A_p^+$ .

Conversely, given  $w_0$  and  $w_1$ , at least one of them belongs to  $RH_\infty^+ \cap A_1^+$  (because  $p = 1$  or  $r = \infty$ ), so one of them is essentially increasing, therefore  $w_0 w_1 \in RH_r^+ \cap A_p^+$  (Lemma 4.3).

Let us now suppose  $p > 1$  and  $r < \infty$ . Let  $w = w_0 w_1$  with  $w_0 \in RH_r^+ \cap A_1^+$ , and  $w_1 \in RH_\infty^+ \cap A_p^+$ . We want to see that  $w \in RH_r^+ \cap A_p^+$ . Note that for  $w_1$  we have

$$\frac{1}{|I|} \int_I w_1^{1-p'} \leq C \left( \frac{1}{|I^-|} \int_{I^-} w_1 \right)^{1-p'} \leq C w_1(a-h)^{1-p'},$$

which implies  $w_1^{1-p'} \in A_1^-$  (Remark 2.10). Let  $v = w_1^{1-p'}$ , then  $w_1 = v^{1-p}$  with  $v \in A_1^-$ , so  $w = w_0 w_1 = w_0 v^{1-p}$  with  $w_0 \in A_1^+$  and  $v \in A_1^-$  (see [7]), and this implies  $w \in A_p^+$ .

Now

$$\begin{aligned} \frac{1}{|I|} \int_I w^r &= \frac{1}{|I|} \int_I w_0^r w_1^r \leq \left( \sup_I w_1 \right)^r C \left( \frac{1}{|I^+|} \int_{I^+} w_0 \right)^r \\ &\leq C \left( \frac{1}{|I^+|} \int_{I^+} w_1 \right)^r \left( \inf_{I^+} w_0 \right)^r \\ &\leq C \left( \frac{1}{|I^+|} \int_{I^+} w_0 w_1 \right)^r, \end{aligned}$$

and by Lemma 2.5 we have  $w \in RH_r^+$ . Conversely, let  $w \in RH_r^+ \cap A_p^+$ , then by Lemma 4.4  $w^r \in A_q^+$  with  $q = r(p-1) + 1$ , there exists  $v_0 \in A_1^+$  and  $v_1 \in A_1^-$  such that  $w^r = v_0 v_1^{1-q}$  (see [7]), or equivalently  $w = v_0^{1/r} v_1^{(1-q)/r} = v_0^{1/r} v_1^{1-p}$ . Let  $w_0 = v_0^{1/r}$  and  $w_1 = v_1^{1-p}$ . We will see that  $w_0 \in RH_r^+ \cap A_1^+$ . We note,

$$\begin{aligned} \frac{1}{|I|} \int_I w_0^r &= \frac{1}{|I|} \int_I v_0 \leq C \inf_{I^+} v_0 \\ &\leq C \left( \frac{1}{|I^+|} \int_{I^+} v_0^{1/r} \right)^r = C \left( \frac{1}{|I^+|} \int_{I^+} w_0 \right)^r, \end{aligned}$$

and also

$$\begin{aligned} \frac{1}{|I|} \int_I w_0 &= \frac{1}{|I|} \int_I v_0^{1/r} \leq \left( \frac{1}{|I|} \int_I v_0 \right)^{1/r} \\ &\leq C \inf_{I^+} v_0^{1/r} = C \inf_{I^+} w_0. \end{aligned}$$

We only have to see now that  $w_1 \in RH_\infty^+ \cap A_p^+$  and we are done.

First we claim

$$(4.5) \quad \text{if } w \in A_1^- \text{ then } w^{-\gamma} \in RH_\infty^+ \text{ for all } \gamma > 0.$$

In fact, by Hölder's inequality we have  $(\frac{1}{|I|} \int_I w)^{-\gamma} \leq \frac{1}{|I|} \int_I w^{-\gamma}$  for any interval  $I = (a, b)$ , and as  $w \in A_1^-$  we have that  $Cw(x) \geq \frac{1}{|I|} \int_I w$  for almost every  $x \in I^-$ , and therefore

$$w(x)^{-\gamma} \leq C \left( \frac{1}{|I|} \int_I w \right)^{-\gamma} \leq \frac{1}{|I|} \int_I w^{-\gamma} \leq C \frac{1}{b-x} \int_x^b w^{-\gamma}.$$

Let  $w_1 = v_1^{1-p}$ . As  $v_1 \in A_1^-$ , then  $w_1 \in RH_\infty^+$ . Moreover,

$$\begin{aligned} \frac{1}{|I|} \int_I w_1 \left( \frac{1}{|I^+|} \int_{I^+} w_1^{1-p'} \right)^{p-1} &= \frac{1}{|I|} \int_I v_1^{1-p} \left( \frac{1}{|I^+|} \int_{I^+} v_1 \right)^{p-1} \\ &\leq \frac{1}{|I|} \int_I v_1^{1-p} (C \inf_I v_1)^{p-1} \\ &\leq \frac{C}{|I|} \int_I v_1^{1-p} v_1^{p-1} \leq C, \end{aligned}$$

i.e.  $w_1 \in A_p^+$ . □

**Factorization Theorem for weights in  $A_\infty^+$ .** A weight  $w$  satisfies  $w \in A_\infty^+$  if, and only if, there exist  $w_1 \in RH_\infty^+$  and  $w_0 \in A_1^+$  such that  $w = w_0 w_1$ .

*P r o o f.* If  $w \in A_\infty^+$  then  $w \in A_q^+$  for some  $1 < q < \infty$ , so there exist  $v_0 \in A_1^+$  and  $v_1 \in A_1^-$  such that  $w = v_0 v_1^{1-q}$ . Let  $w_0 = v_0$  and  $w_1 = v_1^{1-q}$ . By (4.5),  $w_1 \in RH_\infty^+$ . So we are done. Conversely, if  $w_1 \in RH_\infty^+$ , then  $w_1 \in A_q^+$  for some  $1 < q$ , i.e., there exists  $C$  such that

$$\left( \frac{1}{|I|} \int_I w_1 \right)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C,$$

but then

$$(\sup_{I^-} w_1)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq \left( \frac{1}{|I|} \int_I w_1 \right)^{q'-1} \frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C,$$

and we get

$$\frac{1}{|I|} \int_{I^+} w_1^{1-q'} \leq C \inf_{I^-} w_1^{1-q'},$$

and it is easy to see that this inequality implies  $w_1^{1-q'} \in A_1^-$ . Then  $v_1 = w_1^{1-q'} \in A_1^-$ , so  $w = w_0 w_1 = w_0 v_1^{1-q} \in A_q^+ \subset A_\infty^+$ .  $\square$

## 5. CLASSES $A_p^+$ AND $RH_r^+$

In this section we will use Theorems 3.1 and 3.3 and the factorization theorems to obtain the best ranges for the classes  $A_p^+$  and  $RH_r^+$ . As we shall see, the range of the index will depend on the factorization of the weights.

The following theorem gives us the precise range in  $A_p^+$  for weights in  $RH_r^+$ .

**Theorem 5.1.** *Let  $w \in RH_r^+$ ,  $w = w_0 w_1^{\frac{1}{r}}$  with  $w_0 \in RH_\infty^+$  and  $w_1 \in A_1^+$ . Then  $w \in A_p^+$  for all  $p > C$  where  $C = RH_\infty^+(w_0)$ , and this is the best possible range.*

*Proof.* Let  $w_0 \in RH_\infty^+$  and  $w_1 \in A_1^+$ . By Theorem 3.3,  $w_0 \in A_p^+$  for all  $p > C$ . Let  $p > C$ , then there exists  $\varepsilon > 0$  such that  $w_0 \in A_{p-\varepsilon}^+$ , so we choose  $s > 1$  satisfying  $1 - (p - \varepsilon)' = s(1 - p')$ , and by Hölder's inequality

$$\begin{aligned} & \frac{1}{|I^-|} \int_{I^-} w_0 w_1 \left( \frac{1}{|I^+|} \int_{I^+} (w_0 w_1)^{1-p'} \right)^{p-1} \\ & \leq \left( \frac{1}{|I|} \int_I w_0 \right) \left( \frac{1}{|I^-|} \int_{I^-} w_1 \right) \left( \frac{1}{|I^+|} \int_{I^+} w_0^{s(1-p')} \right)^{\frac{(p-1)}{s}} \left( \frac{1}{|I^+|} \int_{I^+} w_1^{s'(1-p')} \right)^{\frac{(p-1)}{s'}} \\ & \leq C. \end{aligned}$$

To see that this is the best range, we consider  $w_0$  as in Theorem 3.3 and  $w_1 = 1$ .  $\square$

**Remark 5.2.** Given  $w \in RH_r^+$  there exist  $u \in RH_\infty^+$ , and  $v \in A_1^+$  such that  $w = uv^{\frac{1}{r}}$ . We only have to consider the factorization theorem and choose  $u = w_1$  and  $v = w_0^r$ . We have to prove that  $v \in A_1^+$ . Keeping in mind that  $w_0 \in RH_r^+ \cap A_1^+$  we have

$$\frac{1}{|I^-|} \int_{|I^-|} v = \frac{1}{|I^-|} \int_{|I^-|} w_0^r \leq C \left( \frac{1}{|I|} \int_{|I|} w_0 \right)^r \leq C w_0^r(x) = Cv(x)$$

for almost every  $x \in I^+$ , i.e.,  $v \in A_1^+$ .

The next theorem shows us the precise range of the higher integrability of  $w \in RH_r^+$ .

**Theorem 5.3.** *Let  $w \in RH_r^+$ ,  $w = uv^{1/r}$  with  $u \in RH_\infty^+$  and  $v \in A_1^+$ . If  $C = A_1^+(v)$  then  $w \in RH_s^+$  for all  $r \leq s < \frac{Cr}{C-1}$ . The range of  $s$  is the best possible.*

*Proof.* Let  $r < s < \frac{Cr}{C-1}$ , let us choose  $q > 1$  such that  $s < \frac{Cr}{q(C-1)}$ . As  $1 < \frac{qs}{r} < \frac{C}{C-1}$ , by Theorem 3.1 we have  $v \in RH_{\frac{qs}{r}}^+$ , and using Hölder's inequality we obtain that  $u^s \in RH_\infty^+$  and  $v \in A_1^+$  which yields

$$\begin{aligned} \frac{1}{|I|} \int_I w^s &= \frac{1}{|I|} \int_I u^s v^{s/r} \leq \left( \frac{1}{|I|} \int_I u^{q's} \right)^{1/q'} \left( \frac{1}{|I|} \int_I v^{qs/r} \right)^{1/q} \\ &\leq \sup_I u^s C \left( \frac{1}{|I^+|} \int_{I^+} v \right)^{s/r} \leq \frac{C}{|I^+|} \int_{I^+} u^s \left( \inf_{I^{++}} v \right)^{s/r} \\ &\leq C \sup_{I^+} u^s \inf_{I^{++}} v^{s/r} \leq C \left( \frac{1}{|I^{++}|} \int_{I^{++}} u \right)^s \inf_{I^{++}} v^{s/r} \\ &\leq C \left( \frac{1}{|I^{++}|} \int_{I^{++}} uv^{1/r} \right)^s = C \left( \frac{1}{|I^{++}|} \int_{I^{++}} w \right)^s, \end{aligned}$$

and we get that  $w \in RH_s^+$  (Lemma 2.5).

To see this is the best range possible, we choose  $v \in A_1^+$  as in Theorem 3.1 and  $u = 1$ , then  $w = v^{1/r} \in RH_s^+$  for all  $r \leq s < \frac{Cr}{C-1}$  ( $C = A_1^+(v)$ ). If  $s = \frac{Cr}{C-1}$  and  $w \in RH_s^+$  then  $v \in RH_{\frac{C}{C-1}}^+$ , but we have seen (Theorem 3.1) that this can not happen.  $\square$

The next theorem shows us which is the best range in  $RH_r^+$  for a given weight in  $A_p^+$ .

**Theorem 5.4.** *Let  $w \in A_p^+$ ,  $w = uv^{1-p}$  with  $u \in A_1^+$ ,  $v \in A_1^-$  and  $C = A_1^+(u)$ , then  $w \in RH_r^+$  for all  $1 < r < \frac{C}{C-1}$ , this range being the best possible.*

*Proof.* By Theorem 3.1 we have  $u \in RH_r^+$  for all  $1 < r < \frac{C}{C-1}$  and we know that  $v^{1-p} \in RH_\infty^+$ , hence

$$\begin{aligned} \frac{1}{|I|} \int_I w^r &\leq \frac{1}{|I|} \int_I u^r \sup_I \left( v^{-r(p-1)} \right) \\ &\leq C \left( \frac{1}{|I^+|} \int_{I^+} u \right)^r \left( \frac{1}{|I^+|} \int_{I^+} v^{1-p} \right)^r \leq C \left( \inf_{I^{++}} u \right)^r \left( \sup_{I^+} v^{1-p} \right)^r \\ &\leq C \left( \inf_{I^{++}} u \right)^r \left( \frac{1}{|I^{++}|} \int_{I^{++}} v^{1-p} \right)^r \leq C \left( \frac{1}{|I^{++}|} \int_{I^{++}} w \right)^r. \end{aligned}$$

By Lemma 2.5 we conclude  $w \in RH_r^+$ .



To see this is the best range we take  $u$  as in Theorem 3.1 and  $v = 1$ . So we have  $w = u \in A_p^+$  and  $w \notin RH_{\frac{C}{C-1}}^+$ .  $\square$

**Corollary 5.5.** *Let  $w = uv^{1-p} \in A_p^+$  with  $u \in A_1^+$ ,  $v \in A_1^-$  and  $C = \max\{A_1^+(u), A_1^-(v)\}$ . Then  $w^\tau \in A_p^+$  for all  $1 \leq \tau < \frac{C}{C-1}$  and the range is the best possible.*

*Proof.* By Theorem 5.4 we have that  $w \in RH_\tau^+$  for all  $1 \leq \tau < \frac{C}{C-1}$  and  $w^{1-p'} \in RH_\tau^-$  for all  $1 \leq \tau < \frac{C}{C-1}$ . Let  $a < d$ , let us choose  $b, c$  such that  $b - a = d - c = \frac{1}{4}(d - a)$ , and we also consider the point  $\frac{c+b}{2}$ . Then we have four intervals, namely,  $I^- = (a, b)$ ,  $I = (b, \frac{b+c}{2})$ ,  $I^+ = (\frac{b+c}{2}, c)$ , and  $I^{++} = (c, d)$ . Now

$$\frac{1}{|I^-|} \int_{I^-} w^\tau \left( \frac{1}{|I^{++}|} \int_{I^{++}} w^{\tau(1-p')} \right)^{p-1} \leq \left( \frac{1}{|I|} \int_I w \right)^\tau \left( \frac{1}{|I^+|} \int_{I^+} w^{1-p'} \right)^{\tau(p-1)},$$

$$\leq C^\tau,$$

thus  $w^\tau \in A_p^+$  (Lemma 2.6). Considering  $u$  as in Theorem 3.1, we see this is the best possible range.  $\square$

Using Theorem 5.4 we will show the exact range of  $q < p$  such that  $w \in A_p^+$  implies  $w \in A_q^+$ .

**Theorem 5.6.** *Let  $w = uv^{1-p} \in A_p^+$  with  $u \in A_1^+$ ,  $v \in A_1^-$  and  $C = A_1^-(v)$ . Then  $w \in A_q^+$  for all  $1 + \frac{(p-1)(C-1)}{C} < q < \infty$  and this is the best range for  $q$ .*

*Proof.* Note that  $w^{1-p'} = vu^{1-p'} \in A_{p'}^-$ , by Theorem 5.4 we have  $w^{1-p'} \in RH_r^-$  for all  $1 < r < \frac{C}{C-1}$ . For the classes  $RH_r^-$  and  $A_p^-$  we have from Lemma 4.4 that  $w^{(1-p')r} \in A_{q'}^-$ , where  $q' = r(p'-1) + 1 = \frac{r}{p-1} + 1$ . But this is the same as  $w^{1-q'} \in A_{q'}^-$ , i.e.,  $w \in A_q^+$  for all  $1 + (p-1)\frac{C-1}{C} < q$ .

To see this is the best range, let  $v(x) = x^{\frac{1-C}{C}}$  if  $x \leq 0$  and equal to 0 if  $x > 0$  and  $u = 1$  for all  $x$ . Note that  $v \in A_1^-$  and  $A_1^-(v) = C$ . Then  $w = v^{1-p} \in A_p^+$  and  $w \in A_q^+$  for all  $q > 1 + (p-1)\frac{C-1}{C}$ . Observe that  $w \notin A_{1+(p-1)\frac{C-1}{C}}^+$ .  $\square$

Finally, the last theorem gives us the best possible range for a weight in  $A_\infty^+$ .

**Theorem 5.7.** *Let  $w \in A_\infty^+$ ,  $w = w_0 w_1$ ,  $w_0 \in A_1^+$ ,  $w_1 \in RH_\infty^+$  and  $C = RH_\infty^+(w_1)$ . Then  $w \in A_p^+$  for all  $p > C$ . The range of  $p$ 's is the best possible.*

Proof. Note that  $w_1 \in RH_\infty^+$  implies  $w_1 \in A_p^+$  for all  $p > C$ , hence

$$\begin{aligned} & \frac{1}{|I|} \int_I w_0 w_1 \left( \frac{1}{|I|^{++}} \int_{I^{++}} (w_0 w_1)^{1-p'} \right)^{p-1} \\ & \leq \sup_I(w_1) \frac{1}{|I|} \int_I w_0 \left( \sup_{I^{++}} w_0^{1-p'} \right)^{p-1} \left( \frac{1}{|I|^{++}} \int_{I^{++}} w_1^{1-p'} \right)^{p-1} \\ & \leq C \frac{1}{|I|^+} \int_{I^+} w_1 \inf_{I^+}(w_0) \sup_{I^{++}}(w_0^{-1}) \left( \frac{1}{|I|^{++}} \int_{I^{++}} w_1^{1-p'} \right)^{p-1} \\ & \leq C \sup_{I^{++}}(w_0^{-1}) \frac{1}{|I^+|} \int_{I^+} w_0 \\ & \leq C \frac{1}{(\inf_{I^{++}} w_0)} \inf_{I^{++}} w_0 \leq C, \end{aligned}$$

and by Lemma 2.6 we have  $w \in A_p^+$  for all  $p > C$ .

To see this is the best range, we consider  $w(x) = 0$  if  $x \leq -1$ ,  $|x|^{C-1}$  if  $-1 < x \leq 0$  and 1 if  $x \geq 0$ .  $\square$

#### References

- [1] *H. Aimar, L. Forzani and F. J. Martín-Reyes*: On weighted inequalities for one-sided singular integrals. Proc. Amer. Math. Soc. *125* (1997), 2057–2064.
- [2] *D. Cruz-Uribe and C. J. Neugebauer*: The structure of reverse Hölder classes. Trans. Amer. Math. Soc. *347* (1995), 2941–2960.
- [3] *D. Cruz-Uribe, C. J. Neugebauer and V. Olesen*: The one-sided minimal operator and the one-sided reverse Hölder inequality. Stud. Math *116* (1995), 255–270.
- [4] *P. Guan and E. Sawyer*: Regularity estimates for oblique derivative problem. Anal. of Mathematics, Second Series *157* (1) (1993), 1–70.
- [5] *F. J. Martín-Reyes*: New proofs of weighted inequalities for the one-sided Hardy-Littlewood maximal functions. Proc. Amer. Math. Soc. *117*, 691–698.
- [6] *F. J. Martín-Reyes, P. Ortega and A. de la Torre*: Weighted inequalities for one-sided maximal functions. Trans. Amer. Math. Soc. *319* (2) (1990), 517–534.
- [7] *F. J. Martín-Reyes, L. Pick and A. de la Torre*:  $A_\infty^+$  condition. Canad. J. Math. *45* (6) (1993), 1231–1244.
- [8] *C. J. Neugebauer*: The precise range of indices for the  $RH_r$  and  $A_p$  weight classes. Preprint.
- [9] *E. Sawyer*: Weighted inequalities for the one-sided Hardy-Littlewood maximal function. Trans. Amer. Math. Soc. *297* (1986), 53–61.

*Author's address*: M. S. Riveros, FaMAF, Universidad Nacional de Córdoba, (5000) Córdoba, Argentina, e-mail: [sriveros@mate.uncor.edu](mailto:sriveros@mate.uncor.edu); A. de la Torre, Análisis Matemático, Facultad de Ciencias, Universidad de Málaga, (29071) Málaga, Spain, e-mail: [torre@anamat.cie.uma.es](mailto:torre@anamat.cie.uma.es).