

Sangwon Park

The general structure of inverse polynomial modules

Czechoslovak Mathematical Journal, Vol. 51 (2001), No. 2, 343–349

Persistent URL: <http://dml.cz/dmlcz/127651>

Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE GENERAL STRUCTURE OF INVERSE
POLYNOMIAL MODULES

SANGWON PARK, Pusan

(Received July 23, 1998)

Abstract. In this paper we compute injective, projective and flat dimensions of inverse polynomial modules as $R[x]$ -modules. We also generalize Hom and Ext functors of inverse polynomial modules to any submonoid but we show Tor functor of inverse polynomial modules can be generalized only for a symmetric submonoid.

Keywords: module, inverse polynomial, homological dimensions, Hom, Ext, Tor

MSC 2000: 16E30, 13C11, 16D80

1. INTRODUCTION

Northcott in [4] considered the module $K[x^{-1}]$ of inverse polynomials over the polynomial ring $K[x]$ (with K a field). The idea for this module came from Macaulay's work in [1]. McKerrow elaborated on Northcott's work (in [3]) and considered the module $M[x^{-1}]$ over $R[x]$ (with R a ring and M a left R -module). An easy modification of argument in Northcott [4] shows that if R is a left noetherian ring and E is an injective left $R[x]$ -module, then $E[x^{-1}]$ is an injective left $R[x]$ -module. In [5] and [6] we considered the behaviors of these so-call Macaulay-Northcott modules when we apply the torsion and extension functors to them. In this paper we will consider various homological dimensions of these modules. We also generalize Hom, Ext and Tor functors of Macaulay-Northcott modules to any submonoid. An interesting result is the fact that $\mathrm{Tor}_i^{R[x^S]}(M[x^{-S}], N[x^{-S}]) \cong \mathrm{Tor}_{i-1}^R(M, N)[x^{-S}]$ for a symmetric submonoid S of N (the set of natural numbers).

Definition 1.1. Let R be a ring and M be a left R -module, then $M[x^{-1}]$ is a left $R[x]$ -module such that

$$x(m_0 + m_1x^{-1} + \dots + m_nx^{-n}) = m_1 + m_2x^{-1} + \dots + m_nx^{-n+1}$$

and such that

$$r(m_0 + m_1x^{-1} + \dots + m_nx^{-n}) = rm_0 + rm_1x^{-1} + \dots + rm_nx^{-n}$$

where $r \in R$. We call $M[x^{-1}]$ a Macaulay-Northcott module.

Definition 1.2. Let \mathcal{C} be the category of left R -module and \mathcal{D} be the category of left $R[x]$ -module. Let $f: {}_R M \rightarrow {}_R N$ be a R -linear map, then $T: \mathcal{C} \rightarrow \mathcal{D}$ defined by $T(M) = M[x^{-1}]$ and $T(f) = f$ (where $f(m_0 + m_1x^{-1} + \dots + m_nx^{-n+1}) = f(m_0) + f(m_1)x^{-1} + \dots + f(m_n)x^{-1}$) is an additive and exact functor between \mathcal{C} and \mathcal{D} . We call T the Macaulay-Northcott functor.

2. THE GENERAL STRUCTURE OF INVERSE POLYNOMIAL MODULES

Lemma 2.1. Let M be an essential extension of N as a left R -module, then $M[x^{-1}]$ is an essential extension of $N[x^{-1}]$ as a left $R[x]$ -module.

Proof. Let $m_0 + m_1x^{-1} + \dots + m_ix^{-i} \in M[x^{-1}]$ w.l.o.g. (without loss of generality) let $m_i \neq 0$, then there is $r_i \in R$, $r_i \neq 0$ such that $m_ir_i \in N$, $m_ir_i \neq 0$. So $r_ix^i(m_0 + m_1x^{-1} + \dots + m_ix^{-i}) = r_im_i \in N[x^{-1}]$. Hence, $M[x^{-1}]$ is an essential extension of $N[x^{-1}]$. \square

Remark 2.2. Let R be a left noetherian ring. If E is an injective envelope of M then $E[x^{-1}]$ is an injective envelope of $M[x^{-1}]$.

Lemma 2.3. If $0 \rightarrow M \rightarrow E^0 \rightarrow E^1 \rightarrow \dots$ is a minimal injective resolution of M as a left R -module then

$$0 \rightarrow M[x^{-1}] \rightarrow E^0[x^{-1}] \rightarrow E^1[x^{-1}] \rightarrow \dots$$

is a minimal injective resolution of $M[x^{-1}]$ as a left $R[x]$ -module.

Proof. This follows from the lemma above and from the exactness of the Macaulay-Northcott functor. \square

Definiton 2.4 ([7]). Let $0 \rightarrow B \xrightarrow{\varepsilon} E^0 \xrightarrow{d^0} E^1 \rightarrow \dots$ be an injective resolution; denote $\text{im } \varepsilon$ by L^0 and, for $n \geq 1$, denote $\text{im } d^{n-1}$ by L^n . For $n \geq 0$, L^n is the n -th cosyzygy.

We will need the following results.

Lemma 2.5 (Theorem 9.8, [7]). *The followings are equivalent for a left R -module N :*

- (1) $\text{id}(N) \leq n$;
- (2) $\text{Ext}^k(M, N) = 0$ for all module M and all $k \geq n + 1$;
- (3) $\text{Ext}^{n+1}(M, N) = 0$ for all modules M ;
- (4) every injective resolution of a left R -module N has an injective $(n - 1)$ st cosyzygy.

Theorem 2.6. *Let R be a left Noetherian ring. Then*

$$\text{id}_{R[x]}(M[x^{-1}]) = \text{id}_R(M).$$

Proof. Suppose $\text{id}_R(M) = n$ and

$$0 \rightarrow M \rightarrow E^0 \rightarrow E^1 \rightarrow \dots \rightarrow E^n \rightarrow 0$$

is an injective resolution of M . Then

$$0 \rightarrow M[x^{-1}] \rightarrow E^0[x^{-1}] \rightarrow E^1[x^{-1}] \rightarrow \dots \rightarrow E^n[x^{-1}] \rightarrow 0$$

is an injective resolution of $M[x^{-1}]$. Let

$$K^n = \ker(E^{n-2} \rightarrow E^{n-1})$$

then K^n is an injective R -module but K^i is not an injective R -module for $i < n$. Thus $K^n[x^{-1}]$ is an injective $R[x]$ -module and $K^i[x^{-1}]$ is not an injective $R[x]$ -module for $i < n$. Therefore, $\text{id}_{R[x]}(M[x^{-1}]) = n$. Suppose $\text{id}_R(M) = \infty$ and

$$0 \rightarrow M \rightarrow E^0 \rightarrow E^1 \rightarrow \dots \rightarrow E^n \rightarrow \dots$$

is an injective resolution of M . Then

$$M[x^{-1}] \rightarrow E^0[x^{-1}] \rightarrow E^1[x^{-1}] \rightarrow \dots \rightarrow E^n[x^{-1}] \rightarrow \dots$$

is an injective resolution of $M[x^{-1}]$. Let

$$K^n = \ker(E^{n-2} \rightarrow E^{n-1})$$

then K^i is not an injective R -module for all i . Thus $K^i[x^{-1}]$ is not an injective $R[x]$ -module for all i . Therefore, $\text{id}_{R[x]}(M[x^{-1}]) = \infty$. Similarly if $\text{id}_{R[x]} M[x^{-1}] = n$, then $\text{id}_R(M) = n$ and if $\text{id}_{R[x]} M[x^{-1}] = \infty$, then $\text{id}_R(M) = \infty$. Hence, $\text{id}_{R[x]}(M[x^{-1}]) = \text{id}_R(M)$. \square

Lemma 2.7. *There is a short exact sequence of $R[x]$ -modules*

$$0 \rightarrow M[x] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-1}] \rightarrow 0.$$

Proof. Let M be a R -module, then $M[x] \subset M[x, x^{-1}]$ is a submodule. Let $\varphi: \frac{M[x, x^{-1}]}{M[x]} \rightarrow M[x^{-1}]$ be defined by

$$\begin{aligned} \varphi((a_0 + a_1x + \dots + a_nx^n + b_1x^{-1} + \dots + b_mx^{-m}) + M[x]) \\ = b_1 + b_2x^{-1} + \dots + b_mx^{-m+1}. \end{aligned}$$

Then φ is an isomorphism. Hence,

$$0 \rightarrow M[x] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-1}] \rightarrow 0$$

is a short exact sequence of $R[x]$ -modules. □

Theorem 2.8. *Let R be a left noetherian ring. Then*

$$\begin{aligned} n \leq \text{pd}_{R[x]}(M[x^{-1}]) \leq n+1 \quad \text{if} \quad \text{pd}_R(M) = n, \\ \text{pd}_{R[x]}(M[x^{-1}]) = \infty \quad \text{if} \quad \text{pd}_R(M) = \infty. \end{aligned}$$

Proof. Suppose $\text{pd}_R(M) = n$ then $\text{pd}_{R[x]}(M[x]) = n$. Consider the following exact sequence

$$0 \rightarrow M[x] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-1}] \rightarrow 0.$$

Then we have the long exact sequence with Ext

$$\begin{aligned} 0 \rightarrow \text{Hom}(M[x^{-1}], N) \rightarrow \text{Hom}(M[x, x^{-1}], N) \rightarrow \text{Hom}(M[x], N) \\ \rightarrow \text{Ext}^1(M[x^{-1}], N) \rightarrow \dots \rightarrow \text{Ext}^{n+1}(M[x^{-1}], N) \\ \rightarrow \text{Ext}^{n+1}(M[x, x^{-1}], N) \rightarrow \text{Ext}^{n+1}(M[x], N) \\ \rightarrow \text{Ext}^{n+2}(M[x^{-1}], N) \rightarrow \text{Ext}^{n+2}(M[x, x^{-1}], N) \rightarrow \dots \end{aligned}$$

for a left R -module N . So $\text{Ext}^{n+1}(M[x], N) = 0$. Since $\text{pd}_R(S^{-1}M) \leq \text{pd}_R(M) + 1$ and $M[x, x^{-1}] \cong S^{-1}M[x]$, where $S = \{1, s, s^2, \dots\}$ for $s \in R$,

$$\text{pd}_{R[x]}(M[x, x^{-1}]) \leq \text{pd}_{R[x]}(M[x]) + 1.$$

So $\text{Ext}^{n+2}(M[x, x^{-1}], N) = 0$. Thus $\text{Ext}^{n+2}(M[x^{-1}], N) = 0$. Therefore,

$$\text{pd}_{R[x]}(M[x^{-1}]) \leq n + 1.$$

Since $\text{Ext}^n(M[x^{-1}], N[x^{-1}]) \cong \text{Ext}^n(M, N)[[x]]$ (Theorem 1.2, [6]), and since $\text{pd}_R(M) = n$, $\text{Ext}^n(M, N)[[x]] \neq 0$, $\text{Ext}^n(M[x^{-1}], N[x^{-1}]) \neq 0$. Therefore,

$$\text{pd}_{R[x]}(M[x^{-1}]) \geq n.$$

Hence, we have

$$n \leq \text{pd}_{R[x]}(M[x^{-1}]) \leq n + 1.$$

Suppose $\text{pd}_R(M) = \infty$. Let $n \in \mathbb{Z}^+$ then $\text{Ext}^{n+1}(M, N) \neq 0$ for some N . But $\text{Ext}^{n+1}(M[x^{-1}], N[x^{-1}]) \cong \text{Ext}^{n+1}(M, N)[[x]]$. Therefore,

$$\text{Ext}^{n+1}(M[x^{-1}], N[x^{-1}]) \neq 0.$$

Hence, we have $\text{pd}_{R[x]}(M[x^{-1}]) = \infty$. □

Theorem 2.9. *Let R be a commutative and noetherian and M be a finitely generated left R -module. Then*

$$\text{pd}_{R[x]}(M[x^{-1}]) = n + 1 \quad \text{if} \quad \text{pd}_R(M) = n.$$

P r o o f. Suppose $\text{pd}_R(M) = n$ then $\text{Tor}_n^R(M, N) \neq 0$ for some finitely generated left R -module N by (II) p. 130 [2]. By Theorem 2.1 [6],

$$\text{Tor}_{n+1}^{R[x]}(M[x^{-1}], N[x^{-1}]) \cong \text{Tor}_n^R(M, N)[x^{-1}].$$

So $\text{Tor}_{n+1}^{R[x]}(M[x^{-1}], N[x^{-1}]) \neq 0$. Therefore, $\text{pd}_{R[x]}(M[x^{-1}]) > n$. Hence, by the previous theorem we conclude that $\text{pd}_{R[x]}(M[x^{-1}]) = n + 1$. □

Theorem 2.10. *If $\text{fd}_R(M) = n$ then $\text{fd}_{R[x]}(M[x^{-1}]) = n + 1$ and if $\text{fd}_R(M) = \infty$ then $\text{fd}_{R[x]}(M[x^{-1}]) = \infty$.*

P r o o f. Suppose $\text{fd}_R(M) = n$ then $\text{fd}_{R[x]}(M[x]) = n$. Consider the following exact sequence $0 \rightarrow M[x] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-1}] \rightarrow 0$. Then we have a long exact sequence with Tor

$$\begin{aligned} & \dots \rightarrow \text{Tor}_{n+2}(N, M[x^{-1}]) \rightarrow \text{Tor}_{n+1}(N, M[x]) \\ & \rightarrow \text{Tor}_{n+1}(N, M[x, x^{-1}]) \rightarrow \text{Tor}_{n+1}(N, M[x^{-1}]) \rightarrow \dots \rightarrow \text{Tor}_1(N, M[x]) \\ & \rightarrow N \otimes M[x] \rightarrow N \otimes M[x, x^{-1}] \rightarrow N \otimes M[x^{-1}] \rightarrow 0. \end{aligned}$$

Then $\text{Tor}_{n+1}(N, M[x]) = 0$. Since $M[x, x^{-1}] \cong S^{-1}M[x]$ where $S = \{1, s, s^2, \dots\}$, $s \in R$ and

$$\text{fd}_{R[x]}(M[x, x^{-1}]) \leq \text{fd}_{R[x]}(M[x]) + 1 = n + 1,$$

$\text{Tor}_{n+2}(N, M[x, x^{-1}]) = 0$. Therefore, $\text{Tor}_{n+2}(N, M[x^{-1}]) = 0$. Hence,

$$\text{fd}_{R[x]}(M[x^{-1}]) \leq n + 1.$$

Since $\text{fd}_R(M) = n$, $\text{Tor}_n(N, M) \neq 0$ for some N . Since

$$\text{Tor}_{n+1}(N[x^{-1}], M[x^{-1}]) \cong \text{Tor}_n(N, M)[x^{-1}],$$

$\text{Tor}_{n+1}(N[x^{-1}], M[x^{-1}]) \neq 0$. Thus $\text{fd}_{R[x]}(M[x^{-1}]) \geq n + 1$. Therefore, we have $\text{fd}_{R[x]}(M[x^{-1}]) = n + 1$. Suppose $\text{fd}_R(M) = \infty$. Let $n \in \mathbb{Z}^+$ then $\text{Tor}_{n+1}(N, M) \neq 0$ for some N . Since

$$\text{Tor}_{n+1}(N[x], M[x^{-1}]) \cong \text{Tor}_{n+1}(N, M)[x^{-1}],$$

$\text{Tor}_{n+1}(N[x], M[x^{-1}]) \neq 0$. Hence, we have $\text{fd}_{R[x]}(M[x^{-1}]) = \infty$. \square

Let $S \subset \mathbb{N}$ (\mathbb{N} the natural numbers) be a submonoid where we assume that for some n_0 , all $n \geq n_0$ are in \mathbb{N} . If S is a symmetric, then it has a conductor $c \in S$, i.e. c is such that the function $n \rightarrow -n + c - 1$ from \mathbb{Z} to \mathbb{Z} maps S bijectively to its complement in \mathbb{Z} .

If S is a submonoid of \mathbb{N} (\mathbb{N} is the set of all natural numbers), then $R[x^S]$ is defined to be the ring of all sums $\sum_{i \in S} r_i x^i$ with $r_i \in R$. If M is a left R -module, then we let $M[x^{-S}]$ consist of all $\sum_{i \in S} m_i x^{-i}$. Then $M[x^{-S}]$ is naturally an $R[x^S]$ -module (where $x^j \cdot x^{-i} = x^{-(i-j)}$ if $i - j \in S$ and $x^j \cdot x^{-i} = 0$ if $i - j \notin S$ for $i, j \in S$).

Theorem 2.11. *Let M be a left R -module and S be a symmetric submonoid of \mathbb{N} (the set of natural numbers). Then there is a short exact sequence*

$$0 \rightarrow M[x^S] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-S}] \rightarrow 0$$

as $R[x^S]$ -modules.

Proof. Let M be a R -module, then $M[x^S] \subset M[x, x^{-1}]$ is a submodule as $R[x^S]$ -modules. Let $\sum_{n \in \mathbb{Z}} a_n x^n = \dots + a_{-1} x^{-1} + a_0 + a_1 x^1 + a_2 x^2 + \dots$ for an element of $M[x, x^{-1}]$. Let $\varphi: M[x, x^{-1}]/M[x^S] \rightarrow M[x^{-S}]$ be defined by

$$\varphi\left(\sum_{n \in \mathbb{Z}} a_n x^n + M[x^S]\right) = \sum_{n \in S} a_{-n+c-1} x^{-n}$$

where c is the conductor of S . Then φ is an isomorphism. Hence,

$$0 \rightarrow M[x^S] \rightarrow M[x, x^{-1}] \rightarrow M[x^{-S}] \rightarrow 0$$

is a short exact sequence as $R[x^S]$ -module. \square

We note that for a nonsymmetric submonoid S , the above theorem does not hold.

Remark 2.12. We easily see that the natural isomorphism

$$\mathrm{Hom}_{R[x]}(M[x^{-1}], N[x^{-1}]) \cong \mathrm{Hom}_R(M, N)[[x]]$$

(Theorem 1.1, [6]) can be extended to the case

$$\mathrm{Hom}_{R[x^S]}(M[x^{-S}], N[x^{-S}]) \cong \mathrm{Hom}_R(M, N)[[x^S]]$$

for any submonoid $S \subset N$, and also for a left noetherian ring R the isomorphism

$$\mathrm{Ext}_{R[x]}^i(M[x^{-1}], N[x^{-1}]) \cong \mathrm{Ext}_R^i(M, N)[[x]]$$

(Theorem 1.2, [6]) can be extended to the case

$$\mathrm{Ext}_{R[x^S]}^i(M[x^{-S}], N[x^{-S}]) \cong \mathrm{Ext}_R^i(M, N)[[x^S]]$$

for any submonoid $S \subset N$. But by the above theorem 2.8, the isomorphism

$$\mathrm{Tor}_i^{R[x]}(M[x^{-1}], N[x^{-1}]) \cong \mathrm{Tor}_{i-1}^R(M, N)[x^{-1}]$$

(Theorem 2.1, [6]) can be extended to the case

$$\mathrm{Tor}_i^{R[x^S]}(M[x^{-S}], N[x^{-S}]) \cong \mathrm{Tor}_{i-1}^R(M, N)[x^{-S}]$$

only for a symmetric submonoid S .

References

- [1] *F. S. Macaulay*: The algebraic theory of modular system. Cambridge Tracts in Math. 19 (1916).
- [2] *H. Matsumura*: Commutative Algebra. W. A. Benjamin, Inc., New York, 1970.
- [3] *A. S. McKerrow*: On the injective dimension of modules of power series. Quart J. Math. Oxford Ser. (2), 25 (1974), 359–368.
- [4] *D. G. Northcott*: Injective envelopes and inverse polynomials. J. London Math. Soc. (2), 8 (1974), 290–296.
- [5] *S. Park*: Inverse polynomials and injective covers. Comm. Algebra 21 (1993), 4599–4613.
- [6] *S. Park*: The Macaulay-Northcott functor. Arch. Math. (Basel) 63 (1994), 225–230.
- [7] *J. Rotman*: An Introduction to Homological Algebra. Academic Press Inc., New York, 1979.

Author's address: Department of Mathematics, Dong-A University, Pusan, Korea 604-714, e-mail: swpark@mail.donga.ac.kr.