

Ján Jakubík

Torsion classes of Specker lattice ordered groups

Czechoslovak Mathematical Journal, Vol. 52 (2002), No. 3, 469–482

Persistent URL: <http://dml.cz/dmlcz/127736>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

TORSION CLASSES OF SPECKER LATTICE ORDERED GROUPS

JÁN JAKUBÍK, Košice

(Received June 3, 1999)

Abstract. In this paper we investigate the relations between torsion classes of Specker lattice ordered groups and torsion classes of generalized Boolean algebras.

Keywords: Specker lattice ordered group, generalized Boolean algebra, torsion class

MSC 2000: 06F15, 06E99

Specker lattice ordered groups have been investigated by Conrad and Darnel [3], [4], [5], and by Conrad and Martinez [7]. Below we write “Specker group” instead of “Specker lattice ordered group”.

The notion of a torsion class of lattice ordered groups was introduced by Martinez [13].

In [5] it was proved that the class \mathcal{S}_G of all Specker groups is a torsion class of lattice ordered groups.

Radical classes of generalized Boolean algebras were studied in [11]. Let Y be a radical class of generalized Boolean algebras; we define Y to be a torsion class of generalized Boolean algebras if it is closed with respect to homomorphic images.

We denote by \mathcal{T}^s the collection of all torsion classes X of lattice ordered groups such that $X \subseteq \mathcal{S}_G$. Further, let \mathcal{T}^b be the collection of all torsion classes of generalized Boolean algebras.

In the present paper we show that there exists a one-to-one mapping φ_0 of \mathcal{T}^s onto \mathcal{T}^b such that, whenever $X_1, X_2 \in \mathcal{T}^s$, then

$$X_1 \subseteq X_2 \Leftrightarrow \varphi_0(X_1) \subseteq \varphi_0(X_2).$$

Further, we prove that \mathcal{T}^s is a large collection (in the sense that there exists an injective mapping of the class of all infinite cardinals into \mathcal{T}^s).

1. PRELIMINARIES

For the sake of completeness we recall some relevant definitions.

We denote by \mathcal{G} the class of all lattice ordered groups. Let $G \in \mathcal{G}$. An element $0 < s \in G$ is called singular if $x \wedge (s - x) = 0$ whenever $0 \leq x \leq s$. The set of all singular elements of G is denoted by $S(G)$; further, we put $S_0(G) = S(G) \cup \{0\}$. Then $S_0(G)$ is a sublattice of the lattice G^+ .

A lattice ordered group G is a Specker group if G is generated as a group by the set $S(G)$. (Cf. [6].)

For each $G \in \mathcal{G}$ let $C(G)$ be the system of all convex ℓ -subgroups of G ; this system is partially ordered by the set-theoretical inclusion. Then $C(G)$ is a complete lattice.

A torsion class of lattice ordered groups is defined to be a nonempty subclass X of \mathcal{G} such that

- (i) X is closed with respect to homomorphisms;
- (ii) if $G_1 \in X$ and $G_2 \in C(G_1)$, then $G_2 \in X$;
- (iii) if $G \in \mathcal{G}$ and $\{G_i\}_{i \in I} \subseteq C(G) \cap X$, then $\bigvee_{i \in I} G_i \in X$.

If X is a nonempty subclass of \mathcal{G} which is closed with respect to isomorphisms and satisfies the conditions (ii), (iii), then X is called a radical class of lattice ordered groups (cf. [10]).

A lattice L is a generalized Boolean algebra if it has the least element 0 and if for each $x \in L$, the interval $[0, x]$ of L is a Boolean algebra.

Let \mathcal{B} be the class of all generalized Boolean algebras. For each $B \in \mathcal{B}$, the system $J(B)$ of all ideals of B (partially ordered by the set-theoretical inclusion) is a complete lattice.

The torsion class of generalized Boolean algebras is defined by conditions which are analogous to the conditions (i), (ii), (iii) above with the distinction that G and $C(G_1)$ are replaced by B and $J(B_1)$.

A nonempty subclass Y of \mathcal{B} which is closed with respect to isomorphisms and satisfies the conditions analogous to (ii) and (iii) (in the above specified sense) is called a radical class of generalized Boolean algebras.

2. AUXILIARY RESULTS

For lattice ordered groups we apply the notation as in Birkhoff [1] and Conrad [2].

It is well-known that an element $0 < s$ of a lattice ordered group G belongs to $S(G)$ if and only if the interval $[0, s]$ of G is a Boolean algebra.

The following lemma is easy to verify (cf. also [5]).

Lemma 2.1. *Let G be a Specker group. Then $S_0(G)$ is a generalized Boolean algebra.*

The following result is known (cf. [4], Proposition 2.6). Let us remark that a simple alternative proof of 2.2 can be performed by applying Carathéodory functions (for this notion, cf. Gofman [8] and the author [9], [12]).

Lemma 2.2. *Let B be a generalized Boolean algebra. There exists a Specker group G such that $B = S_0(G)$.*

Let $G \in \mathcal{G}$. An indexed system $(a_i)_{i \in I}$ of elements of G^+ is called disjoint if $a_{i(1)} \wedge a_{i(2)} = 0$ whenever $i(1)$ and $i(2)$ are distinct elements of I .

Let Z be the additive group of all integers. If $G \in \mathcal{G}$, $x \in G$ and if 0 is the neutral element of Z , then we define $0x$ to be the neutral element of G . (We do not distinguish typographically the neutral element of G and the neutral element of Z ; from the context it will be clear which of these elements is taken into consideration.)

From [5], Proposition 1.2 we obtain

Lemma 2.3. *The following conditions for G are equivalent:*

- (i) G is a Specker group.
- (ii) For each $0 \neq x \in G$ there exist a disjoint system (x_i) ($i = 1, 2, \dots, n$) of elements of $S(G)$ and integers α_i ($i = 1, 2, \dots, n$) such that

$$(1) \quad x = \alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n.$$

Under the notation as in 2.3 we say that (1) is a representation of the element x . A simple calculation yields

Lemma 2.4. *Let (1) be a representation of an element x of a Specker group G . Then $x > 0$ if and only if $\alpha_i > 0$ for $i = 1, 2, \dots, n$. Further, $x \in S(G)$ if and only if $\alpha_i \in \{0, 1\}$ for each $i = 1, 2, \dots, n$ and if there is $i \in \{1, 2, \dots, n\}$ with $\alpha_i \neq 0$.*

Let us extend the definition of the representation so that for $x = 0$ we consider (1) to be a representation of x if $\alpha_i = 0$ ($i = 1, 2, \dots, n$). Moreover, for any $x \in G$ the relation $x_i = 0$ for some $i \in \{1, 2, \dots, n\}$ will be also allowed.

Lemma 2.5. *Let x and y be elements of a Specker group G . Then there exist $t_1, t_2, \dots, t_k \in S(G)$ and integers $\gamma_1, \gamma_2, \dots, \gamma_k, \gamma'_1, \gamma'_2, \dots, \gamma'_k$ such that x and y have representations*

$$\begin{aligned} x &= \gamma_1 t_1 + \gamma_2 t_2 + \dots + \gamma_k t_k, \\ y &= \gamma'_1 t_1 + \gamma'_2 t_2 + \dots + \gamma'_k t_k. \end{aligned}$$

Proof. In view of 2.3 there exist representations

$$\begin{aligned}x &= \alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n, \\y &= \beta_1 y_1 + \beta_2 y_2 + \dots + \beta_m y_m.\end{aligned}$$

Denote

$$v = \left(\bigvee x_i\right) \vee \left(\bigvee y_j\right)$$

with $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m$. Hence v belongs to $S_0(G)$. Thus $[0, v]$ is a Boolean algebra.

Let x_{n+1} be the complement of the element $x_1 \vee x_2 \vee \dots \vee x_n$ in the Boolean algebra $[0, v]$ and let y_{m+1} be defined analogously. We put

$$I = \{1, 2, \dots, n+1\}, \quad J = \{1, 2, \dots, m+1\}, \quad \alpha_{n+1} = 0 = \beta_{m+1}.$$

Hence we have representations

$$x = \sum_{i \in I} \alpha_i x_i, \quad y = \sum_{j \in J} \beta_j y_j.$$

Denote $x_i \wedge y_j = z_{ij}$ for $i \in I$ and $j \in J$. We obtain

$$v = \bigvee_{i \in I} x_i = \bigvee_{j \in J} y_j,$$

whence for each $i \in I$ we have

$$\begin{aligned}x_i &= x_i \wedge v = x_i \wedge \left(\bigvee_{j \in J} y_j\right) = \bigvee_{j \in J} (x_i \wedge y_j) \\&= \bigvee_{j \in J} z_{ij} = \sum_{j \in J} z_{ij},\end{aligned}$$

since the indexed system $(z_{ij})_{j \in J}$ is disjoint. Analogously,

$$y_j = \sum_{i \in I} z_{ij}.$$

Therefore we get

$$(2) \quad x = \sum_{i \in I, j \in J} \alpha_i z_{ij},$$

$$(3) \quad y = \sum_{i \in I, j \in J} \beta_i z_{ij},$$

and (2), (3) are representations of x and y , respectively. This completes the proof. \square

Let B be a generalized Boolean algebra and let A be an ideal of B . In view of 2.2 there exists a Specker group G such that $S_0(G) = B$. We denote by G_1 the set of all $x \in G$ for which there exists a representation (1) such that $x_i \in A$ for $i = 1, 2, \dots, n$.

Lemma 2.6. G_1 is a convex ℓ -subgroup of G and $S_0(G_1) = A$.

Proof. a) Let $0 < x \in G_1$ and $y \in G$, $0 < y \leq x$. We can assume that (1) is a representation of x and that $x_i \in A$ for $i = 1, 2, \dots, n$. Further, in view of 2.4, $\alpha_i \geq 0$ for $i = 1, 2, \dots, n$.

There exists a representation of the element y having the form

$$y = \beta_1 y_1 + \beta_2 y_2 + \dots + \beta_m y_m$$

with $y_j \in S(G)$ and $\beta_j > 0$ for $j = 1, 2, \dots, m$.

Consider the element y_1 . Put

$$\begin{aligned} x_0 &= x_1 + x_2 + \dots + x_n = \vee x_i \quad (i = 1, 2, \dots, n), \\ k &= \max\{\alpha_1, \alpha_2, \dots, \alpha_n\}. \end{aligned}$$

We have $y_1 \leq x$, whence $y_1 \leq kx_0$. Further, $x_0 \in A$.

In view of the Riesz decomposition property (cf., e.g., Conrad [2], p. 0,19) there are $y_{11}, y_{12}, \dots, y_{1k} \in G$ such that $0 \leq y_{1t} \leq x_0$ for $t = 1, 2, \dots, k$ and

$$y_1 = y_{11} + y_{12} + \dots + y_{1k}.$$

The relation $y_1 \in S(G)$ yields that the system (y_{1t}) ($t = 1, 2, \dots, k$) is disjoint. Thus

$$y_1 = y_{11} \vee y_{12} \vee \dots \vee y_{1k} \leq x_0.$$

Hence $y_1 \in A$. Similarly, $y_2, \dots, y_m \in A$. We conclude that $y \in G_1$.

b) If $x \in G$ and if x has a representation (1), then $-x$ has the representation

$$-x = (-\alpha_1)x_1 + \dots + (-\alpha_n)x_n.$$

Hence if x belongs to G_1 , then $-x$ belongs to G_1 as well.

c) Let x and y be elements of G_1 . We apply the same notation as in the proof of 2.5 and we can assume that all x_i and all y_j belong to A . Then all z_{ij} belong to A .

From (2) and (3) we conclude that $x + y$ has the representation

$$x + y = \sum_{i \in I, j \in J} (\alpha_i + \beta_j) z_{ij}.$$

Thus $x + y \in G_1$.

Let $i \in I$ and $j \in J$. Put

$$\gamma_{ij} = \min\{\alpha_i, \beta_j\}, \quad \delta_{ij} = \max\{\alpha_i, \beta_j\}.$$

Then

$$\begin{aligned} x \wedge y &= \sum_{i \in I, j \in J} \gamma_{ij} z_{ij}, \\ x \vee y &= \sum_{i \in I, j \in J} \delta_{ij} z_{ij}, \end{aligned}$$

whence $x \wedge y, x \vee y \in G_1$. Therefore in view of b), G_1 is an ℓ -subgroup of G . This fact and a) yield that G_1 is a convex ℓ -subgroup of G .

d) The relation $A \subseteq S_0(G_1)$ is obviously valid. Let $0 < x \in S_0(G_1)$. We apply the notation as above. We can assume that $x_i > 0$ and $\alpha_i > 0$ for $i = 1, 2, \dots, n$.

Since x is singular, the system $(\alpha_i x_i)$ ($i = 1, 2, \dots, n$) is disjoint and all $\alpha_i x_i$ belong to $S(G)$. In view of 2.4, $\alpha_i = 1$ for $i = 1, 2, \dots, n$. Hence

$$x = x_1 + x_2 + \dots + x_n = x_1 \vee x_2 \vee \dots \vee x_n \in A.$$

□

The following result is well-known.

Lemma 2.7. *Let $G \in \mathcal{G}$ and let $\{G_i\}_{i \in I}$ be a nonempty system of elements of $C(G)$. Put $H = \bigvee_{i \in I} G_i$. Then H is the set of all elements $h \in G$ which can be expressed in the form*

$$h = h_1 + h_2 + \dots + h_n,$$

where $h_j \in \bigcup_{i \in I} G_i$ for each $j \in \{1, 2, \dots, n\}$. If $h > 0$, then there are h_i with the mentioned property such that $h_j > 0$ for $j = 1, 2, \dots, n$.

Lemma 2.8. *Let $B \in \mathcal{B}$ and let $\{A_i\}_{i \in I}$ be a nonempty system of elements of $J(B)$. Put $A = \bigvee_{i \in I} A_i$. Then A is the set of all elements $a \in B$ which can be expressed in the form*

$$a = a_1 \vee a_2 \vee \dots \vee a_n,$$

where $a_j \in \bigcup_{i \in I} A_i$ for each $j \in \{1, 2, \dots, n\}$.

The proof is simple and will be omitted.

Lemma 2.9. Let $G \in \mathcal{G}$, $G_i \in C(G)$ ($i \in I$), $\bigvee_{i \in I} G_i = H$, $B_i = S_0(G_i)$. Then $S_0(H) = \bigvee_{i \in I} B_i$, where $\bigvee_{i \in I} B_i$ is taken with respect to the lattice $J(S_0(G))$.

P r o o f. Let $i \in I$. We have $G_i \in C(G)$. From this relation we infer that

$$S_0(G_i) = S_0(H) \cap G_i,$$

whence $B_i \subseteq S_0(H)$. From this and from the fact that $S_0(H)$ is an ideal of $S_0(G)$ we obtain

$$\bigvee_{i \in I} B_i \subseteq S_0(H).$$

Let $0 < h \in S_0(H)$. Then in view of 2.7 there are $h_1, h_2, \dots, h_k \in \bigcup_{i \in I} G_i$ such that $0 < h_t$ ($t = 1, 2, \dots, k$) and $h = h_1 + h_2 + \dots + h_k$. Since h is singular in G all h_1, h_2, \dots, h_k are singular in G , hence for each $t \in \{1, 2, \dots, k\}$ there is $i(t) \in I$ such that $h_t \in S_0(G_{i(t)}) = B_{i(t)}$. Moreover, the system $(h_t)_{t=1,2,\dots,k}$ is disjoint. Thus

$$h = h_1 \vee h_2 \vee \dots \vee h_k.$$

Therefore $h \in \bigvee_{i \in I} B_i$. □

3. THE MAPPING φ

For each Specker group G we put

$$\varphi(G) = S_0(G).$$

From 2.3 we conclude

Lemma 3.1. If G_1 and G_2 are Specker groups such that $\varphi(G_1)$ is isomorphic to $\varphi(G_2)$, then G_1 and G_2 are isomorphic.

Let K^s and K^b be the collection of all nonempty classes of Specker groups or of generalized Boolean algebras, respectively. For each $X \in K^s$ we put

$$(1) \quad \varphi(X) = \{\varphi(G) : G \in X\} = Y.$$

Hence φ is a mapping of K^s into K^b .

Lemma 3.1.1. φ is a one-to-one mapping of K^s onto K^b such that, if $X_1, X_2 \in K^s$, then

$$(2) \quad X_1 \subseteq X_2 \Leftrightarrow \varphi(X_1) \subseteq \varphi(X_2).$$

Proof. In view of 2.2, φ is an epimorphism, and according to 3.1, φ is a monomorphism. The validity of (2) is then obvious. \square

Let X and Y be as in (1). If X satisfies the condition (ii) from Section 1 then we say that it is closed with respect to convex ℓ -subgroups. Under the analogous assumption on Y we say that Y is closed with respect to ideals.

If the condition (iii) from Section 1 is fulfilled for X then X is said to be closed under joins; the same term will be applied for Y under the analogous assumption.

Lemma 3.2. Let $X \in K^s$, $Y = \varphi(X)$. Then the following conditions are equivalent:

- (i) X is closed with respect to convex ℓ -subgroups;
- (ii) Y is closed with respect to ideals.

Proof. a) Assume that (i) is valid. Let $B \in Y$ and let A be an ideal of B . There exists $G \in X$ with $B = S_0(G)$. Let G_1 be as in 2.6. Then $G_1 \in X$, hence $S_0(G_1) \in Y$. In view of 2.6, $S_0(G_1) = A$. Therefore (ii) holds.

b) Suppose that the condition (ii) is satisfied. Let $G \in X$ and $G_1 \in C(X)$. Then

$$S_0(G_1) = S_0(G) \cap G_1,$$

whence $S_0(G_1)$ is an ideal of $S_0(G)$. Thus $S_0(G_1)$ belongs to Y . Therefore G_1 is an element of X . This yields that the condition (i) is valid. \square

The following assertions 3.3 and 3.4 slightly sharpen some results of [12], Section 3; in fact, several steps in the proofs are the same.

Lemma 3.3. Let $X \in K^s$ be closed with respect to convex ℓ -subgroups, $Y = \varphi(X)$. Then the following conditions are equivalent:

- (i) X is closed under joins;
- (ii) Y is closed under joins.

Proof. a) Let (i) be valid. Let B be a generalized Boolean algebra and let $\{B_i\}_{i \in I}$ be a nonempty subset of $J(B)$ such that all B_i belong to Y .

In view of 2.2 there is a Specker group G such that $S_0(G) = B$. Further, according to 2.9, for each B_i there is $G_i \in C(G)$ with $S_0(B_i) = G_i$. Hence $G_i \in X$ for each

$i \in I$. The condition (i) yields that the lattice ordered group $H = \bigvee_{i \in I} G_i$ belongs to X . Now from 2.9 we conclude that $\bigvee_{i \in I} B_i$ is an element of Y . Hence (ii) holds.

b) Assume that (ii) is satisfied. Let G be a lattice ordered group and let $\emptyset \neq \{G_i\}_{i \in I} \subseteq C(G)$ such that $G_i \in X$ for each $i \in I$. Then $B_i = S_0(G_i) \in Y$ for each $i \in I$. Put $B = S_0(G)$. We have $\{B_i\}_{i \in I} \subseteq J(B)$. In view of (ii), $\bigvee_{i \in I} B_i \in Y$. Thus 2.9 yields that $\bigvee_{i \in I} G_i$ belongs to X . Hence (i) holds. \square

Corollary 3.4. *Let X and Y be as in 3.2. The following conditions are equivalent:*

- (i) X is a radical class of lattice ordered groups;
- (ii) Y is a radical class of generalized Boolean algebras.

We denote by \mathcal{R}^s the collection of all radical classes X of lattice ordered groups such that $X \subseteq \mathcal{S}_G$. Further, let \mathcal{R}^b be the collection of all radical classes of generalized Boolean algebras. Let φ_0 be the mapping φ reduced to the collection \mathcal{R}^s .

Lemma 3.5. *φ_0 is a one-to-one mapping of \mathcal{R}^s onto \mathcal{R}^b such that if $X_1, X_2 \in \mathcal{R}^s$, then*

$$X_1 \subseteq X_2 \Leftrightarrow \varphi_0(X_1) \subseteq \varphi_0(X_2).$$

Proof. This is a consequence of 3.1.1 and 3.4. \square

4. TORSION CLASSES

Let G be a Specker group. Put $B = S_0(G)$. Let A be an ideal in B and let G_1 be as in 2.6. In view of 2.6 we have $S_0(G_1) = A$. If G_2 is another convex ℓ -subgroup of G and if $S_0(G_2) = A$, then $G_2 = G_1$. In fact, all elements of G_2 are linear combinations with integral coefficients of elements of $A \subseteq G_1$, whence $G_2 \subseteq G_1$; similarly, $G_1 \subseteq G_2$. Hence there is a one-to-one correspondence ψ between the elements of $C(G)$ and the elements of $J(B)$; this correspondence is given by

$$\psi(H) = S_0(H),$$

where H runs over $C(G)$.

Since G is abelian, each element $H \in C(G)$ is an ℓ -ideal of G and thus it is a kernel of a congruence ϱ_1 on G , and each congruence on G can be constructed in this way.

Similarly, each ideal A of B is kernel of a congruence on the generalized Boolean algebra B and in this way we obtain all congruences on B .

Let G_1 and A be as above. Let us construct the factor lattice ordered group $\overline{G} = G/G_1$ and the factor generalized Boolean algebra $\overline{B} = B(A)$.

For $g \in G$ and $b \in B$ we put

$$\begin{aligned}\overline{g} &= g + G_1 = \{g_1 \in G: g_1 \varrho_1 g\}, \\ \tilde{b} &= \{b_1 \in B: b_1 \varrho_2 b\},\end{aligned}$$

where ϱ_1 (and ϱ_2) is the congruence relation on G generated by G_1 (or the congruence relation on B generated by A , respectively).

Lemma 4.1. *Let $b, b_1 \in B$. Then $b \varrho_2 b_1$ if and only if $b \varrho_1 b_1$.*

P r o o f. Let $b \varrho_2 b_1$. Denote

$$u = b \wedge b_1, \quad v = b \vee b_1.$$

Then $u \varrho_2 v$. Let u' be the complement of u in the interval $[0, v]$. Thus

$$u' = u' \wedge v, \quad 0 = u' \wedge u,$$

whence $0 \varrho_2 u'$. It is well-known that the kernel of ϱ_2 is the ideal A of B . Thus $u' \in A$ and so $0 \varrho_1 u'$. Since

$$u = 0 \vee u, \quad v = u' \vee u$$

we obtain $u \varrho_1 v$ and this yields $b \varrho_1 b_1$.

Conversely, suppose that $b \varrho_1 b_1$. Then by analogous steps as above (with ϱ_1 and ϱ_2 interchanged) we conclude that $b \varrho_2 b_1$. \square

Lemma 4.2. *Let $x \in G$. The following conditions are equivalent:*

- (i) $\overline{x} \in S_0(\overline{G})$;
- (ii) *there exists $x_1 \in \overline{x}$ such that $x_1 \in S_0(G)$.*

P r o o f. The implication (ii) \Rightarrow (i) is obvious. Assume that (i) is valid.

The case $\overline{x} = \overline{0}$ is trivial. Suppose that $\overline{x} > \overline{0}$. Then without loss of generality we can assume that $x > 0$. Hence there exist $s_1, s_2, \dots, s_k \in S(G)$ and positive integers n_1, n_2, \dots, n_k such that

$$(1) \quad x = n_1 s_1 + n_2 s_2 + \dots + n_k s_k$$

and, moreover, the set $\{s_1, s_2, \dots, s_k\}$ is disjoint.

If all s_i belong to G_1 then $\bar{x} = \bar{0}$, which is a contradiction. Then the elements s_i belonging to G_1 can be omitted in (1); assume that $s_1, s_2, \dots, s_m \notin G_1$ and $s_{m+1}, \dots, s_k \in G_1$. Put

$$x_1 = n_1 s_1 + n_2 s_2 + \dots + n_m s_m.$$

We obtain $x_1 \in G_1$ and $0 < x_1 \in \bar{x}$.

Assume that $s_i \notin G_1$ and $n_i \geq 2$ for some $i \in 1, 2, \dots, m$. Then $\bar{0} < 2\bar{s}_i \leq \bar{x}$, whence $2\bar{s}_i \in S_0(\bar{G})$, but $\bar{0} < \bar{s}_i = \bar{s}_i \wedge \bar{s}_i = \bar{s}_i \wedge (2\bar{s}_i - \bar{s}_i)$, hence $2\bar{s}_i$ fails to be singular. Then $2\bar{s}_i \notin S_0(\bar{G})$, which is a contradiction. Thus $n_i = 1$ and hence x_1 can be written in the form

$$x_1 = s_1 \vee s_2 \vee \dots \vee s_m$$

with $s_1, s_2, \dots, s_m \in S_0(G)$. Therefore $x_1 \in S_0(G)$. □

Let f be a mapping of $S_0(\bar{G})$ into $B/A = \bar{B}$ which is defined as follows. For $\bar{x} \in S_0(\bar{G})$ we put

$$f(\bar{x}) = \tilde{x}_1,$$

where x_1 is as in 4.2.

If $x_1, x_2 \in B$ and both x_1 and x_2 satisfy the condition (ii) from 4.2, then $\bar{x}_1 = \bar{x} = \bar{x}_2$. Thus in view of 4.1 we obtain $\tilde{x}_1 = \tilde{x}_2$. Hence the mapping f is correctly defined.

Let $\tilde{z} \in \tilde{B}$. Then $f(\bar{z}) = \tilde{z}$, hence f is an epimorphism. Suppose that $\bar{x}, \bar{y} \in S_0(\bar{G})$ and $f(\bar{x}) = f(\bar{y})$. In other words, we have $f(\bar{x}) = \tilde{x}_1, f(\bar{y}) = \tilde{y}_1$ and $\tilde{x}_1 = \tilde{y}_1$. Then 4.1 yields that $\bar{x}_1 = \bar{y}_1$. Since $\bar{x}_1 = \bar{x}$ and $\bar{y}_1 = \bar{y}$ we get $\bar{x} = \bar{y}$. Therefore f is a monomorphism.

Further, in view of 4.1 we conclude that the mapping f is regular with respect to the lattice operations (i.e., if $x, y \in S_0(\bar{G})$, then $f(x \vee y) = f(x) \vee f(y)$, and similarly for the operation \wedge). Thus we have

Lemma 4.3. *f is an isomorphism of the generalized Boolean algebra $S_0(\bar{G})$ onto the generalized Boolean algebra \bar{B} .*

Now let X be a nonempty class of Specker groups and $Y = \varphi(X)$, where φ is as in Section 3.

Lemma 4.4. *The following conditions are equivalent:*

- (i) X is closed with respect to homomorphic images.
- (ii) Y is closed with respect to homomorphic images.

Proof. a) Assume that (i) is valid. Let $B \in Y$ and let A be an element of $J(B)$. We have to verify that $\overline{B} = B/A$ belongs to Y .

There exists $G \in X$ with $\varphi(G) = B$. Let G_1 be as in 2.6. In view of (i) we have $G/G_1 = \overline{G} \in X$. Hence $S_0(\overline{G}) \in Y$. According to 4.3, $S_0(\overline{G})$ is isomorphic to \overline{B} . Therefore $\overline{B} \in Y$.

b) Conversely, suppose that (ii) holds. Let $G \in X$ and $G_1 \in C(G)$. We have to verify that $\overline{G} = G/G_1$ belongs to X .

Denote $B = S_0(G)$. Thus $\varphi(G) = S_0(G) = B \in Y$. According to 4.3, $S_0(\overline{G})$ is a homomorphic image of B and hence, in view of (ii), $S_0(\overline{G})$ belongs to Y . Since

$$\varphi(\overline{G}) = S_0(\overline{G})$$

we obtain that \overline{G} must belong to X . □

Let \mathcal{T}^s and \mathcal{T}^b be the collection of all torsion classes of Specker groups or the collection of all torsion classes of generalized Boolean algebras, respectively.

Further, let φ_1 be the mapping φ reduced to the collection \mathcal{T}^s .

From 3.5 and 4.4 we conclude

Theorem 4.5. φ_1 is a one-to-one mapping of \mathcal{T}^s onto \mathcal{T}^b such that for $X_1, X_2 \in \mathcal{T}^s$ we have

$$X_1 \subseteq X_2 \Leftrightarrow \varphi_1(X_1) \subseteq \varphi_1(X_2).$$

Let K be the class of all infinite cardinals. For each $\alpha \in K$ we denote by $\mathcal{A}(\alpha)$ the class of all generalized Boolean algebras B such that, whenever $[x, y]$ is an interval of B , then

$$\text{card}[x, y] \leq \alpha.$$

It is obvious that in the definition of $\mathcal{A}(\alpha)$ it suffices to take into account the intervals $[x, y]$ with $x = 0$.

Lemma 4.6. Let $\alpha \in K$. Then $\mathcal{A}(\alpha)$ is a radical class of generalized Boolean algebras.

Proof. In view of the definition, $\mathcal{A}(\alpha)$ is closed with respect to ideals. It remains to verify that it is closed with respect to joins. Let $B \in \mathcal{B}$ and $\{A_i\}_{i \in I}$ be as in 2.8. Suppose that all A_i belong to $\mathcal{A}(\alpha)$. We apply the notation as in 2.8. We have to show that A belongs to $\mathcal{A}(\alpha)$.

Let $a \in B$. In view of 2.8 and according to Lemma 3.1 from [11] we infer that the element a can be written in the form

$$a = y_1 \vee y_2 \vee \dots \vee y_n$$

such that $y_j \in \bigcup A_i$ ($i \in I$) for each $j = 1, 2, \dots, n$, and $y_{j(1)} \wedge y_{j(2)} = 0$ whenever $j(1)$ and $j(2)$ are distinct elements of the set $\{1, 2, \dots, n\}$.

Then Lemma 3.2 of [11] yields that

$$[0, a] \simeq [0, y_1] \times [0, y_2] \times \dots \times [0, y_n],$$

whence $\text{card}[0, a] \leq \alpha$ and thus $A \in \mathcal{A}(\alpha)$. □

Lemma 4.7. *Let $\alpha \in K$, $B \in \mathcal{B}$ and let $[x, y]$ be an interval of B with $\text{card}[x, y] \leq \alpha$. Let A be an ideal of B ; put $\overline{B} = B/A$. For $z \in B$ let \overline{z} be the class in B/A containing the element z . Then $\text{card}[\overline{x}, \overline{y}] \leq \alpha$.*

Proof. Consider the mapping $f = [x, y] \rightarrow [\overline{x}, \overline{y}]$ defined by $f(z) = \overline{z}$ for each $z \in [x, y]$. Let $t \in B$, $\overline{t} \in [\overline{x}, \overline{y}]$. Put $t_1 = (t \vee x) \wedge y$. Then $t_1 \in [x, y]$ and $\overline{t_1} = \overline{t}$, whence $f(t_1) = \overline{t}$. Therefore the mapping f is surjective. Thus $\text{card}[\overline{x}, \overline{y}] \leq \text{card}[x, y] \leq \alpha$. □

From 4.6 and 4.7 we conclude

Proposition 4.8. *Let $\alpha \in K$. Then $\mathcal{A}(\alpha)$ is a torsion class of generalized Boolean algebras.*

For $\alpha \in K$ let B_α be the free Boolean algebra with α free generators. Then

- (i) $B_\alpha \in \mathcal{A}(\alpha)$;
- (ii) if $\beta \in K$ and $\beta > \alpha$, then $B_\beta \notin \mathcal{A}(\alpha)$.

We put $f_1(\alpha) = \mathcal{A}(\alpha)$ for each $\alpha \in K$. In view of 4.8, (i) and (ii) we have

Lemma 4.9. *f_1 is an injective mapping of the class K into \mathcal{T}^b .*

Let φ_1 be as in 4.5. For each $\alpha \in K$ we set

$$f_2(\alpha) = \varphi_1^{-1}(f_1(\alpha)).$$

From 4.5 and 4.9 we obtain

Theorem 4.10. *f_2 is an injective mapping of the class K into \mathcal{T}^s .*

References

- [1] *G. Birkhoff*: Lattice Theory. Third Edition, Providence, 1967.
- [2] *P. Conrad*: Lattice Ordered Groups. Tulane University, 1970.
- [3] *P. Conrad and M. R. Darnel*: Lattice-ordered groups whose lattices determine their additions. *Trans. Amer. Math. Soc.* 330 (1992), 575–598.
- [4] *P. F. Conrad and M. R. Darnel*: Generalized Boolean algebras in lattice-ordered groups. *Order* 14 (1998), 295–319.
- [5] *P. F. Conrad and M. R. Darnel*: Subgroups and hulls of Specker lattice-ordered groups. *Czechoslovak Math. J* 51 (2001), 395–413.
- [6] *P. F. Conrad and D. McAlister*: The completion of an ℓ -group. *J. Austral. Math. Soc.* 9 (1969), 182–208.
- [7] *P. F. Conrad and J. Martinez*: Signatures and S -discrete lattice ordered groups. *Algebra Universalis* 29 (1992), 521–545.
- [8] *C. Gofman*: Remarks on lattice ordered groups and vector lattices. I. Carathéodory functions. *Trans. Amer. Math. Soc.* 88 (1958), 107–120.
- [9] *J. Jakubík*: Cardinal properties of lattice ordered groups. *Fund. Math.* 74 (1972), 85–98.
- [10] *J. Jakubík*: Radical mappings and radical classes of lattice ordered groups. *Symposia Math.* 21. Academic Press, New York-London, 1977, pp. 451–477.
- [11] *J. Jakubík*: Radical classes of generalized Boolean algebras. *Czechoslovak Math. J.* 48 (1998), 253–268.
- [12] *J. Jakubík*: Radical classes of complete lattice ordered groups. *Math. Slovaca* 49 (1999), 417–424.
- [13] *J. Martinez*: Torsion theory for lattice ordered groups. *Czechoslovak Math. J.* 25 (1975), 284–299.

Author's address: Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia, e-mail: kstefan@saske.sk.