

Angelina Y. M. Chin

A note on semilocal group rings

Czechoslovak Mathematical Journal, Vol. 52 (2002), No. 4, 749–755

Persistent URL: <http://dml.cz/dmlcz/127761>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON SEMILOCAL GROUP RINGS

ANGELINA Y. M. CHIN, Kuala Lumpur

(Received October 27, 1999)

Abstract. Let R be an associative ring with identity and let $J(R)$ denote the Jacobson radical of R . R is said to be *semilocal* if $R/J(R)$ is Artinian. In this paper we give necessary and sufficient conditions for the group ring RG , where G is an abelian group, to be semilocal.

Keywords: semilocal, group ring

MSC 2000: 16L30

1. INTRODUCTION

All rings considered in this paper are associative with identity. Given a ring R and a group G , we will denote the group ring of G over R by RG . If H is a subgroup of G then ωH will denote the right ideal of RG generated by $\{1-h \mid h \in H\}$. In particular, if H is a normal subgroup of G then ωH is an ideal of RG and $RG/\omega H \cong R(G/H)$. If $H = G$, then ωG is called the *augmentation ideal* of RG and is written as Δ . It is well-known that $R \cong RG/\Delta$. If I is an ideal of R then IG is the ideal of RG generated by the subset I and $(R/I)G \cong RG/IG$. These results and notation may be found in Connell's paper (see [2]).

For any ring R , the Jacobson radical of R will be denoted by $J(R)$ and the characteristic of R by $\text{char } R$. By an Artinian ring we mean a ring that is both left and right Artinian. If R is a ring such that $R/J(R)$ is Artinian then we say that R is *semilocal*. By $p > 0$ we mean that p is a prime number.

Our main result in this paper is as follows:

Theorem 1. *Let R be a ring and G an abelian group. Then RG is semilocal if and only if*

- (i) R is semilocal and G is finite, or
- (ii) R is semilocal and $G \cong G_p \times H$, where G_p is an infinite p -group, H is finite, the order of H is relatively prime to p and $R/J(R)$ is of characteristic $p > 0$.

We remark that if R is a commutative ring and G is an abelian group, Gulliksen, Ribenboim and Viswanathan [4] and Renault [8] have shown that conditions (i) and (ii) in Theorem 1 are necessary and sufficient for RG to be semilocal. Theorem 1 is thus an extension of their result. In the case when $R = \mathbb{F}$ is a field and G is an arbitrary group, the question on whether $\mathbb{F}G$ is semilocal implies that G is locally finite or a finite extension of a p -group (where $p = \text{char } \mathbb{F} > 0$) has been of some interest. Theorem 1 shows that the answer to this question is in the affirmative if G is abelian. S.M. Woods has in fact shown in [9], Theorem 3.2 that G must be torsion if RG is semilocal. J.M. Goursaud [3] and D.S. Passman [7] independently proved that if $\mathbb{F}G$ is semilocal and G is finite, then G is a finite extension of a p -group. J. Lawrence [6] proved that if \mathbb{F} is a field transcendental over the algebraic closure of its prime subfield and $\mathbb{F}G$ is semilocal, then G is a finite extension of a p -group where $p = \text{char } \mathbb{F}$.

2. PRELIMINARIES

For the sake of completeness we first deal with some preliminaries of the proof of Theorem 1.

Theorem 2.1 ([2]). *Let R be a ring and G a group. Then RG is Artinian if and only if R is Artinian and G is finite.*

Theorem 2.2 ([2], [1]). *Let R be a ring and G a group. Then RG is regular if and only if*

- (a) R is regular;
- (b) G is locally finite;
- (c) the order of every finite subgroup of G is a unit in R .

Proposition 2.3. *Let R be a ring and G a group. If R is semilocal and G is finite then RG is semilocal.*

Proof. Since $R/J(R)$ is Artinian and G is finite, so $(R/J(R))G$ is Artinian (by Theorem 2.1). Since G is locally finite, it follows from [2], Proposition 9 that $J(R)G \subseteq J(RG)$. Now consider the mapping $\pi: RG/J(R)G \rightarrow RG/J(RG)$ defined as follows:

$$\pi(x + J(R)G) = x + J(RG), \quad x \in RG.$$

The mapping π is well-defined since $J(R)G \subseteq J(RG)$. It is easy to verify that π is a ring epimorphism. Then since $RG/J(R)G \cong (R/J(R))G$ is Artinian, so is $RG/J(RG)$. Hence RG is semilocal. \square

The proof of the following proposition is straightforward and will be left to the reader.

Proposition 2.4. *Any homomorphic image of a semilocal ring is semilocal.*

Proposition 2.5. *Let R be a ring and G a group. If RG is semilocal then R is semilocal and G is a torsion group.*

Proof. Since $R \cong RG/\Delta$ and RG is semilocal, it follows from Proposition 2.4 that R is semilocal. The assertion that G is a torsion group follows from the proof of Theorem 3.2 in [9]. \square

Lemma 2.6. *Let R_1, \dots, R_n be rings. Then $R = \prod_{i=1}^n R_i$ is semilocal if and only if each R_i is semilocal.*

Proof. We first note that

$$(2.1) \quad R/J(R) = \prod_{i=1}^n R_i / J\left(\prod_{i=1}^n R_i\right) = \prod_{i=1}^n R_i / \prod_{i=1}^n J(R_i) \cong \prod_{i=1}^n R_i / J(R_i).$$

Now if R is semilocal then $R/J(R)$ is Artinian and so is $\prod_{i=1}^n R_i / J(R_i)$ (by (2.1)). Therefore $R_i / J(R_i)$ is Artinian and hence R_i is semilocal ($i = 1, \dots, n$).

Conversely, if each R_i is semilocal, then each $R_i / J(R_i)$ is Artinian. Hence $\prod_{i=1}^n R_i / J(R_i)$ is Artinian and so is $R/J(R)$ (by (2.1)). Therefore R is semilocal and this completes the proof. \square

Lemma 2.7. *Let R be a ring. If $\bar{R} = R/J(R)$ is semilocal, so is R .*

Proof. Since \bar{R} is semiprimitive and semilocal, so $\bar{R} \cong \bar{R}/J(\bar{R})$ is Artinian; hence R is semilocal. \square

For any ring R and positive integer n , we let $M_n(R)$ denote the ring of $n \times n$ matrices over R .

Lemma 2.8. *A ring R is semilocal if and only if $M_n(R)$ is semilocal.*

Proof. It is well-known that

$$M_n(R/J(R)) \cong M_n(R)/M_n(J(R)) = M_n(R)/J(M_n(R)).$$

The result then follows immediately from the fact that a ring R is Artinian if and only if $M_n(R)$ is Artinian (see [5], p. 71). \square

Lemma 2.9. *Let R be a ring and G a group. Then*

$$M_n(R)G \cong M_n(RG).$$

Proof. Let $\theta: M_n(R)G \rightarrow M_n(RG)$ be the mapping defined as follows: For any $A_1g_1 + \dots + A_sg_s \in M_n(R)G$, let

$$\theta(A_1g_1 + \dots + A_sg_s) = (b_{ij}),$$

where $b_{ij} = a_{ij}^{(1)}g_1 + \dots + a_{ij}^{(s)}g_s$ and $a_{ij}^{(m)}$ is the entry in the i -th row and j -th column of A_m , $m = 1, \dots, s$. It may be verified routinely that θ is a ring isomorphism. Hence $M_n(R)G \cong M_n(RG)$. \square

Remark. It is known that if R is a completely reducible ring, then R is isomorphic to a finite direct product of full matrix rings over division rings, that is,

$$R \cong M_{n_1}(D_1) \times \dots \times M_{n_k}(D_k)$$

where D_i is a division ring ($i = 1, \dots, k$). We shall refer to the D_i 's ($i = 1, \dots, k$) as division rings associated with R .

Proposition 2.10. *Let R be a ring and G a group. If RG is semilocal, so is DG for each division ring D associated with $R/J(R)$.*

Proof. Assume that RG is semilocal. By Proposition 2.4 we have that $(R/J(R))G \cong RG/J(R)G$ and $R \cong RG/\Delta$ are semilocal. Therefore, $R/J(R)$ is completely reducible and hence, isomorphic to a finite direct product of full matrix rings over division rings, that is,

$$R/J(R) \cong M_{n_1}(D_1) \times \dots \times M_{n_k}(D_k) = \prod_{i=1}^k M_{n_i}(D_i)$$

for some division rings D_1, \dots, D_k . It follows that

$$\begin{aligned} (R/J(R))G &\cong \left(\prod_{i=1}^k M_{n_i}(D_i) \right) G \cong \prod_{i=1}^k M_{n_i}(D_i)G \\ &\cong \prod_{i=1}^k M_{n_i}(D_iG) \quad (\text{by Lemma 2.9}). \end{aligned}$$

Since $(R/J(R))G$ is semilocal, so is $\prod_{i=1}^k M_{n_i}(D_iG)$ (by Proposition 2.4). It follows from Lemma 2.6 that each $M_{n_i}(D_iG)$ is semilocal and hence by Lemma 2.8, D_iG is semilocal ($i = 1, \dots, k$). \square

Proposition 2.11. *If D is a division ring of characteristic $p > 0$ and G is an abelian group which is a finite extension of a p -group, then DG is semilocal.*

Proof. By assumption we have that G/G_p is finite for some p -subgroup G_p of G . If $G_p = \{1\}$, then G is finite and it follows easily that DG is semilocal. Now assume that $G_p \neq \{1\}$ and let $g \in G_p$, $g \neq 1$. Then $g^{p^n} = 1$ for a positive integer n and therefore, $(1 - g)^{p^n} = 0$. It follows that $1 - g$ is a nilpotent element. Since $1 - g$ lies in the centre of DG , so the ideal generated by $1 - g$ is nilpotent (hence nil). Then, since all nil ideals of DG are contained in $J(DG)$, so $1 - g \in J(DG)$. It thus follows that $\omega G_p \subseteq J(DG)$. Now consider the mapping $\pi: DG/\omega G_p \rightarrow DG/J(DG)$ defined as follows:

$$\pi(x + \omega G_p) = x + J(DG), \quad x \in DG.$$

Since $\omega G_p \subseteq J(DG)$, π is well-defined. It is easily verified that π is a ring epimorphism. Note that $D(G/G_p)$ is Artinian since D is Artinian and G/G_p is finite. Then since $DG/\omega G_p \cong D(G/G_p)$ is Artinian, so is $DG/J(DG)$. Hence DG is semilocal. \square

3. PROOF OF THEOREM 1

We are now ready for the proof of the main theorem.

Proof of Theorem 1. (\Rightarrow): Suppose that RG is semilocal. Since $R \cong RG/\Delta$, it follows from Proposition 2.4 that R is semilocal. By Proposition 2.10 we have that DG is semilocal for each division ring D associated with $R/J(R)$. Let D be one of those division rings and let $p = \text{char } D$. We consider the following cases:

Case 1: $p = 0$. In this case, the order of every finite subgroup of G is a unit in D . By Proposition 2.5 we know that G is torsion. Then, since G is abelian, it follows that G is locally finite. Since D is regular, it follows from Theorem 2.2 that DG is regular. Hence $J(DG) = \{0\}$ and therefore $DG \cong DG/J(DG)$ is Artinian. We thus have that G is finite, that is (i) occurs.

Case 2: $p > 0$. Since G is an abelian torsion group we may write $G \cong G_p \times H$ where G_p is the Sylow p -subgroup of G and the order of every element of H is prime to p . Clearly, the order of every finite subgroup of H is a unit in D . Furthermore,

H is locally finite (since G is locally finite) and D is regular. It follows that DH is regular (by Theorem 2.2); hence $J(DH) = \{0\}$. Since

$$DH \cong D(G/G_p) \cong DG/\omega G_p$$

is semilocal (by Proposition 2.4), so $DH \cong DH/J(DH)$ is Artinian and hence H is finite. If G_p is finite, then (i) occurs.

Now suppose that G_p is infinite. We show that each of the division rings associated with the completely reducible ring $R/J(R)$ has the characteristic p . Suppose that there exists a division ring D' associated with $R/J(R)$ such that $\text{char } D' = q$ and $q \neq p$. If $q = 0$, then by the same argument as in Case 1 we have that G is finite. But this is impossible since $G_p \subseteq G$ and G_p is infinite. If $q > 0$, then since G is an abelian torsion group, we may write $G \cong G_q \times H'$, where G_q is the Sylow q -subgroup of G and the order of every element of H' is prime to q . By the same argument as in the preceding paragraph we can show that H' is finite. But since $G_p \subseteq H'$, this implies that G_p is finite; a contradiction. Hence if G_p is infinite, then each of the division rings associated with $R/J(R)$ is of the characteristic $p > 0$. It follows then that $\text{char } R/J(R) = p$ and hence, (ii) occurs.

(\Leftarrow): Suppose that (i) occurs. It follows readily from Proposition 2.3 that RG is semilocal.

Now suppose that (ii) occurs. Then $R/J(R)$ is a completely reducible ring and therefore it is isomorphic to a finite direct product of full matrix rings over division rings, that is,

$$R/J(R) \cong M_{n_1}(D_1) \times \dots \times M_{n_k}(D_k) = \prod_{i=1}^k M_{n_i}(D_i)$$

for some division rings D_1, \dots, D_k . Therefore,

$$(3.1) \quad \begin{aligned} RG/J(R)G &\cong (R/J(R))G \cong \prod_{i=1}^k M_{n_i}(D_i)G \\ &\cong \prod_{i=1}^k M_{n_i}(D_iG) \quad (\text{by Lemma 2.9}). \end{aligned}$$

Since $\text{char } R/J(R) = p > 0$, so each D_i is a division ring of the characteristic $p > 0$. Then since G is a finite extension of a p -group, it follows from Proposition 2.11 that each D_iG is semilocal. From Lemma 2.8 we have that $M_{n_i}(D_iG)$ is semilocal for each i . Therefore $\prod_{i=1}^k M_{n_i}(D_iG)$ is semilocal (by Lemma 2.6) and it follows from (3.1)

that $RG/J(R)G$ is semilocal. Now since G is locally finite, so $J(R)G \subseteq J(RG)$ (by [2], Proposition 9). Consider the mapping $\pi: RG/J(R)G \rightarrow RG/J(RG)$ defined as follows:

$$\pi(\alpha + J(R)G) = \alpha + J(RG), \quad \alpha \in RG.$$

By routine verification, π is a well-defined ring epimorphism. Since $RG/J(R)G$ is semilocal, so is $RG/J(RG)$ (by Proposition 2.4). It then follows from Lemma 2.7 that RG is semilocal.

This completes the proof of the theorem. □

References

- [1] *M. Auslander*: On regular group rings. Proc. Amer. Math. Soc. 8 (1957), 658–664.
- [2] *I. G. Connell*: On the group ring. Canad. J. Math. 15 (1963), 650–685.
- [3] *J. M. Goursaud*: Sur les anneaux de groupes semi-parfaits. Canad. J. Math. 25 (1973), 922–928.
- [4] *K. Gulliksen, P. Ribenboim and T. M. Viswanathan*: An elementary note on group rings. J. Reine Angew. Math. 242 (1970), 148–162.
- [5] *J. Lambek*: Lectures on Rings and Modules. Blaisdell, Waltham, Mass., 1966.
- [6] *J. Lawrence*: Semilocal group rings and tensor products. Michigan Math. J. 22 (1975), 309–313.
- [7] *D. S. Passman*: Advances in group rings. Israel J. Math. 19 (1974), 67–107.
- [8] *G. Renault*: Sur les anneaux de groupes. C. R. Acad. Sci. Paris Ser. A 273 (1971), 84–87.
- [9] *S. M. Woods*: Some results on semi-perfect group rings. Canad. J. Math. 28 (1974), 121–129.

Author's address: Institute of Mathematical Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia, e-mail: acym@mnt.math.um.edu.my.