

Anna de Simone; Daniele Mundici; Mirko Navara
A Cantor-Bernstein theorem for σ -complete MV-algebras

Czechoslovak Mathematical Journal, Vol. 53 (2003), No. 2, 437–447

Persistent URL: <http://dml.cz/dmlcz/127812>

Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A CANTOR-BERNSTEIN THEOREM
FOR σ -COMPLETE MV-ALGEBRAS

A. DE SIMONE, Napoli, D. MUNDICI, Florence, and M. NAVARA, Praha

(Received June 14, 2000)

Abstract. The Cantor-Bernstein theorem was extended to σ -complete boolean algebras by Sikorski and Tarski. Chang's MV-algebras are a nontrivial generalization of boolean algebras: they stand to the infinite-valued calculus of Łukasiewicz as boolean algebras stand to the classical two-valued calculus. In this paper we further generalize the Cantor-Bernstein theorem to σ -complete MV-algebras, and compare it to a related result proved by Jakubík for certain complete MV-algebras.

Keywords: Cantor-Bernstein theorem, MV-algebra, boolean element of an MV-algebra, partition of unity, direct product decomposition, σ -complete MV-algebra

MSC 2000: 06D35, 06D30, 06C15, 03G20

1. INTRODUCTION

The Cantor-Bernstein theorem states that, if a set X can be embedded into a set Y , and vice versa, then there is a one-one map of X onto Y . The theorem was proved by Dedekind in 1887, conjectured by Cantor in 1895, and again proved by Bernstein in 1898, [6, p. 85].

For any boolean algebra A , let $[0, a]$ denote the boolean algebra of all $x \in A$ such that $0 \leq x \leq a$, equipped with the restriction of the join and meet of A , where the complement of $y \in [0, a]$ is the meet of a with the complement $\neg y$ of y in A . (Note that Sikorski [8, p. 29] writes $A|a$ instead of $[0, a]$.)

Research supported by COST ACTION 15 on Many-valued logics for computer science applications, Project on Logic of the Italian MURST, Project VS 96049 of the Czech Ministry of Education and Grant 201/97/0437 of the Grant Agency of the Czech Republic.

Sikorski [9] and Tarski [10] proved the following generalization of the (Dedekind)-Cantor-Bernstein theorem: For any two σ -complete boolean algebras A and B and elements $a \in A$ and $b \in B$, if B is isomorphic to $[0, a]$ and A is isomorphic to $[0, b]$, then A and B are isomorphic. To obtain the Cantor-Bernstein theorem it suffices to assume that A and B are the powersets of X and Y , respectively, with the natural set-theoretic boolean operations.

Our aim in this paper is to further generalize the Cantor-Bernstein theorem to MV-algebras—the latter being an interesting “non-commutative” extension of boolean algebras (see [7] for a precise formulation of this). Since, as proved in [3] and [5], the σ -completeness assumption is indispensable already in the boolean algebraic setup, our Cantor-Bernstein theorem shall be proved for σ -complete MV-algebras.

2. MV-ALGEBRAS

An *MV-algebra* $A = (A, 0, \oplus, \neg)$ is an algebra where the operation $\oplus: A \times A \rightarrow A$ is associative and commutative with 0 as the neutral element, the operation $\neg: A \rightarrow A$ satisfies the identities $\neg\neg x = x$ and $x \oplus \neg 0 = \neg 0$, and, in addition,

$$(1) \quad y \oplus \neg(y \oplus \neg x) = x \oplus \neg(x \oplus \neg y).$$

Example 2.1. The real unit interval $[0, 1]$ equipped with the operations $x \oplus y = \min(1, x + y)$ and $\neg x = 1 - x$ is an MV-algebra.

Following common usage, for any elements x, y of an MV-algebra we will use the abbreviations $1 = \neg 0$, $x \odot y = \neg(\neg x \oplus \neg y)$ and $x \ominus y = x \odot \neg y$. We will denote by (A, \vee, \wedge) the underlying distributive lattice of A , where $x \vee y = x \oplus \neg(x \oplus \neg y)$ and $x \wedge y = x \odot \neg(x \odot \neg y)$. With reference to the underlying order of A , for any element $a \in A$ we let the interval $[0, a]$ be defined by

$$[0, a] = \{x \in A \mid 0 \leq x \leq a\}.$$

An MV-algebra A is σ -complete (*complete*) iff every sequence (every family, respectively) of elements of A has supremum in A with respect to the underlying order of A .

As shown by Chang, boolean algebras coincide with MV-algebras satisfying the equation $x \oplus x = x$. In this case the operation \oplus coincides with \vee , and the operation \odot coincides with \wedge .

An element a in an MV-algebra A is called *boolean* iff $a \oplus a = a$. We let $\mathbf{B}(A)$ denote the set of boolean elements of A . It is not hard to see that the operations of A make $\mathbf{B}(A)$ a boolean algebra. As shown in Corollary 3.3 below, if A is a σ -complete

MV-algebra, then $\mathbf{B}(A)$ is a σ -complete boolean algebra, and the σ -infinitary operations of $\mathbf{B}(A)$ agree with the restrictions of the corresponding operations of A .

A *homomorphism* between two MV-algebras is a map that sends zero to zero, and preserves the operations \oplus and \neg . A one-one surjective homomorphism is called an *isomorphism*.

For further information on MV-algebras we refer to [1], [2] and [7].

Definition 2.2. Let A be an MV-algebra and z a fixed, but otherwise arbitrary, element of A . Let the map $h_z: A \rightarrow [0, z]$ be defined by

$$(2) \quad h_z(x) = x \wedge z.$$

Further, we define the operation $\neg_z: [0, z] \rightarrow [0, z]$ by

$$(3) \quad \neg_z x = z \odot \neg x = z \ominus x,$$

and the operation $\oplus_z: [0, z] \times [0, z] \rightarrow [0, z]$ by

$$(4) \quad x \oplus_z y = (x \oplus y) \wedge z.$$

A moment's reflection shows that the ranges of both operations \neg_z and \oplus_z coincide with $[0, z]$.

Proposition 2.3. Let A be an MV-algebra and $b \in A$. We then have

- (i) for each element $b \in A$, the structure $([0, b], \oplus_b, \neg_b, 0)$ is an MV-algebra.
- If, in addition, b is a boolean element of A then
- (ii) $\neg_b x = b \wedge \neg x$ for all $x \in [0, b]$;
 - (iii) the interval $[0, b]$ (as well as the interval $[0, \neg b]$) is an ideal of A ;
 - (iv) The map h_b defined in (2) is a homomorphism of A onto $[0, b]$ whose kernel coincides with $[0, \neg b]$;
 - (v) The MV-algebra $[0, b]$ is isomorphic to the quotient MV-algebra $A/[0, \neg b]$;
 - (vi) $[0, b]$ is a subalgebra of A iff $b = 1$ iff $[0, b] = A$.

Proof. (i) For every $x \in [0, b]$ we have

$$\neg_b \neg_b x = b \odot \neg(b \odot \neg x) = b \wedge x = x$$

and

$$x \oplus_b \neg_b 0 = x \oplus_b b = (x \oplus b) \wedge b = b = \neg_b 0.$$

Associativity of \oplus_b follows from the identities

$$\begin{aligned}(x \oplus_b y) \oplus_b z &= (((x \oplus y) \wedge b) \oplus z) \wedge b \\ &= ((x \oplus y \oplus z) \wedge (b \oplus z)) \wedge b \\ &= (x \oplus y \oplus z) \wedge b = \dots = x \oplus_b (y \oplus_b z).\end{aligned}$$

With reference to (1) we shall now prove the identity

$$(5) \quad y \oplus_b \neg_b(y \oplus_b \neg_b x) = x \oplus_b \neg_b(x \oplus_b \neg_b y) \quad \text{for all } x, y \in [0, b].$$

First, using distributivity of \odot over \vee , we transform a part of the expression on the left-hand side of (5) as follows:

$$\begin{aligned}\neg_b(y \oplus_b \neg_b x) &= b \odot \neg((y \oplus (b \odot \neg x)) \wedge b) \\ &= b \odot ((\neg y \odot \neg(b \odot \neg x)) \vee \neg b) = b \odot \neg y \odot \neg(b \odot \neg x) \\ &= \neg y \odot (b \wedge x) = \neg y \odot x = \neg(y \oplus \neg x).\end{aligned}$$

We can now simplify the left-hand term in (5) as follows:

$$y \oplus_b \neg_b(y \oplus_b \neg_b x) = (y \oplus \neg(y \oplus \neg x)) \wedge b = (y \vee x) \wedge b = y \vee x,$$

which settles (5). The remaining verifications needed to show that $[0, b]$ is an MV-algebra are all trivial.

Following now the proof of [2, Proposition 6.4.1], let us assume that $b \in \mathbf{B}(A)$. Then condition (ii) is an immediate consequence of the definition of \neg_b and of the fact that \odot coincides with \wedge whenever one of its arguments is boolean, [2, Theorem 1.5.3]. Similarly, (iii) follows from the definition of a boolean element, [2, Corollary 1.5.6], and we also see that \oplus_b coincides with the restriction of \oplus to $[0, b]$. To prove (iv), for all $x, y \in A$ we can write $(x \wedge b) \oplus (y \wedge b) = ((x \wedge b) \oplus y) \wedge ((x \wedge b) \oplus b)$. From $(x \wedge b) \oplus b = (x \wedge b) \vee b = b$ we get $(x \wedge b) \oplus (y \wedge b) = (x \oplus y) \wedge (b \oplus y) \wedge b = (x \oplus y) \wedge b$. We conclude that $h_b(x \oplus y) = h_b(x) \oplus h_b(y) = h_b(x) \oplus_b h_b(y)$. The rest is trivial. The proof of (v) and (vi) is the same as in [2, Proposition 6.4.3]. \square

Remarks. As shown by (ii) above, whenever b is a boolean element of A , there is no discrepancy between our present definition of \neg_b and the definition in [2, (6.4)].

If in a boolean algebra B we denote by \mathcal{I} the principal ideal generated by $\neg b$, then $\mathcal{I} = [0, \neg b]$ and the algebra $[0, b]$ is isomorphic to B/\mathcal{I} via the map $x \in [0, b] \mapsto x/\mathcal{I} \in B/\mathcal{I}$. Condition (v) is a generalization of this fact to MV-algebras.

If a is not a boolean element of A , then $[0, a]$ need not be a homomorphic image of A . For an example, let $A = \{0, 1/2, 1\}$ be a subalgebra of the MV-algebra $[0, 1]$

of Example 2.1. Then $[0, 1/2] = \{0, 1/2\}$ is not a homomorphic image of A , because A has no other ideals than $\{0\}$. One more example is given in 5.2 below.

On the other hand, the existence of a homomorphism of A onto $[0, a]$ need not imply that a is a boolean element of A . As a matter of fact, in the MV-algebra $[0, 1]$ of Example 2.1, multiplication by $1/2$ is a homomorphism of $[0, 1]$ onto the interval MV-algebra $[0, 1/2]$, but the element $1/2$ is not boolean in $[0, 1]$.

The proof of the following result is immediate.

Lemma 2.4. *Let A and B be MV-algebras and let $\alpha: A \rightarrow B$ be an isomorphism of A onto B . For any $a \in A$, the restriction of the map α to the interval $[0, a]$ of A is an isomorphism of the MV-algebra $[0, a]$ onto the interval $[0, \alpha(a)]$ of B , once these two intervals are equipped with the MV-algebraic operations of Definition 2.2 and Proposition 2.3 (i).*

Corollary 2.5. *For each $a \in \mathbf{B}(A)$, the mapping $x \mapsto (x \wedge a, x \wedge \neg a)$ is an isomorphism of A onto the product MV-algebra $[0, a] \times [0, \neg a]$.*

Proof. The same as for [2, Lemma 6.4.5]. □

3. PARTITIONS OF UNITY AND DECOMPOSITIONS

In Lemma 3.4 below we will give an infinitary generalization of Corollary 2.5. To this purpose, we prepare

Notation. We set $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and $\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$.

Lemma 3.1. *Let A be a σ -complete MV-algebra. Let $x_1, x_2, \dots \in A$. Then the following countably infinitary de Morgan identities hold:*

$$(6) \quad \bigwedge_{n \in \mathbb{N}} x_n = \neg \bigvee_{n \in \mathbb{N}} \neg x_n$$

and

$$(7) \quad \bigvee_{n \in \mathbb{N}} x_n = \neg \bigwedge_{n \in \mathbb{N}} \neg x_n.$$

Proof. Let $a = \bigwedge_{n \in \mathbb{N}} x_n$ and $b = \bigvee_{n \in \mathbb{N}} \neg x_n$. For all $n \in \mathbb{N}$ we have $a \leq x_n$, whence $\neg a \geq \neg x_n$. Therefore $\neg a \geq \bigvee_{n \in \mathbb{N}} \neg x_n = b$. Similarly, for all $n \in \mathbb{N}$ we have $b \geq \neg x_n$, $\neg b \leq x_n$, whence $\neg b \leq \bigwedge_{n \in \mathbb{N}} x_n = a$ and $b \geq \neg a$, which settles (6). The proof of (7) is similar. □

As an immediate consequence we get the following infinitary distributive laws:

Lemma 3.2. *Let A be a σ -complete MV-algebra. Let $x_1, x_2, \dots \in A$. Then for each $x \in A$ we have*

$$(8) \quad x \wedge \bigvee_{n \in \mathbb{N}} x_n = \bigvee_{n \in \mathbb{N}} (x \wedge x_n)$$

and

$$(9) \quad x \vee \bigwedge_{n \in \mathbb{N}} x_n = \bigwedge_{n \in \mathbb{N}} (x \vee x_n).$$

Proof. This is an easy adaptation of the proof of [2, Lemma 6.6.4]. □

Corollary 3.3. *Let A be a σ -complete MV-algebra. Then*

(i) $\mathbf{B}(A)$ is a σ -complete boolean algebra. As a matter of fact, for any sequence $b_1, b_2, \dots \in \mathbf{B}(A)$ we have

$$(10) \quad \bigvee_{n \in \mathbb{N}} b_n \in \mathbf{B}(A)$$

and

$$(11) \quad \bigwedge_{n \in \mathbb{N}} b_n \in \mathbf{B}(A).$$

(ii) For every $b \in \mathbf{B}(A)$, letting $h_b: A \rightarrow [0, b]$ be as in (2), it follows that $[0, b]$ is a σ -complete MV-algebra, and h_b preserves all existing infima and suprema. Therefore, for all $x_1, x_2, \dots \in A$ we can write

$$(12) \quad h_b \left(\bigvee_{n \in \mathbb{N}} x_n \right) = \bigvee_{n \in \mathbb{N}} h_b(x_n)$$

and

$$(13) \quad h_b \left(\bigwedge_{n \in \mathbb{N}} x_n \right) = \bigwedge_{n \in \mathbb{N}} h_b(x_n).$$

Proof. An easy adaptation of the proof of [2, Corollary 6.6.5]. □

From Lemma 3.2 we obtain the desired generalization of Corollary 2.5, given by the following variant of [2, Lemma 6.6.6].

Lemma 3.4. *Let A be a σ -complete MV-algebra. Suppose that a sequence $b_1, b_2, \dots \in \mathbf{B}(A)$ satisfies the following conditions:*

$$\text{(Partition of unity)} \quad \bigvee_{n \in \mathbb{N}} b_n = 1 \quad \text{and} \quad b_j \wedge b_k = 0 \quad \text{for all } j \neq k.$$

Then the map $x \mapsto (x \wedge b_1, x \wedge b_2, \dots) = (x \wedge b_n)_{n \in \mathbb{N}}$ is an isomorphism of A onto the product MV-algebra $\prod_{n \in \mathbb{N}} [0, b_n]$.

4. MV-ALGEBRAIC CANTOR-BERNSTEIN THEOREM

In this section we prove the following MV-algebraic generalization of the Cantor-Bernstein theorem.

Theorem 4.1. *Let A and B be σ -complete MV-algebras. Let $a \in \mathbf{B}(A)$, $b \in \mathbf{B}(B)$, and assume α to be an isomorphism of A onto the interval algebra $[0, b]$ of B , and β an isomorphism of B onto the interval algebra $[0, a]$ of A . Then A and B are isomorphic.*

Proof. Skipping all trivialities, we may safely assume $0 < a < 1$ and $0 < b < 1$. Also, A and B can be safely assumed disjoint. We can now define sequences $a_0, a_1, a_2, \dots \in A$ and $b_0, b_1, b_2, \dots \in B$ by the following inductive stipulation:

$$\begin{aligned} a_0 &= 1, & b_0 &= 1, \\ a_{n+1} &= \beta(b_n), & b_{n+1} &= \alpha(a_n). \end{aligned}$$

For each $n = 0, 1, 2, \dots$, both elements a_n and b_n are boolean. From the assumed injectivity of α and β we obtain

$$(14) \quad a_0 > a_1 > a_2 > \dots \quad \text{and} \quad b_0 > b_1 > b_2 > \dots$$

Let $a_\infty \in A$ and $b_\infty \in B$ be given by $a_\infty = \bigwedge_{n \in \mathbb{N}_0} a_n$ and $b_\infty = \bigwedge_{n \in \mathbb{N}_0} b_n$. The existence of a_∞ and b_∞ is ensured by the assumed σ -completeness of A and B . By Corollary 3.3(i) both a_∞ and b_∞ are boolean elements. For all $n \in \mathbb{N}_0$ we have the identities $a_{n+2} = (\beta \circ \alpha)(a_n)$ and $b_{n+2} = (\alpha \circ \beta)(b_n)$. Since the mapping $\beta \circ \alpha$ is an isomorphism of A onto $[0, a_2]$, it preserves countable infima and suprema. Since

for each $n = 0, 1, 2, \dots$ the underlying orders of the interval MV-algebras $[0, a_n]$ and $[0, a_{n+1}]$ agree, we have

$$(\beta \circ \alpha)(a_\infty) = (\beta \circ \alpha)\left(\bigwedge_{n \in \mathbb{N}} a_n\right) = \bigwedge_{n \in \mathbb{N}} (\beta \circ \alpha)(a_n) = \bigwedge_{n \in \mathbb{N}} a_{n+2} = a_\infty.$$

Similarly, $b_\infty = (\alpha \circ \beta)(b_\infty)$. One similarly obtains

$$(15) \quad \alpha(a_\infty) = b_\infty \quad \text{and} \quad \beta(b_\infty) = a_\infty.$$

In particular, $a_\infty = 0$ iff $b_\infty = 0$. For each $n = 0, 1, 2, \dots$ let us define $d_n = a_n \ominus a_{n+1} = a_n \wedge \neg a_{n+1}$ and $e_n = b_n \ominus b_{n+1} = b_n \wedge \neg b_{n+1}$. Then for each $n = 0, 1, 2, \dots$ we have

$$(16) \quad \alpha(d_{2n}) = e_{2n+1} \quad \text{and} \quad \beta(e_{2n}) = d_{2n+1}.$$

A straightforward computation shows that, for any two distinct $m, n \in \mathbb{N}_0$, $d_m \wedge d_n = 0 = e_m \wedge e_n$.

Lemma 3.1 together with (14) yields

$$\bigvee_{n \in \mathbb{N}} d_{n-1} = \bigvee_{n \in \mathbb{N}} \bigvee_{k=1}^n d_{k-1} = \bigvee_{n \in \mathbb{N}} (1 \ominus a_n) = \bigvee_{n \in \mathbb{N}} \neg a_n = \neg \bigwedge_{n \in \mathbb{N}} a_n = \neg a_\infty.$$

It follows that the sequence $(a_\infty, d_0, d_1, d_2, \dots)$ is a partition of unity in $\mathbf{B}(A)$. Analogously, the sequence $(b_\infty, e_0, e_1, e_2, \dots)$ is a partition of unity in $\mathbf{B}(B)$. By Lemma 3.4, the map

$$x \mapsto (x \wedge a_\infty, x \wedge d_0, x \wedge d_1, x \wedge d_2, \dots)$$

is an isomorphism of A onto the product MV-algebra $[0, a_\infty] \times [0, d_0] \times [0, d_1] \times [0, d_2] \times \dots$. Similarly, the map

$$y \mapsto (y \wedge b_\infty, y \wedge e_0, y \wedge e_1, y \wedge e_2, \dots)$$

is an isomorphism of B onto $[0, b_\infty] \times [0, e_0] \times [0, e_1] \times [0, e_2] \times \dots$. By Lemma 2.4 and (15), the restriction of α to $[0, a_\infty]$ is an isomorphism of $[0, a_\infty]$ onto $[0, b_\infty]$, in symbols (and with a slight abuse of notation),

$$\alpha: [0, a_\infty] \cong [0, b_\infty].$$

Another application of Lemma 2.4 together with (16) yields, for each $n = 0, 1, 2, \dots$, an isomorphism

$$\alpha: [0, d_{2n}] \cong [0, e_{2n+1}].$$

Similarly, from the isomorphism $\beta: [0, e_{2n}] \cong [0, d_{2n+1}]$ one obtains an isomorphism

$$\beta^{-1}: [0, d_{2n+1}] \cong [0, e_{2n}]$$

for each $n = 0, 1, 2, \dots$. It is now easy to obtain an isomorphism of $[0, a_\infty] \times [0, d_0] \times [0, d_1] \times [0, d_2] \times \dots$ onto $[0, b_\infty] \times [0, e_0] \times [0, e_1] \times [0, e_2] \times \dots$, whence one has the desired isomorphism of A onto B . \square

If A happens to be a boolean algebra, the above theorem reduces to the boolean-algebraic Cantor-Bernstein theorem stated in the introduction, and proved by Sikorski and Tarski.

5. A RELATED RESULT BY JAKUBÍK

In his paper [4], Jakubík proved a different form of Cantor-Bernstein theorem for MV-algebras. In this section we shall compare Jakubík's result with our Theorem 4.1.

A *lattice isomorphism* between two MV-algebras A and B is a one-one map of A onto B that preserves the underlying lattice structures of A and B . We say that A and B are *lattice isomorphic* iff there is a lattice isomorphism between A and B .

Let $\mathcal{D} \subseteq [0, 1]$ be the MV-algebra consisting of all rational numbers in $[0, 1]$ whose denominator is $1, 2, 4, 8, 16, \dots$. Let \mathcal{Q} be the subalgebra of $[0, 1]$ consisting of all rational numbers in $[0, 1]$. Then \mathcal{D} and \mathcal{Q} are lattice isomorphic (as denumerable, densely ordered chains with two endpoints) but they are not isomorphic MV-algebras. As a matter of fact, the equation $x \oplus x = \neg x$ has a solution in \mathcal{Q} , but does not have any solution in \mathcal{D} . Thus, the existence of a lattice isomorphism between two MV-algebras need not imply that the two MV-algebras are isomorphic. Trivially, if two MV-algebras are isomorphic then their underlying lattices are isomorphic.

For any MV-algebra A let us consider the following property:

$$(*) \quad \text{If } a \in A \text{ and } [0, a] \text{ is a boolean algebra, then } a \in \mathbf{B}(A).$$

Jakubík proved

Theorem 5.1 [4]. *Let A and B be complete MV-algebras satisfying condition (*). Suppose that for some $a \in A$, $b \in B$, A is lattice isomorphic to $[0, b]$ and B is lattice isomorphic to $[0, a]$. Then A and B are isomorphic as MV-algebras.*

The rest of this section is devoted to a comparison between Jakubík's theorem and our Theorem 4.1. To this aim, we present an example that simultaneously shows the necessity of condition (*) in Jakubík's Theorem 5.1 and the necessity of the assumption that a and b are boolean in our Theorem 4.1.

Example. Let $\mathcal{K} = \{0, 1/2, 1\}$ be the uniquely determined three-element subalgebra of the MV-algebra $[0, 1]$ from Example 2.1. Denote by A the product of denumerably many copies of \mathcal{K} ,

$$A = \mathcal{K} \times \mathcal{K} \times \mathcal{K} \times \dots$$

With pointwise defined operations, A is a complete MV-algebra. Let elements $a, b \in A$ be defined by

$$\begin{aligned} a &= (1/2, 1, 1, 1, \dots), \\ b &= (0, 1, 1, 1, \dots). \end{aligned}$$

Then $B = [0, a]$ equipped with the operations from Definition 2.2 is a complete MV-algebra which is (isomorphic to) an interval of A . On the other hand, A is isomorphic to $[0, b]$ via the isomorphism $\alpha: A \rightarrow [0, b]$ defined by

$$\alpha((x_1, x_2, x_3, \dots)) = (0, x_1, x_2, x_3, \dots).$$

A fortiori, B is lattice isomorphic to an interval of A , and A is lattice isomorphic to the interval $[0, b]$ of B . Nevertheless, A and B are not isomorphic MV-algebras. Indeed, the element $c = (1/2, 0, 0, \dots)$ is an atom of B (minimal nonzero element) and it also belongs to the boolean algebra $\mathbf{B}(B)$, while no atom of A is boolean.

Trivially, the interval $[0, c] = \{0, c\}$ is a boolean algebra, but the atom c is not boolean in A , and condition (*) is not satisfied. On the other hand, all the other assumptions of Theorem 5.1 are satisfied. This shows the necessity of assumption (*) in Jakubík's Theorem 5.1. The present example also shows that our Theorem 4.1 would no longer hold without assuming the elements a and b therein to be boolean.

Note that Theorem 4.1 also holds for MV-algebras not satisfying condition (*). We can, for example, apply it to the MV-algebras A and B of the above example. On the other hand, the assumption that a and b are boolean is not needed in Theorem 5.1.

We finally remark that Theorem 5.1 is stated for complete MV-algebras, while our result here is valid for a larger class of σ -complete MV-algebras.

Altogether, Theorems 5.1 and 4.1 are incomparable.

References

- [1] *R. Cignoli and D. Mundici*: An invitation to Chang's MV-algebras. In: *Advances in Algebra and Model Theory* (M. Droste, R. Göbel, eds.). Gordon and Breach Publishing Group, Reading, UK, 1997, pp. 171–197.
- [2] *R. Cignoli, I. M. L. D'Ottaviano and D. Mundici*: Algebraic Foundations of Many-valued Reasoning. *Trends in Logic*. Vol. 7. Kluwer Academic Publishers, Dordrecht, 1999.
- [3] *W. Hanf*: On some fundamental problems concerning isomorphism of boolean algebras. *Math. Scand.* 5 (1957), 205–217.
- [4] *J. Jakubík*: Cantor-Bernstein theorem for MV-algebras. *Czechoslovak Math. J.* 49(124) (1999), 517–526.
- [5] *S. Kinoshita*: A solution to a problem of Sikorski. *Fund. Math.* 40 (1953), 39–41.
- [6] *A. Levy*: Basic Set Theory. *Perspectives in Mathematical Logic*. Springer-Verlag, Berlin, 1979.
- [7] *D. Mundici*: Interpretation of AF C^* -algebras in Łukasiewicz sentential calculus. *J. Funct. Anal.* 65 (1986), 15–63.
- [8] *R. Sikorski*: Boolean Algebras. Springer-Verlag. *Ergebnisse Math. Grenzgeb.*, Berlin, 1960.
- [9] *R. Sikorski*: A generalization of a theorem of Banach and Cantor-Bernstein. *Colloq. Math.* 1 (1948), 140–144 and 242.
- [10] *A. Tarski*: Cardinal Algebras. Oxford University Press, New York, 1949.

Authors' addresses: A. De Simone, Dip. di Matematica e Statistica, Università degli Studi di Napoli "Federico II", Complesso Monte S. Angelo, Via Cintia, 80126 Napoli, Italy, e-mail: annades@unina.it; D. Mundici, Department of Mathematics, University of Florence, Viale Morgagni 67/A, 50134 Florence, Italy, e-mail: mundici@math.unifi.it; M. Navara, Center for Machine Perception, Department of Cybernetics, Faculty of Electrical Engineering, Czech Technical University, Technická 2, 166 27 Praha 6, Czech Republic, e-mail: navara@cmp.felk.cvut.cz.