

Raúl Naulin

On the instability of linear nonautonomous delay systems

Czechoslovak Mathematical Journal, Vol. 53 (2003), No. 3, 497–514

Persistent URL: <http://dml.cz/dmlcz/127818>

Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE INSTABILITY OF LINEAR
NONAUTONOMOUS DELAY SYSTEMS

RAÚL NAULIN, Cumaná

(Received May 25, 2000)

Abstract. The unstable properties of the linear nonautonomous delay system $x'(t) = A(t)x(t) + B(t)x(t - r(t))$, with nonconstant delay $r(t)$, are studied. It is assumed that the linear system $y'(t) = (A(t) + B(t))y(t)$ is unstable, the instability being characterized by a nonstable manifold defined from a dichotomy to this linear system. The delay $r(t)$ is assumed to be continuous and bounded. Two kinds of results are given, those concerning conditions that do not include the properties of the delay function $r(t)$ and the results depending on the asymptotic properties of the delay function.

Keywords: Liapounov instability, h -instability, instability of delay equations, nonconstant delays

MSC 2000: 34D20, 34D05

1. INTRODUCTION

In this paper sufficient conditions for the solutions of the equation

$$(1) \quad x'(t) = A(t)x(t) + B(t)x(t - r(t)), \quad x \in \mathbb{C}^n, \quad t \geq t_0,$$

to be unstable are given. The problem of stability and instability, in the case of constant matrices A , B and a constant delay, r has been studied by many authors [5], [10]. For a good acquaintance with the subject, as well as for the application to mathematical ecology, the reader is referred to the monograph [7]. The constant case is frequently studied by means of the location of the roots of the so called characteristic polynomial: $P(\lambda) = \det(\lambda I - A - Be^{-r\lambda})$. An outstanding result

This research was supported by Proyecto CI-5-1003-0934/00 and Proyecto CI-5-1003-0936/00.

ensures that the solutions of the equation

$$x'(t) = Ax(t) + Bx(t - r)$$

are unstable if there exists a root of the characteristic polynomial with a positive real part. This algebraic method cannot be applied to the nonautonomous Eq. (1).

The unstable properties of Eq. (1) have not been sufficiently studied [5], [7], [10], [11]. This contrasts with the evolution of this theory for ordinary differential equations, to which we may mention the classical result of Perron [1] and Coppel's theorem on instability [3] for nonautonomous systems.

In this paper we will give results on instability and asymptotic instability of Eq. (1), relying on the ideas of the paper [12]. We will distinguish the case when B is integrable, where, under suitable conditions, the instability can be obtained for a general bounded delay $r(t)$. The case when B is not integrable is treated in our paper by means of conditions of boundedness and integrability of the function $r(t)B(t)$. In a section of examples we will apply the results obtained to different classes of delay differential equations.

2. BASIC DEFINITIONS AND NOTATION

The symbol \mathbf{V} will denote the linear space \mathbb{R}^n or \mathbb{C}^n ; $|x|$ stands for a fixed norm of the vector x , and the corresponding matrix norm of a matrix A will be denoted by $|A|$. We will assume that the function $r(t): [t_0, \infty) \rightarrow [0, \sigma]$, $\sigma > 0$, is continuous. Throughout, we will denote $J = [t_0, \infty)$, $J_\sigma = [t_0 - \sigma, \infty)$; the functions $A, B: J_\sigma \rightarrow \mathbf{V}$, are assumed to be continuous; Φ will denote the fundamental matrix of

$$(2) \quad y'(t) = (A(t) + B(t))y(t)$$

satisfying $\Phi(t_0) = I$, where the matrix I denotes the identity; for an interval $I_{a,b}$ of real numbers, we will denote by $C(I_{a,b})$ the space of continuous and bounded functions defined on $I_{a,b}$ with values on \mathbf{V} ; if $x \in C(J_\sigma)$ and $t \geq t_0$, then $x_t \in C([- \sigma, 0])$ will denote the function $x_t(s) = x(t + s)$, $s \in [- \sigma, 0]$; in the space $C([- \sigma, 0])$, the following norm will be used:

$$|\varphi|_\sigma = \max_{s \in [- \sigma, 0]} |\varphi(s)|;$$

$x(t; t_0, \varphi)$ will denote the unique solution of the problem

$$(3) \quad \begin{cases} x'(t) = A(t)x(t) + B(t)x(t - r(t)), & t \geq t_0, \\ x_{t_0} = \varphi, & \varphi \in C([- \sigma, 0]); \end{cases}$$

the letters h, k, p, q will denote positive continuous functions, and $h^{-1}(t) = 1/h(t)$; for $f \in C(J_\sigma)$, let $|f|^\infty := \sup\{|f(t)| : t \in J_\sigma\}$, $C_h(J_\sigma) := \{f : J_\sigma \rightarrow \mathbf{V} : h^{-1}f \in C(J_\sigma)\}$; for $f \in C_h(J_\sigma)$, let $|f|_h := |h^{-1}f|^\infty$ and $B_h[0, \varrho] = \{f \in C_h(J_\sigma) : |f|_h \leq \varrho\}$; $C_h(J_\sigma)$, called the space of h -bounded functions, endowed with the norm $|\cdot|_h$ is a Banach space; $L^1(J)$ will denote the space of integrable functions defined on J , with norm $|f|^1 = \int_{t_0}^\infty |f(s)| ds$; $L_h^1(J)$ is the space of h -integrable functions, that is $f \in L_h^1$ iff $|f|_h^1 := \int_{t_0}^\infty h^{-1}(s)|f(s)| ds < \infty$.

The instability of Eq. (2) will be characterized by means of the following notion of dichotomy [13]:

Definition 1. We say that Eq. (2) has a weak $([h, p], [k, q])$ -dichotomy on J_σ , iff there exist a constant K and a projection matrix P such that

$$(4) \quad \begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq Kh(t)p(s), & t_0 - \sigma \leq s \leq t, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq Kk(t)q(s), & t_0 - \sigma \leq t \leq s, \end{aligned}$$

and

$$(5) \quad \begin{cases} h(t)p(s) \leq Ck(t)q(s), & t \geq s \geq t_0 - \sigma, \\ k(t)q(s) \leq Ch(t)p(s), & s \geq t \geq t_0 - \sigma, \end{cases}$$

where $C \geq 1$ is a constant.

Remark 1. If $h = k, p = q$, then we say that Eq. (1) possesses an $[h, p]$ -dichotomy. If Eq. (1) has an $([h, p], [k, q])$ -dichotomy, then condition (5) implies that Eq. (1) has an $[h, p]$ -dichotomy, and also a $[k, q]$ -dichotomy, each with the same projection matrix P and constant CK .

Remark 2. If $p(t) = h^{-1}(t), q(t) = k^{-1}(t)$ [19], [15], [16], then we say that Eq. (1) possesses an (h, k) -dichotomy. In this case the hypotheses (5) reduce to the requirement:

$$(6) \quad h(t)h^{-1}(s) \leq Ck(t)k^{-1}(s), \quad t \geq s \geq t_0 - \sigma.$$

The case of an (h, h) -dichotomy will be termed an h -dichotomy. If Eq. (1) possesses an (h, k) -dichotomy, then (6) implies that Eq. (1) has an h -dichotomy, and also has a k -dichotomy, each with the same projection matrix P and constant CK .

We call the attention to the use of square brackets to denote an $[h, p]$ -dichotomy. This is made deliberately to distinguish this dichotomy from the notation of the (h, k) -dichotomies in the sense of Pinto [19], where we use parentheses instead.

We will use the following subspaces of initial conditions to Eq. (2):

$$\mathbf{V}_h = \{\xi \in \mathbf{V}: \Phi(t)\xi \in C_h\}, \quad \mathbf{V}_{h,0} = \{\xi \in \mathbf{V}_h: \lim_{t \rightarrow \infty} h(t)^{-1}\Phi(t)\xi = 0\}.$$

The forthcoming Theorem A follows in a way similar to the proof of Proposition 2.2 in [4] (see also [15], [17], [14]), and Theorem B follows from a result on admissibility of a pair of functional spaces [4], [14].

Theorem A. *Let us assume that Eq. (2) has the weak dichotomy (4)–(5), then Eq. (2) has an $([h, p], [k, q])$ -dichotomy with a projection Q , iff*

$$\mathbf{V}_{k,0} \subset Q[\mathbf{V}] \subset \mathbf{V}_h.$$

Theorem B. *If the system*

$$x'(t) = A(t)x(t)$$

has a weak $[h, p]$ -dichotomy and $hpB \in L^1$, then the system

$$y'(t) = [A(t) + B(t)]y(t)$$

has a weak $[h, p]$ -dichotomy.

Throughout, the functions h, k will be assumed to have a bounded growth.

Definition 2. We say that the function $h: J_\sigma \rightarrow (0, \infty)$ is of class $G_{\sigma, M}$, for a positive number M , iff

$$h(s)h^{-1}(t) \leq M, \quad s \in [t - \sigma, t + \sigma], \quad t \geq t_0.$$

We will use the following definitions of instability:

Definition 3. We say that the null solution of Eq. (2) is h -stable on the interval J , iff for every $\varepsilon > 0$ there exists a $\delta > 0$ ($\delta = \delta(t_0, \varepsilon)$) such that if $\varphi \in C([-\sigma, 0])$ and $|\varphi|_\sigma < \delta$, then the solution $x(t; t_0, \varphi)$ exists on all J and $h^{-1}(t)|x(t; t_0, \varphi)| < \varepsilon$ for all $t \geq t_0$. In addition to the above property, if for every $|\varphi|_\sigma < \delta$ we have

$$(7) \quad \lim_{t \rightarrow \infty} h^{-1}(t)x(t; t_0, \varphi) = 0,$$

then the null solution of Eq. (2) is called h -asymptotically stable.

Definition 4. We say that the null solution of Eq. (2) is h -unstable on the interval J , iff there exists an $\varepsilon > 0$ such that for every $\delta > 0$ there exist an initial value function $\varphi_\delta \in C([-\sigma, 0])$, $|\varphi_\delta|_\sigma < \delta$, and a $\tau_\delta \geq t_0$ such that $|x(\tau_\delta; t_0, \varphi_\delta)| \geq \varepsilon$.

3. INSTABILITY UNDER INTEGRABLE CONDITIONS

It is convenient to write Eq. (1) in the equivalent form

$$(8) \quad x'(t) = (A(t) + B(t))x(t) + B(t)(x(t - r(t)) - x(t)).$$

Regarding this equation, for $t \geq t_0$ let us define the operator

$$\begin{aligned} \mathcal{U}[x](t) &= \int_{t_0}^t \Phi(t)P\Phi^{-1}(s)B(s)(x(s - r(s)) - x(s)) \, ds \\ &\quad - \int_t^\infty \Phi(t)(I - P)\Phi^{-1}(s)B(s)(x(s - r(s)) - x(s)) \, ds. \end{aligned}$$

We call \mathcal{U} the dichotomic operator associated to Eq. (1). Note that \mathcal{U} applies to functions of the space $C(J_\sigma)$. Since the function $\mathcal{U}[y]$ is not defined on $[t_0 - \sigma, t_0]$, we complete this definition in the following manner:

$$\mathcal{T}[x](t) = \begin{cases} \mathcal{U}[x](t_0), & t \in [t_0 - \sigma, t_0], \\ \mathcal{U}[x](t), & t \geq t_0. \end{cases}$$

Lemma 1. *If Eq. (2) has the dichotomy (4)–(5), where $h \in G_{\sigma, M}$, then $hpB \in L^1$ implies that the operator $\mathcal{T}: C_h(J_\sigma) \rightarrow C_h(J_\sigma)$ is continuous:*

$$(9) \quad |\mathcal{T}[x]|_h \leq 2KCM^2 \int_{t_0}^\infty h(s)p(s)|B(s)| \, ds |x|_h.$$

Moreover, if

$$(10) \quad 2KCM^2 \int_{t_0}^\infty h(s)p(s)|B(s)| \, ds < 1,$$

then \mathcal{T} acts as a contraction.

Proof. For $t \geq t_0$ we have the estimate

$$\begin{aligned} |h^{-1}(t)\mathcal{T}[x](t)| &\leq K \int_{t_0}^t p(s)|B(s)||x(s - r(s)) - x(s)| \, ds \\ &\quad + K \int_t^\infty h^{-1}(t)k(t)q(s)|B(s)||x(s - r(s)) - x(s)| \, ds \\ &\leq KC(M + 1) \int_{t_0}^\infty h(s)p(s)|B(s)| \, ds |x|_h. \end{aligned}$$

For $t \in [t_0 - \sigma, t_0]$ we may write

$$\begin{aligned} |h^{-1}(t)\mathcal{T}[x](t)| &= |h^{-1}(t)h(t_0)h^{-1}(t_0)\mathcal{T}[x](t_0)| \\ &\leq KCM(M + 1) \int_{t_0}^\infty h(s)p(s)|B(s)| \, ds |x|_h. \end{aligned}$$

From these estimates the assertion of the lemma follows, because $M \geq 1$. □

Theorem 1. *If Eq. (2) has the dichotomy (4)–(5), where the function h is of class $G_{\sigma,M}$, and (10) is satisfied, then the null solution of Eq. (1) is h -unstable if $\mathbf{V}_h \neq \mathbf{V}$.*

Proof. Let us assume the contrary, then for $\varepsilon > 0$ there exists a $\delta > 0$ such that $\varphi \in C[-\sigma, 0]$, $|\varphi|_\sigma < \delta$ imply $|h^{-1}(t)x(t; t_0, \varphi)| < \varepsilon, \forall t \geq t_0$. Let φ be an initial value function satisfying

$$(11) \quad \varphi = \text{const.} = x_0, \quad 0 < |x_0| < \delta, \quad x_0 \in (I - P)[\mathbf{V}].$$

We will show that $x(t; t_0, \varphi)$ is not h -bounded. This is enough to accomplish the proof of the theorem. We define

$$y(t) = x(t; t_0, \varphi) - \mathcal{T}[x(\cdot; t_0, \varphi)](t), \quad t \geq t_0 - \sigma.$$

By Lemma 1 the function $y(t)$ belongs to $C_h(J_\sigma)$. Besides, $y(t)$ is a solution of Eq. (2) on the interval $[t_0, \infty)$. Hence $y(t_0) \in \mathbf{V}_h$. Due to Theorem A we may assume that $y(t_0) \in P[\mathbf{V}]$. Moreover, we have

$$y(t_0) = x_0 + (I - P) \int_{t_0}^{\infty} \Phi^{-1}(s)B(s)(x(s - r(s)) - x(s)) ds \in (I - P)[\mathbf{V}],$$

implying $y(t_0) = 0$, and consequently $y(t) = 0, \forall t \geq t_0$. But in this case $x(\cdot; t_0, \varphi)$ satisfies the integral equation

$$x(t; t_0, \varphi) = \mathcal{T}(x(\cdot, t_0, \varphi))(t), \quad t \geq t_0,$$

whence

$$x(t; t_0, \varphi) = \mathcal{T}(x(\cdot, t_0, \varphi))(t), \quad t \geq t_0 - \sigma.$$

Thus, any solution $x(\cdot; t_0, \varphi)$, where φ satisfies (11), is a fixed point of the dichotomic operator $\mathcal{T}: C_h(J_\sigma) \rightarrow C_h(J_\sigma)$. But condition (10) implies that the operator \mathcal{T} is a contraction. Since \mathcal{T} is linear, we have $x(\cdot; t_0, \varphi) = 0$, which yields the contradiction $\varphi(0) = x(t_0; t_0, \varphi) = x_0 = 0$. \square

If the function h is bounded away from null ($h(t) \geq \alpha > 0, \forall t$), then the conditions of Theorem 1 imply that the solutions of Eq. (1) are unstable in the sense of Liapounov.

Corollary 1. *If Eq. (2) has the dichotomy (4)–(5), where the function k is of class $G_{\sigma,M}$, and*

$$2KCM^2 \int_{t_0}^{\infty} k(s)q(s)|B(s)| ds < 1$$

is satisfied, then the null solution of Eq. (1) is k -unstable if $\mathbf{V}_k \neq \mathbf{V}$.

Proof. Condition (5) implies that Eq. (2) has a $[k, q]$ -dichotomy. The rest of the proof follows in the same way as that of Theorem 1. \square

If the function $B(t)$ is integrable then the equation

$$y'(t) = (\text{diag}\{-1, t^{-1}, t\} + B(t))y(t), \quad t \geq t_0 = 1,$$

has an (e^{-t}, t) -dichotomy. According to Theorem 1 the null solution of

$$(12) \quad x'(t) = \text{diag}\{-1, t^{-1}, t\}x(t) + B(t)x(t - r(t))$$

is e^{-t} -unstable for every bounded delay $r(t)$. This does not imply the Liapounov instability. Nevertheless, Eq. (12) has a k -dichotomy with $k(t) = t$. The condition $\mathbf{V}_k \neq \mathbf{V}$ is certainly satisfied, therefore Corollary 1 yields the Liapounov instability of Eq. (12).

The proof of Theorem 1 shows that no solution $x(t; t_0, \varphi)$ of Eq. (1) satisfying (11) is h -bounded. A natural question arises: For which initial value functions may we expect that the solution $x(t; t_0, \varphi)$ is not h -bounded? To answer this question, let us consider the set $\mathcal{S} \subset C[-\sigma, 0]$ of initial value functions defined by the following properties:

$$(13) \quad |h_{t_0}^{-1}\varphi|_{\sigma} = |h^{-1}(t_0)\varphi(0)| \neq 0, \quad \varphi(0) \in (I - P)[\mathbf{V}].$$

Theorem 2. *Under the conditions of Theorem 1, every solution $x(t; t_0, \varphi)$, with $\varphi \in \mathcal{S}$ is h -unbounded.*

Proof. Assume that $\varphi \in \mathcal{S}$ and $x(t; t_0, \varphi)$ is h -bounded. By repeating the first lines of the proof of Theorem 1 we obtain

$$x(t; t_0, \varphi) = \mathcal{T}(x(\cdot; t_0, \varphi))(t), \quad t \geq t_0.$$

From this identity and properties (9), (13) we conclude

$$\sup_{[t_0, \infty)} |h^{-1}(t)x(\cdot; t_0, \varphi)| \leq 2KCM^2 |hpB|^1 |x(\cdot; t_0, \varphi)|_h.$$

Since $\varphi \in \mathcal{S}$, the last estimate implies

$$|x(\cdot; t_0, \varphi)|_h \leq 2KCM^2 |hpB|^1 |x(\cdot; t_0, \varphi)|_h,$$

whence $x(t; t_0, \varphi) = 0$, $t \geq t_0 - \sigma$, because of condition (10). Hence $\varphi = 0$, a contradiction with $\varphi \in \mathcal{S}$. \square

If the condition $\mathbf{V}_h \neq \mathbf{V}$ of Theorem 1 is not satisfied, then the following theorem provides the answer to the problem of instability:

Theorem 3. *If Eq. (2) has the dichotomy (4)–(5), where the function h is of class $G_{\sigma, M}$, and (10) is satisfied, then the null solution of Eq. (1) is not asymptotically h -stable if $\mathbf{V}_{h,0} \neq \mathbf{V}_h$ (respectively, the null solution of Eq. (1) is not asymptotically k -stable if $\mathbf{V}_{k,0} \neq \mathbf{V}_k$ and $2KCM^2|kqB|^1 < 1$).*

Proof. According to Remark 1, we may handle the dichotomy (4)–(5) as an $[h, p]$ -dichotomy. Let us assume that the null solution of Eq. (1) is asymptotically h -stable. Then for $\varepsilon = 1$ there exists a positive δ such that $|\varphi|_\sigma < \delta$ implies $|h^{-1}(t)x(t; t_0, \varphi)| < 1$, $t \geq t_0$, and (7) is satisfied. Let ϱ be a positive number such that $\varrho|h_{t_0}|_\sigma < \delta$ and let γ be a positive number such that

$$\gamma + 2KCM^2|hpB|^1\varrho \leq \varrho.$$

Fixing a vector $y_0 \in \mathbf{V}_h \setminus \mathbf{V}_{h,0}$ with the property $|\Phi y_0|_h < \gamma$, we introduce an operator \mathcal{F} defined by

$$(14) \quad \mathcal{F}[x](t) = \Phi(t)y_0 + \mathcal{T}[x](t), \quad t \geq t_0 - \sigma.$$

Due to the choice of γ we have the property

$$\mathcal{F}: B_h[0, \varrho] \rightarrow B_h[0, \varrho].$$

By virtue of (10) the operator \mathcal{F} is contractive in this ball. Let x be the unique fixed point of \mathcal{F} in $B_h[0, \varrho]$. This function is a solution of Eq. (1). Due to Theorem A we may assume that the projection P defining the $[h, p]$ -dichotomy satisfies

$$\lim_{t \rightarrow \infty} h^{-1}(t)\Phi(t)P = 0.$$

This property implies the identity

$$(15) \quad x(t) = \Phi(t)y_0 + o(h)(t),$$

where $o(h)$ denotes a function satisfying $\lim_{t \rightarrow \infty} h^{-1}(t)o(h)(t) = 0$. As $x = \mathcal{F}[x]$, the estimate (9), for $t \in [t_0 - \sigma, t_0]$, yields

$$|x(t)| \leq h(t)|\Phi y_0|_h + h(t)|\mathcal{T}[x]|_h \leq (\gamma + 2KCM^2|hpB|^1\varrho)|h_{t_0}|_\sigma \leq \varrho|h_{t_0}|_\sigma < \delta.$$

Therefore $|x_{t_0}| < \delta$, implying $x(t) = o(h)(t)$. However

$$\lim_{t \rightarrow \infty} h^{-1}(t)\Phi(t)y_0 \neq 0.$$

The last relation and (15) are contradictory. □

The forthcoming Theorem 4 and Theorem 5 are consequences of Theorem 1 and Theorem 3, respectively.

Theorem 4. *If Eq. (2) has an (h, k) -dichotomy, where the function h is of class $G_{\sigma, M}$, and*

$$(16) \quad 2KCM^2 \int_{t_0}^{\infty} |B(s)| \, ds < 1,$$

then the null solution of Eq. (1) is h -unstable if $\mathbf{V}_h \neq \mathbf{V}$.

Theorem 5. *If Eq. (2) has an (h, k) -dichotomy, where the function h is of class $G_{\sigma, M}$ and (16) is satisfied, then the null solution of Eq. (1) is not asymptotically h -stable if $\mathbf{V}_{h,0} \neq \mathbf{V}_h$.*

4. INSTABILITY FOR NONINTEGRABLE COEFFICIENTS

The previous result is of a limited interest, since the condition (10) does not involve the time lag function $r(t)$. Let $h \in G_{\sigma, M}$. In order to incorporate the properties of $r(t)$ into the statements of our theorems, following the ideas in [8], we introduce the set \mathcal{M}_h consisting of the functions belonging to $C_h(J_\sigma)$ such that

$$(\mathbf{M}_h) \quad h^{-1}(t)|x(t) - x(t')| \leq M\beta(t)(t - t')|x|_h, \quad t - \sigma \leq t' \leq t, \quad t \geq t_0, \quad t' \geq t_0,$$

where, for convenience, the constant M is the same as in Definition 2, and the function β is defined by

$$\beta(t) = \max\{1, |(A + B)_t|_\sigma\}.$$

By standard arguments, we can prove that \mathcal{M}_h is a closed set in $C_h(J_\sigma)$.

Lemma 2. *Assume that Eq. (2) has the dichotomy (4)–(5), $h \in G_{\sigma, M}$. If*

$$(17) \quad 2KCM^2\{|\beta r h p B|^1 + |\beta r h p B|^\infty\} < 1,$$

then

$$(18) \quad \mathcal{T}: \mathcal{M}_h \rightarrow \mathcal{M}_h.$$

P r o o f. The definition of the operator \mathcal{T} implies, for $t \geq t_0$ and $x \in C_h(J_\sigma)$, that

$$\begin{aligned} |h^{-1}(t)\mathcal{T}[x](t)| &\leq KC \int_{t_0}^{\infty} p(s)|B(s)||x(s-r(s))-x(s)| ds \\ &\leq KCM \int_{t_0}^{\infty} \beta(s)r(s)h(s)p(s)|B(s)| ds|x|_h. \end{aligned}$$

On the other hand, for $t \in [t_0 - \sigma, t_0]$ we have

$$\begin{aligned} |h^{-1}(t)\mathcal{T}[x](t)| &= |h^{-1}(t)h(t_0)h^{-1}(t_0)\mathcal{T}[x](t_0)| \\ &\leq KCM^2 \int_{t_0}^{\infty} \beta(s)r(s)h(s)p(s)|B(s)| ds|x|_h. \end{aligned}$$

Hence, we have proved that condition (17) implies $\mathcal{T}: C_h(J_\sigma) \rightarrow C_h(J_\sigma)$.

In order to verify the property $[\mathbf{M}_h]$, we write

$$\mathcal{T}[x](t) - \mathcal{T}[x](t') = I_1(t) + I_2(t) + I_3(t) - I_4(t), \quad t - \sigma \leq t' \leq t,$$

where

$$\begin{aligned} I_1(t) &= \int_{t'}^t \Phi(t)P\Phi^{-1}(s)B(s)(x(s) - x(s-r(s))) ds, \\ I_2(t) &= \int_{t_0}^{t'} [\Phi(t) - \Phi(t')]P\Phi^{-1}(s)B(s)(x(s) - x(s-r(s))) ds, \\ I_3(t) &= \int_{t'}^t \Phi(t')(I - P)\Phi^{-1}(s)B(s)(x(s) - x(s-r(s))) ds, \\ I_4(t) &= \int_t^{\infty} [\Phi(t) - \Phi(t)](I - P)\Phi^{-1}(s)B(s)(x(s) - x(s-r(s))) ds. \end{aligned}$$

From condition (4) we have

$$\begin{aligned} (19) \quad |h^{-1}(t)I_1(t)| &\leq KM \int_{t'}^t h(s)p(s)|B(s)|\beta(s)r(s) ds|x|_h \\ &\leq KM\beta(t)|\beta r h p B|^\infty(t-t')|x|_h. \end{aligned}$$

On the interval $s \leq t' \leq t$, $t-t' \leq \sigma$, we obtain the estimate

$$\begin{aligned} (20) \quad |(\Phi(t) - \Phi(t'))P\Phi^{-1}(s)| &= \left| \int_{t'}^t (A+B)(u)\Phi(u) du P\Phi^{-1}(s) \right| \\ &\leq K\beta(t) \int_{t'}^t h(u)p(s) du \\ &\leq KM\beta(t)h(t)p(s)(t-t'). \end{aligned}$$

Inserting the estimate (20) into the definition of function I_2 , we may write

$$(21) \quad |h^{-1}(t)I_2(t)| \leqslant K M^2 \beta(t) \int_{t_0}^{\infty} \beta(s) h(s) p(s) |B(s)| r(s) \, ds (t - t') |x|_h.$$

Further, we have

$$(22) \quad |h^{-1}(t)I_3(t)| \leqslant K C M \left(\int_{t'}^t h(t') h^{-1}(t) h(s) p(s) \beta(s) r(s) |B(s)| \, ds \right) |x|_h \\ \leqslant K C M \beta(t) |h p \beta r B|^\infty (t - t') |x|_h.$$

Finally, relying on (20) we conclude

$$(23) \quad |h^{-1}(t)I_4(t)| \leqslant K C M^2 \beta(t) |h p \beta r B|^1 (t - t') |x|_h.$$

From (19), (21), (22) and (23) we obtain

$$|\mathcal{S}[x](t) - \mathcal{S}[x](t')| \leqslant 2 K C M^2 \beta(t) h(t) (|h p \beta r B|^\infty + |h p \beta r B|^1) (t - t') |x|_h.$$

Therefore condition (17) implies (18). □

Lemma 3. *If $h \in G_{\sigma, M}$ and $y_0 \in \mathbf{V}_h$, then $\Phi y_0 \in \mathcal{M}_h$.*

Proof. We have to verify the property $[\mathbf{M}_h]$. If $t - \sigma \leqslant t' \leqslant t$, $t \geqslant t_0$, then

$$h^{-1}(t) |\Phi(t)y_0 - \Phi(t')y_0| = \left| h^{-1}(t) \int_{t'}^t (A(s) + B(s)) \Phi(s) y_0 \, ds \right| \\ \leqslant h^{-1}(t) \int_{t'}^t |A(s) + B(s)| h(s) h^{-1}(s) |\Phi(s) y_0| \, ds \\ \leqslant M \beta(t) (t - t') |\Phi y_0|_h,$$

whence the proof of the lemma follows. □

Theorem 6. *Let us assume that Eq. (1) has the dichotomy (4)–(5) and the functions h, k are of class $G_{\sigma, M}$. If*

$$(24) \quad 2 K C M^2 \{ |k q \beta r B|^1 + |k q \beta r B|^\infty \} < 1$$

and $\mathbf{V}_h \neq \mathbf{V}_k$, then the null solution of Eq. (1) is h -unstable.

Proof. Let us assume that the null solution of Eq. (1) is h -stable. Then for $\varepsilon = 1$ there exists a $\delta > 0$ such that $|x(\cdot, t_0, \varphi)|_h < 1$ if $|\varphi|_\sigma < \delta$. Let ϱ be a small number such that

$$(25) \quad \varrho |k_{t_0}|_\sigma < \delta.$$

For a small $\gamma > 0$ satisfying

$$\gamma + 2KCM^2\{|kq\beta rB|^1 + |qk\beta rB|^\infty\}\varrho \leq \varrho$$

we fix an initial condition $y_0 \in \mathbf{V}_k \setminus \mathbf{V}_h$ such that

$$|y(\cdot, t_0, y_0)|_k \leq \gamma.$$

Let us consider the integral equation $x = \mathcal{F}[x]$, where the operator \mathcal{F} is defined by (14). From Lemma 2 and Lemma 3 (where instead of h we put k) we have $\mathcal{F}: \mathcal{M}_k \rightarrow \mathcal{M}_k$. Let $B_k^*[0, \varrho]$ be the closed ball with center $x = 0$ and radius ϱ contained in \mathcal{M}_k . In view of the choice of the number γ we have

$$\mathcal{F}: B_k^*[0, \varrho] \rightarrow B_k^*[0, \varrho].$$

In virtue of (17), the operator \mathcal{F} contracts the points of the ball $B_k^*[0, \varrho]$. Let x be a fixed point of the operator \mathcal{F} . A straightforward calculation shows that x is a solution of Eq. (1). Moreover, $t \in [t_0 - \sigma, t_0]$ and (25) yield

$$|x(t)| \leq k(t)(|\Phi y_0|_k + |\mathcal{T}[x]|_k) \leq (\gamma + 2KCM^2|kqB|^1\varrho)|k_{t_0}|_\sigma \leq \varrho|k_{t_0}|_\sigma < \delta,$$

implying that x is an h -bounded function. Therefore $\mathcal{T}[x]$ is h -bounded. The last assertion is proved in the following way: the second estimate in the proof of Lemma 2 shows that this is certainly satisfied if $\beta r h p B \in L^1[t_0, \infty)$, but this follows from (24), because $h p \leq C k q$. Since

$$x(t) = y(t, t_0, y_0) + \mathcal{T}[x](t),$$

we obtain that the function $y(\cdot, t_0, y_0)$ must be h -bounded. However, this contradicts the choice of y_0 . □

The forthcoming Lemma 4 and Theorem 7 follow from Lemma 2 and Theorem 6 in the particular case of a (h, k) -dichotomy.

Lemma 4. *Assume that Eq. 2 has an (h, k) -dichotomy and $h \in G_{\sigma, M}$. If*

$$(26) \quad 2KCM^2\{|\beta r B|^1 + |\beta r B|^\infty\} < 1,$$

then

$$\mathcal{T}: \mathcal{M}_h \rightarrow \mathcal{M}_h.$$

Theorem 7. *Let us assume that Eq. (2) has an (h, k) -dichotomy and the functions h, k belong to $G_{\sigma, M}$. If condition (26) is satisfied and $\mathbf{V}_h \neq \mathbf{V}_k$, then the null solution of Eq. (1) is h -unstable.*

Finally, before we start the section of examples, we emphasize that the stability analysis of Eq. (1) via an appropriate ordinary differential equation is not new. It was used, for example, by Cooke [2], and recently by Györi and Pituk in an interesting paper [8].

5. APPLICATIONS

We present three examples of independent interest.

5.1. Instability of $x^{(n)}(t) = b(t)x(t - r(t))$.

To start, let us begin with the second order equation

$$(27) \quad x''(t) = b(t)x(t - r(t)), \quad 0 \leq r(t) \leq 1, \quad t \geq t_0 = 1,$$

whose vectorial form is

$$\begin{pmatrix} y(t) \\ x(t) \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} y(t) \\ x(t) \end{pmatrix} + b(t) \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} y(t - r(t)) \\ x(t - r(t)) \end{pmatrix}.$$

If $|(u, v)| = |u| + |v|$ is the norm in \mathbb{R}^2 , then the linear system

$$\begin{pmatrix} y(t) \\ x(t) \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} y(t) \\ x(t) \end{pmatrix}$$

with the fundamental matrix $\Phi(t) = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ has the weak dichotomy

$$\begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq 2s, & 1 - \sigma \leq s \leq t, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq 2t, & 1 - \sigma \leq t \leq s, \end{aligned} \quad P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

satisfying (5) with functions $h(t) = 1$, $p(t) = t$, $k(t) = t$, $q(t) = 1$, and constant $C = 1$. If the function $tb(t)$ is integrable, then, according to Theorem B, the perturbed equation

$$x''(t) = b(t)x(t)$$

has a weak dichotomy of type $([1, t], [t, 1])$. According to Theorem 1, the null solution of Eq. (27) is unstable if $tb(t) \in L^1$, because $\mathbf{V}_h \neq \mathbf{V}$. Since $\mathbf{V}_k \neq \mathbf{V}_{k,0}$, even Theorem 3 is applicable to this example if $tb(t) \in L^1$; the null solution of Eq. (27)

is t -asymptotic unstable. In this example, the property $\mathbf{V}_h \neq \mathbf{V}_k$ suggests the possibility of applying Theorem 6, for which we require the sufficient condition

$$(28) \quad \lim_{t \rightarrow \infty} \left(\int_t^\infty |s \max\{1, |b(s)|\} r(s) b(s)| ds + |t \max\{1, |b(t)|\} r(t) b(t)| \right) = 0$$

equivalent to

$$g \in L^1, \quad \lim_{t \rightarrow \infty} g(t) = 0, \quad g(t) := \max\{1, |b(t)|\} r(t) b(t),$$

in order to establish property (24). Under this condition the null solution of Eq. (27) is unstable in the sense of Liapounov.

The general equation

$$x^{(n)}(x) = b(x)x(t - r(t))$$

can be treated similarly, the linear system

$$x'(t) = (I + I_1)x(t), \quad I_1 = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}$$

having the family of weak dichotomies

$$\begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq Kt^{m-1}s^m, \quad t \geq s \geq 1, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq Kt^m s^{m-1}, \quad s \geq t \geq 1, \end{aligned}$$

where

$$P = \text{diag}\{\overbrace{1, \dots, 1}^m, 0, \dots, 0\}, \quad 0 < m < n.$$

5.2 Euler equations.

As a second example we study the instability of the Euler equation

$$(29) \quad y'' + \frac{\mu}{t^2}y = b(t)y(t - r(t)),$$

where μ is a real parameter satisfying $|\mu| < 1/4$ (the case $|\mu| \geq 1/4$ can be dealt with similarly as we do in the text). In this case the fundamental matrix corresponding to the linear equation

$$y'' + \frac{\mu}{t^2}y = 0$$

has the form

$$(30) \quad \Phi(t) = \begin{pmatrix} t^{\lambda_-} & t^{\lambda_+} \\ \lambda_- t^{\lambda_- - 1} & \lambda_+ t^{\lambda_+ - 1} \end{pmatrix}, \quad \lambda_{\mp} = \frac{1}{2} \mp \sqrt{\frac{1}{4} - \mu}.$$

For the projection matrix $P = \text{diag}\{1, 0\}$ we have the following estimates satisfying (4)–(5):

$$(31) \quad \begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq Kt^{\lambda_-}s^{\lambda_+}, \quad t \geq s, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq Kt^{\lambda_+}s^{\lambda_-}, \quad s \geq t. \end{aligned}$$

We aim at applying Theorem 3. For $h(t) = t^{\lambda_-}$, $k(t) = t^{\lambda_+}$, we observe that $\mathbf{V}_k \neq \mathbf{V}_{k,0}$. By Theorem B, the equation

$$y'' + \frac{\mu}{t^2}y = b(t)y$$

has a $[t^{\lambda_+}, t^{\lambda_-}]$ -dichotomy if $tb(t) \in L^1$. According to Theorem 3 we have that the null solution of Eq. (29), under condition $tb(t)$, is not asymptotically t^{λ_+} -stable (this implies the Liapounov instability). Such a consequence cannot be obtained if $b(t) = O(t^{-2})$. Let us study this case more carefully. Assume the existence of a constant ν such that the function $(\nu - t^2b(t))t^{-1}$ is integrable. Then Eq. (29) can be written in the form

$$(32) \quad x''(t) + \frac{\mu - \nu}{t^2}x(t) + \frac{\nu - t^2b(t)}{t^2}x(t) = b(t)(x(t - r(t)) - x(t)).$$

If $4(\mu - \nu) < 1$, then the equation

$$\begin{pmatrix} y(t) \\ x(t) \end{pmatrix}' = \frac{\mu - \nu}{t^2} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} y(t) \\ x(t) \end{pmatrix}$$

has the dichotomy (31), with a fundamental matrix (30) defined with values λ_{\mp} , where instead of μ it is necessary to write $\mu - \nu$. By Theorem B, the dichotomy (31) is preserved if $(\nu - t^2b(t))t^{-1}$ is integrable. Now Theorem 6 will apply to Eq. (32) if the condition (24) is fulfilled. In the present example a sufficient condition to obtain (24) is (28).

An example of a function satisfying these conditions is $b(t) = 1/(1 + t^2)$, with $\nu = 1$. Note that $tb(t)$ is not integrable.

In examples regarding Euler equations it is worth mentioning that the classical change of the time scale $t = e^{\tau}$ in Eq. (29) would not work, since it is not clear how to define the function $y(\tau) = x(e^{\tau})$ from $x(t - r(t))$.

5.3. Instability of $x'(t) = Ax(t) + B(t)x(t - r(t))$.

Let us consider the system

$$(33) \quad x'(t) = Ax(t) + B(t)x(t - r(t)),$$

where the function A is constant. Throughout, $\sigma(A)$ will denote the set of eigenvalues of the matrix A ; $\sigma_+ = \{\lambda \in \sigma(A) : \operatorname{Re} \lambda > 0\}$. If $\sigma_+(A) \neq \emptyset$, we define numbers α, ν such that

$$0 < \alpha < \mu - \nu < \nu < \mu = \min\{\operatorname{Re} \lambda : \operatorname{Re} \lambda > 0\}.$$

The change of the variable

$$(34) \quad x(t) = e^{\nu t} z(t) = e^{(\nu - \alpha)t} e^{\alpha t} z(t),$$

reduces Eq. (33) to the form

$$(35) \quad z'(t) = (A - \nu I + e^{-r(t)} B(t))z(t) + e^{-r(t)} B(t)(z(t - r(t)) - z(t)),$$

the system

$$(36) \quad u'(t) = (A - \nu I)u(t)$$

having an $(e^{-\alpha t}, e^{\beta t})$ -dichotomy. If $B \in L^1$, then the system

$$(37) \quad u'(t) = (A - \nu I + e^{-r(t)} B(t))u(t)$$

has the same $(e^{-\alpha t}, e^{\beta t})$ -dichotomy [4], [15]. It is clear that the function $e^{-\alpha t}$ is of class $G_{\sigma, M}$ for some positive constant M . All conditions of Theorem 1 are fulfilled for large values of the initial moment t_0 . Thus, the null solution $z = 0$ is $e^{-\alpha t}$ -unstable, which implies, in view of (34), the Liapounov instability of the solutions of (33). We may write the above outcomes in

Theorem 8. *If $\sigma_+(A)$ is not empty and $B(t)$ is integrable, then for any bounded delay function $r(t)$, the null solution of Eq. (1) is unstable.*

A more interesting result is obtained for a nonintegrable and bounded function $B(t)$. First we recall a known result on the stability of exponential dichotomies [4], [16]:

Theorem C. *Let us assume that Eq. (36) allows the exponential dichotomy*

$$(38) \quad \begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq K_1 e^{-\alpha(t-s)}, & t_0 - \sigma \leq s \leq t, & \alpha > 0, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq K_2 e^{\beta(t-s)}, & t_0 - \sigma \leq t \leq s, & \beta > 0. \end{aligned}$$

Then the condition

$$(39) \quad |e^{-r(\cdot)}B|^\infty \left\{ \frac{K_1}{\alpha} + \frac{K_2}{\beta} \right\} < \frac{1}{2}$$

implies that Eq. (37) has the exponential dichotomy

$$(40) \quad \begin{aligned} |\Phi(t)P\Phi^{-1}(s)| &\leq Ke^{-\alpha_1(t-s)}, & t_0 - \sigma \leq s \leq t, \\ |\Phi(t)(I - P)\Phi^{-1}(s)| &\leq Ke^{\beta_1(t-s)}, & t_0 - \sigma \leq t \leq s, \end{aligned}$$

where K is a constant and $0 < \alpha_1 < \alpha$, $0 < \beta_1 < \beta$.

We desire to apply Theorem 6 to Eq. (35), where Eq. (36) has the dichotomy (38). If (39) is fulfilled, then Eq. (37) has the dichotomy (40). The condition (17) requires the estimate

$$|\beta r(\cdot)e^{-r(\cdot)}B|^\infty \leq |r(\cdot)e^{-r(\cdot)}B|^\infty (1 + |A| + |e^{-r(\cdot)}B|^\infty).$$

For the functions $h(t) = e^{-\alpha_1 t}$, $k(t) = e^{\beta_1 t}$, the constant M in Definition 2 can be chosen to be $M = e^{\max\{\alpha, \beta\}\sigma}$. The property (5) is established with $C = 1$. Thus, condition (17) is satisfied if

$$2Ke^{2\alpha_1\sigma}(1 + |A| + |e^{-r(\cdot)}B|^\infty)(|re^{-r(\cdot)}B|^1 + |r(\cdot)e^{-r(\cdot)}B|^\infty) < 1.$$

The condition $\mathbf{V}_h \neq \mathbf{V}_k$ of Theorem 6 is certainly satisfied. Note that α_1 can be taken as close to null as desired. Thus, this last condition follows from

$$(41) \quad 2K(1 + |A| + |e^{-r(\cdot)}B|^\infty)(|r(\cdot)e^{-r(\cdot)}B|^1 + |r(\cdot)e^{-r(\cdot)}B|^\infty) < 1.$$

From Theorem 6 we obtain that conditions (39), (41) imply the $e^{-\alpha_1 t}$ -instability of Eq. (35). Since $\alpha_1 < \alpha$, then the null solution of Eq. (35) is $e^{-\alpha t}$ -unstable, implying, according to (34), the instability of Eq. (33). From these conclusions we obtain

Theorem 9. *If the matrix A allows an eigenvalue with a positive real part and $B(t) \sim C_1 t^{-\gamma_1}$, $r(t) \sim C_2 t^{-\gamma_2}$, where γ_1, γ_2 are nonnegative constants such that $\gamma_1 + \gamma_2 > 1$, then the null solution of the equation with bounded delay (33) is unstable.*

Acknowledgment. The author wishes to express his gratitude to the anonymous referee for his meticulous reading and improvements to the final form of this paper.

References

- [1] *E. A. Coddington and N. Levinson*: Theory of Ordinary Differential Equations. McGill-Hill, New York, 1975.
- [2] *K. L. Cooke*: Functional differential equations close to differential equations. Bull. Amer. Math. Soc. *72* (1966), 285–288.
- [3] *W. A. Coppel*: On the stability of ordinary differential equations. J. London Math. Soc. *39* (1964), 255–260.
- [4] *W. A. Coppel*: Dichotomies in Stability Theory. Lecture Notes in Mathematics Vol. 629. Springer Verlag, Berlin, 1978.
- [5] *L. E. Elsgoltz and S. B. Norkin*: Introduction to the Theory of Differential Equations with Deviating Arguments. Nauka, Moscow, 1971. (In Russian.)
- [6] *J. Gallardo and M. Pinto*: Asymptotic integration of nonautonomous delay-differential systems. J. Math. Anal. Appl. *199* (1996), 654–675.
- [7] *K. Gopalsamy*: Stability and Oscillations in Delay Differential Equations of Populations Dynamics. Kluwer, Dordrecht, 1992.
- [8] *I. Györi and M. Pituk*: Stability criteria for linear delay differential equations. Differential Integral Equations *10* (1997), 841–852.
- [9] *N. Rouche, P. Habets and M. Laloy*: Stability Theory by Liapounov’s Second Method. App. Math. Sciences *22*. Springer, Berlin, 1977.
- [10] *J. K. Hale*: Theory of Functional Differential Equations. Springer-Verlag, New York, 1977.
- [11] *J. K. Hale and S. M. Verduyn Lunel*: Introduction to Functional Differential Equations. Springer-Verlag, New York, 1993.
- [12] *R. Naulin*: Instability of nonautonomous differential systems. Differential Equations Dynam. Systems *6* (1998), 363–376.
- [13] *R. Naulin*: Weak dichotomies and asymptotic integration of nonlinear differential systems. Nonlinear Studies *5* (1998), 201–218.
- [14] *R. Naulin*: Functional analytic characterization of a class of dichotomies. Unpublished work (1999).
- [15] *R. Naulin and M. Pinto*: Roughness of (h, k) -dichotomies. J. Differential Equations *118* (1995), 20–35.
- [16] *R. Naulin and M. Pinto*: Admissible perturbations of exponential dichotomy roughness. J. Nonlinear Anal. TMA *31* (1998), 559–571.
- [17] *R. Naulin and M. Pinto*: Projections for dichotomies in linear differential equations. Appl. Anal. *69* (1998), 239–255.
- [18] *M. Pinto*: Non autonomous semilinear differential systems: Asymptotic behavior and stable manifolds. Preprint (1997).
- [19] *M. Pinto*: Dichotomy and asymptotic integration. Contributions USACH (1992), 13–22.

Author’s address: R. Naulin, Departamento de Matemáticas, Universidad de Oriente, Cumaná 6101-A, Apartado 245, Venezuela, e-mail: rnaulin@sucre.udo.edu.ve.