

Abdelmalek Azizi

Weak multiplication modules

Czechoslovak Mathematical Journal, Vol. 53 (2003), No. 3, 529–534

Persistent URL: <http://dml.cz/dmlcz/127820>

Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

WEAK MULTIPLICATION MODULES

A. AZIZI, Shiraz

(Received June 29, 2000)

Abstract. In this paper we characterize weak multiplication modules.

Keywords: prime submodules, weak multiplication modules, rank of modules

MSC 2000: 13E05

1. INTRODUCTION

Throughout this paper all rings are commutative with identity and all modules are unitary. A proper submodule N of a module M over a ring R is said to be prime (P -prime) if $ra \in N$ for $r \in R$ and $a \in M$ implies that either $a \in N$ or $r \in (N : M) = P$ (see, for example, [4], [6]). The set of all prime submodules in an R -module M is denoted $\text{Spec}_R M$ or $\text{Spec } M$.

Recall that if R is an integral domain with the quotient field K , the rank of an R -module M ($\text{rank } M$ or $\text{rank}_R M$) is defined to be the maximal number of elements of M linearly independent over R . We have $\text{rank } M =$ the dimension of the vector space KM over K , that is $\text{rank } M = \text{rank}_K KM$ ([7]).

An R -module M is called a multiplication module if for every submodule N of M we have $N = IM$, where I is an ideal of R ([3]).

2. WEAK MULTIPLICATION MODULES

Definition. An R -module M is called a *weak multiplication module* if $\text{Spec } M = \emptyset$ or for every prime submodule N of M we have $N = IM$, where I is an ideal of R .

One can easily show that if M is a weak multiplication module, then $N = (N : M)M$ for every prime submodule N of M ([1]).

As is seen in [1], Q is a weak multiplication Z -module which is not a multiplication module.

If R is a ring (not necessarily an integral domain) and M is an R -module, the subset $T(M)$ of M is defined by

$$T(M) = \{m \in M \mid \exists 0 \neq r \in R \text{ such that } rm = 0\}.$$

Obviously, if R is an integral domain, then $T(M)$ is a submodule of M .

It is well known that if R is a ring in which every proper ideal is prime, then R is a field. Compare it with the following result.

Proposition 2.1. *Let R be a ring and $O \neq M$ an R -module, then R is a field if and only if every proper submodule of M is a prime submodule of M and $T(M) \neq M$.*

Proof. \Rightarrow Is obvious.

\Leftarrow Let $a \in M - T(M)$, so $\text{Ann}(a) = O$. In view of the assumption, it is easy to see that every proper submodule of the R -module $M^* = Ra$ is a prime submodule of M^* and $M^* = Ra \cong R$ as R -modules, therefore every proper ideal of R is a prime ideal, hence R is a field. \square

Note. The condition $T(M) \neq M$ in the previous result is necessary. For example, let R be a ring which is not a field and let m be a maximal ideal of R , then for the R -module $M = \frac{R}{m}$ every proper submodule is prime, indeed the only proper submodule of M is $\frac{m}{m}$ which is prime as well.

Lemma 2.2. *Let P be a prime ideal of R , let S be a multiplicatively closed set such that $P \cap S = \emptyset$ and let M be an R -module. Then there exists a one-to-one correspondence between the P -prime submodules of M and the $S^{-1}P$ -prime submodules of $S^{-1}M$.*

Proof. See [5, Proposition 1]. \square

Lemma 2.3. *An R -module M is a weak multiplication module if and only if the R_P -module M_P is a weak multiplication module for every prime (or maximal) ideal P of R .*

Proof. Let M be a weak multiplication R -module and N a prime submodule of M_P where P is a prime ideal of R . According to Lemma 2.2, we know that $N \cap M$ is a prime submodule of M . So $N \cap M = IM$, therefore $N = (N \cap M)_P = I_P M_P$.

Conversely, let N be a prime submodule of M . We show that $(\frac{N}{(N:M)M})_P = O$ for every maximal ideal P .

If $(N : M) \subseteq P$, then by Lemma 2.2, N_P is a prime submodule, so $N_P = (N_P : M_P)M_P$, and by Corollary 1 of [5], $(N_P : M_P) = (N : M)_P$. Hence $\left(\frac{N}{(N:M)M}\right)_P = \frac{N_P}{(N:M)_P M_P} = \frac{N_P}{(N_P:M_P)M_P} = O$. If $(N : M) \not\subseteq P$, then clearly $N_P = M_P$ and $(N : M)_P = R_P$, so obviously

$$\left(\frac{N}{(N : M)M}\right)_P = \frac{N_P}{(N : M)_P M_P} = \frac{M_P}{M_P} = O.$$

□

Proposition 2.4. *If M is a weak multiplication module over an integral domain, then*

- (i) *If M is a non-zero torsion-free module, then $\text{rank } M = 1$.*
- (ii) *If M is a torsion module, then $\text{rank } M = 0$.*
- (iii) *M is either torsion or torsion-free.*

Proof. (i) First let $O \neq M$ be a vector space which is a weak multiplication module. If $\text{rank } M > 1$, then let $O \neq W \subset M$. According to Proposition 2.1, W is a prime submodule of M , and since M is a weak multiplication module, $W = IM$ where I is an ideal of the field R . So $I = O$ or $I = R$, which is a contradiction. Hence $\text{rank } M \leq 1$, and since $O \neq M$, then $\text{rank } M = 1$.

Now in the general case, if M is a non-zero torsion-free R -module, then $KM \neq O$, where K is the quotient field of R . By Lemma 2.3, KM is a weak multiplication K -module (vector space), and as we have proved above, $\text{rank}_K KM = 1$. Hence $\text{rank } M = \text{rank}_K KM = 1$.

(ii) Suppose that M is a torsion module, then $KM = O$ and therefore $\text{rank } M = \text{rank}_K KM = 0$.

(iii) If $T(M) \neq M$, we show that $T(M) = O$. If $T(M) \neq O$, then $KM \neq 0$ and by Lemma 2.3, KM is a non-zero weak multiplication K -module, so by part (i), $\text{rank}_K KM = 1$, that is $\text{rank } M = \text{rank}_K KM = 1$. It is easy to see that $T(M)$ is a prime submodule of M , so $T(M) = (T(M) : M)T(M)$ and since $T(M) \neq O$, $(T(M) : M) \neq O$. Let $0 \neq r \in (T(M) : M)$. Since $\text{rank } M = 1$, let $\{x\}$ be a linearly independent set in M . Now, $rx \in rM \subseteq T(M)$, so there exists $0 \neq r_1 \in R$ such that $r_1 rx = 0$, and this is a contradiction, because $\{x\}$ is linearly independent. □

Proposition 2.5. *A finitely generated module is a multiplication module if and only if it is locally cyclic.*

Proof. See [3, Proposition 5]. □

Theorem 2.6. *Let R be a local ring with a maximal ideal m and let M be a finitely generated R -module. If $\{\bar{u}_1, \bar{u}_2, \bar{u}_3, \dots, \bar{u}_n\}$ is a basis of the vector space $\bar{M} = \frac{M}{mM}$ over the field $\frac{R}{m}$, then $\{u_1, u_2, u_3, \dots, u_n\}$ is a minimal basis of M .*

Proof. See [7, Theorem 2.3]. □

Theorem 2.7. *Every finitely generated weak multiplication module is a multiplication module.*

Proof. Suppose that M is a finitely generated weak multiplication R -module. We show that M is locally cyclic, and by Proposition 2.5, M is a multiplication module. By localization and Lemma 2.3, we can assume that M is a finitely generated weak multiplication R -module where R is a local ring. Let m be the only maximal ideal of R . Obviously $\frac{M}{mM}$ is a finitely generated weak multiplication $\frac{R}{m}$ -module. If $mM = M$, then by Nakayama's Lemma $M = O$, so it is cyclic.

If $mM \neq M$, then $\text{rank}_{R/m} \frac{M}{mM} = 1$, by Proposition 2.4 (i) and by Theorem 2.6, M is a cyclic R -module. □

Theorem 2.8. *If R is a ring, then the following are equivalent.*

- (i) $\dim R = 0$.
- (ii) *For every weak multiplication R -module M , if $T(M) = 0$, then M is cyclic.*
- (iii) *For every weak multiplication R -module M , if $T(M) = 0$, then M is a multiplication module.*

Proof. (i) \Rightarrow (ii). First let R be a field. Let M be a torsion-free weak multiplication R -module. If $M = 0$, then M is cyclic. So let $0 \neq M$. M is a non-zero weak multiplication vector space over the field R . According to Proposition 2.4 (i), we have $\text{rank } M = 1$. That is $M \cong R$, and evidently M is cyclic.

Now we prove the general case. Let $0 \neq M$. It is easy to see that $T(M) = 0$ is a prime submodule of M . Hence $(T(M) : M)$ is a prime ideal of R and since $\dim R = 0$, $\frac{R}{(T(M):M)}$ is a field. Since $T(M) = 0$, one can easily show that $M \cong \frac{M}{0} = \frac{M}{T(M)}$ is a torsion-free weak multiplication $\frac{R}{(T(M):M)}$ -module. So M is a torsion-free weak multiplication module over the field $\frac{R}{(T(M):M)}$. And as we have proved above M is a cyclic $\frac{R}{(T(M):M)}$ -module and clearly M is a cyclic R -module.

(ii) \Rightarrow (iii). Is obvious.

(iii) \Rightarrow (i). Let P be a prime ideal of R . It is enough to prove that $\frac{R}{P}$ is a field.

If K is the quotient field of the integral domain $\frac{R}{P}$, then by Theorem 1 in [5], $\text{Spec}_{\frac{R}{P}}(K) = \{O\}$. So K is a torsion-free weak multiplication $\frac{R}{P}$ -module. Therefore by assumption it is a multiplication module. And since $\frac{R}{P} \leq K$, we have $\frac{R}{P} = IK$, where I is a non-zero ideal of $\frac{R}{P}$ and obviously $IK = K$. Hence $\frac{R}{P} = K$, and this completes the proof. □

Corollary 2.9. *If R is an integral domain, then the following are equivalent.*

- (i) R is a field.
- (ii) Every weak multiplication R -module is cyclic.
- (iii) Every weak multiplication R -module is a multiplication module.

Proof. If R is a field, then since every weak multiplication R -module is a vector space, it is a torsion-free weak multiplication R -module, so the proof follows by Theorem 2.8. □

Lemma 2.10. *Let R be a ring and M an R -module whose annihilator is contained in only finitely many maximal ideals m_1, m_2, \dots, m_n of R . If M_{m_i} is a cyclic R_{m_i} -module for $1, 2, \dots, n$, then M is a cyclic R -module.*

Proof. See Lemma 3 of [3]. □

In [3, Proposition 8], Barnard proved:

Every finitely generated Artinian multiplication R -module M is cyclic. In this case we know that $\frac{R}{\text{Ann } M}$ is an Artinian ring and obviously M is a multiplication $\frac{R}{\text{Ann } M}$ -module. So the following result is a generalization of this result.

Proposition 2.11. *Every weak multiplication module over an Artinian ring is cyclic.*

Proof. Let M' be a weak multiplication module over an Artinian ring R' . We prove that M' is locally cyclic and by Lemma 2.10, M' is cyclic. Let P be a prime ideal. Put $M'_P = M$ and $R'_P = R$. So R is a local Artinian ring and by Lemma 2.3, M is a weak multiplication R -module. Suppose that P is the only prime ideal of R , then $P^n = O$ for some natural number n . If $PM = M$, obviously $O = P^n M = M$, so let $PM \neq M$. $\frac{M}{PM}$ is a weak multiplication $\frac{R}{P}$ -module. Therefore, by Proposition 2.4 (i), we have $\text{rank}_{\frac{R}{P}} \frac{M}{PM} = 1$. That means PM is a maximal submodule of M . If $x \in M - PM$, then $PM \subset PM + Rx \subseteq M$, and therefore $PM + Rx = M$. Thus $O = P^n \frac{M}{Rx} = P \frac{M}{Rx} = \frac{M}{Rx}$, so $M = Rx$. □

Proposition 2.12. *If m is a maximal ideal of the ring R which is a minimal prime ideal and $m \neq m^2$, then the following are equivalent.*

- (i) m is a weak multiplication R -module.
- (ii) There is no ideal between m^2 and m .
- (iii) $\text{Spec}_R m = \{m^2\}$.

Proof. By localization and Lemma 2.3 we can assume that R is a local ring with the only prime ideal m .

(i) \Rightarrow (ii). Let m be a weak multiplication R -module. If $m^2 \subseteq I \subset m$ where I is an ideal of R , we show that I is a prime submodule of m . Let $r_1 r_2 \in I$, where $r_1 \in R$ and $r_2 \in m$. Suppose that $r_2 \notin I$, then r_1 is not a unit, hence $r_1 \in m$, hence $r_1 m \subseteq m^2 \subset I$, that is I is a prime submodule of m .

Since m is a weak multiplication module, and I is a prime submodule, then $I = mm_1$ for some ideal m_1 of R . If $m_1 = R$, then $I = mm_1 = m$, which is impossible. So $m_1 \subseteq m$, that is $m^2 \subseteq I = mm_1 \subseteq m^2$, thus there is no ideal between m^2 and m .

(ii) \Rightarrow (iii). Suppose that there is no ideal between m^2 and m . If I is a prime submodule of the R -module m , then $(I : m)$ is a prime ideal. Further, since m is the only prime ideal of R , we have $(I : m) = m$. Therefore $m^2 \subseteq I \subset m$, and by assumption $I = m^2$, hence $\text{Spec}_R m = \{m^2\}$.

(iii) \Rightarrow (i) Is clear.

The following theorem is a known result, but we will also prove it by the above result. \square

Corollary 2.13. *If R is a local Artinian ring and m is a maximal ideal of R , then m is cyclic if and only if $\text{rank}_{\frac{R}{m}} \frac{m}{m^2} \leq 1$.*

Proof. \Rightarrow Is obvious.

\Leftarrow If $\text{rank}_{\frac{R}{m}} \frac{m}{m^2} = 0$, then $m^2 = m$, and by Nakayama's lemma we have $m = 0$. If $\text{rank}_{\frac{R}{m}} \frac{m}{m^2} = 1$, then there is no ideal between m^2 and m , so by Proposition 2.12, m is a weak multiplication R -module and the proof follows by Proposition 2.11. \square

Acknowledgments. The author would like to thank Professor L. Grunenfelder of Dalhousie University in Canada for his helpful advice and Professor H. Sharif of Shiraz University in Iran for his encouragement.

References

- [1] *S. Abu-Saemeh*: On dimension of finitely generated modules. *Comm. Algebra* 23 (1995), 1131–1144.
- [2] *A. Azizi and H. Sharif*: On prime submodules. *Honam Math. J.* 21 (1999), 1–12.
- [3] *A. Barnard*: Multiplication modules. *J. Algebra* 71 (1981), 174–178.
- [4] *C. P. Lu*: Prime submodules of modules. *Comment. Math. Univ. St. Paul.* 33 (1984), 61–69.
- [5] *C. P. Lu*: Spectra of modules. *Comm. Algebra* 23 (1995), 3741–3752.
- [6] *R. L. McCasland and M. E. Moore*: Prime submodules. *Comm. Algebra* 20 (1992), 1803–1817.
- [7] *H. Matsumura*: *Commutative Ring Theory*. Cambridge University Press, Cambridge, 1990.

Author's address: Department of Mathematics, College of Sciences, Shiraz University, Shiraz 71454, Iran, e-mail: razizi@susc.ac.ir.