

Mehmet Bektaş

Totally real submanifolds in a quaternion space form

Czechoslovak Mathematical Journal, Vol. 54 (2004), No. 2, 341–346

Persistent URL: <http://dml.cz/dmlcz/127892>

Terms of use:

© Institute of Mathematics AS CR, 2004

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

TOTALLY REAL SUBMANIFOLDS
IN A QUATERNION SPACE FORM

MEHMET BEKTAŞ, Elaziğ

(Received June 22, 2001)

Abstract. In this paper, we prove a theorem for n -dimensional totally real minimal submanifold immersed in quaternion space form.

Keywords: totally real submanifold, quaternion space form

MSC 2000: 53C40, 53C56

1. INTRODUCTION

Let $M(c)$ denote a $4n$ -dimensional quaternion space form of quaternion sectional curvature c and let $P(H)$ denote the $4n$ -dimensional quaternion projective space of constant quaternion sectional curvature 4. Let N be an n -dimensional Riemannian manifold isometrically immersed in $M(c)$. We call N a totally real submanifold of $M(c)$ if each tangent 2-plane of N is mapped into a totally real plane in $M(c)$. Chen and Houh [3] gave an inequality for totally real submanifolds of $M(c)$. In particular they considered the case when N was totally geodesic. Later, Sun [6] considered the case when N was pseudo-umbilical and extended the theorem of Chen and Houh; he obtained two integral inequalities for compact totally real submanifolds of $M(c)$. There is other literature studying totally real submanifolds [4], [8], [9].

In this paper, we make use of Yau's maximum principle to study the complete totally real minimal submanifolds with Ricci curvature bounded from below and use the method of proof which is given in [7]. Thus we obtain the following result.

Main theorem. *Let $M(c)$ denote a $4n$ -dimensional quaternion space form of quaternion sectional curvature c . Let N be an n -dimensional totally real minimal manifold immersed in $M(c)$ with Ricci curvature bounded from below. Then*

either N is totally geodesic or

$$(1.1) \quad \inf \tau \leq \frac{1}{12}(3n-2)(n+1)c$$

where τ is the scalar curvature of N .

2. LOCAL FORMULAS

We use the same notations and terminology as in [3], [6] unless otherwise stated. Let $M(c)$ denote a $4n$ -dimensional quaternion space form of quaternion sectional curvature c and let N be an n -dimensional totally real submanifold of $M(c)$. We choose a local field of orthonormal frames,

$$\begin{aligned} e_1, \dots, e_n, \quad e_{I(1)} = Ie_1, \dots, e_{I(n)} = Ie_n, \quad e_{J(1)} = Je_1, \dots, e_{J(n)} = Je_n, \\ e_{K(1)} = Ke_1, \dots, e_{K(n)} = Ke_n, \end{aligned}$$

is such a way that when restricted to N , e_1, \dots, e_n are tangent to N . Here I, J, K are the almost Hermitian structure and satisfy

$$IJ = -JI = K, \quad JK = -KJ = I, \quad KI = -IK = J, \quad I^2 = J^2 = K^2 = -1.$$

We shall use the following convention on the range of indices:

$$\begin{aligned} A, B, \dots = 1, \dots, n, \quad I(1), \dots, I(n), \quad J(1), \dots, J(n), \quad K(1), \dots, K(n), \\ \alpha, \beta, \dots = I(1), \dots, I(n), \quad J(1), \dots, J(n), \quad K(1), \dots, K(n), \\ i, j, \dots = 1, \dots, n, \quad \Phi = I, J, K. \end{aligned}$$

Let $\{w_A\}$ be the dual frame field. Then the structure equations of $M(c)$ are given by

$$\begin{aligned} dw_A &= - \sum_B w_{AB} \wedge w_B, \quad w_{AB} + w_{BA} = 0, \\ (2.1) \quad dw_{AB} &= - \sum_C w_{AC} \wedge w_{CB} + \frac{1}{2} \sum_{CD} K_{ABCD} w_C \wedge w_D, \\ K_{ABCD} &= \frac{c}{4} (\delta_{AC} \delta_{BD} - \delta_{AD} \delta_{BC} + I_{AC} I_{BD} - I_{AD} I_{BC} + 2I_{AB} I_{CD} + J_{AC} J_{BD} \\ &\quad - J_{AD} J_{BC} + 2J_{AB} J_{CD} + K_{AC} K_{BD} - K_{AD} K_{BC} + 2K_{AB} K_{CD}). \end{aligned}$$

Restricting these forms to N , we get the following structure equations of the immersion:

$$\begin{aligned}
 (2.2) \quad & w_\alpha = 0, \quad w_{\alpha i} = \sum_j h_{ij}^\alpha w_j, \quad h_{ij}^\alpha = h_{ji}^\alpha, \quad h_{jk}^{\Phi(i)} = h_{ik}^{\Phi(j)} = h_{ij}^{\Phi(k)}, \\
 & dw_{ij} = - \sum_k w_{ik} \wedge w_{kj} + \frac{1}{2} \sum_k R_{ijkl} w_k \wedge w_l, \\
 & R_{ijkl} = K_{ijkl} + \sum_\alpha (h_{ik}^\alpha h_{jl}^\alpha - h_{il}^\alpha h_{jk}^\alpha), \\
 (2.3) \quad & dw_{\alpha\beta} = - \sum_\gamma w_{\alpha\gamma} \wedge w_{\gamma\beta} + \frac{1}{2} \sum_{ij} R_{\alpha\beta ij} w_i \wedge w_j, \\
 & R_{\alpha\beta ij} = K_{\alpha\beta ij} + \sum_k (h_{ik}^\alpha h_{kj}^\beta - h_{ik}^\beta h_{kj}^\alpha).
 \end{aligned}$$

We call $h = \sum_{ij\alpha} h_{ij}^\alpha w_i w_j e_\alpha$ the second fundamental form of the immersed manifold N . We denote by $S = \sum_{ij\alpha} (h_{ij}^\alpha)^2$ the square of the length of h .

If N is minimal in $M(c)$, i.e., *trace* $h = 0$, then by the equations (2.1) and (2.2), we have

$$(2.4) \quad \tau = \frac{c}{4}n(n+1) - S$$

where S is the scalar curvature of N . We define h_{ij}^α and h_{ijkl}^α by

$$(2.5) \quad \sum_k h_{ijk}^\alpha w_k = dh_{ij}^\alpha - \sum_l h_{il}^\alpha w_l j - \sum_l h_{lj}^\alpha w_l i + \sum_\beta h_{ij}^\beta w_{\alpha\beta}$$

and

$$\sum_l h_{ijkl}^\alpha w_l = dh_{ijk}^\alpha - \sum_l h_{ljk}^\alpha w_l i - \sum_l h_{ilk}^\alpha w_l j - \sum_l h_{ijl}^\alpha w_l k + \sum_\beta h_{ijk}^\beta w_{\alpha\beta},$$

respectively, where

$$h_{ijk}^\alpha = h_{ikj}^\alpha$$

and

$$h_{ijkl}^\alpha - h_{ijlk}^\alpha = \sum_m h_{im}^\alpha R_{mjkl} + \sum_m h_{mj}^\alpha R_{mikl} - \sum_\beta h_{ij}^\beta R_{\alpha\beta kl}.$$

Let H_α and Δ denote the $n \times n$ matrix (h_{ij}^α) and the Laplacian on N , respectively. By a simple calculation, we have (cf. [1], [2], [5])

$$(2.6) \quad \frac{1}{2}\Delta S = \sum_{ij\alpha} (h_{ijk}^\alpha)^2 + \frac{c}{4}(n+1)S + \sum_{\alpha\beta} \text{tr}(H_\alpha H_\beta - H_\beta H_\alpha)^2 - \sum_{\alpha\beta} (\text{tr } H_\alpha H_\beta)^2.$$

In order to prove the main theorem, we need the following Lemmas.

Lemma 1 [6]. Let $H_i, i \geq 2$ be symmetric $n \times n$ matrices, $S_i = \text{tr } H_i^2, S = \sum_i S_i$.

Then

$$(2.7) \quad \sum_{ij} \text{tr}(H_i H_j - H_j H_i)^2 - \sum_{ij} (\text{tr } H_i H_j)^2 \geq -\frac{3}{2} S^2$$

and the equality holds if and only if either all $H_i = 0$ or there exist two H_i different from zero. Moreover, if $H_1 \neq 0, H_2 \neq 0, H_i = 0, i \neq 1, 2$, then $S_1 = S_2$ and there exists an orthogonal $n \times n$ matrix T such that

$$T H_1^t T = \begin{pmatrix} \sqrt{\frac{S_1}{2}} & 0 & \dots & 0 \\ 0 & -\sqrt{\frac{S_1}{2}} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}, \quad T H_2^t T = \begin{pmatrix} 0 & \sqrt{\frac{S_1}{2}} & \dots & 0 \\ \sqrt{\frac{S_1}{2}} & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}.$$

Lemma 2 [7]. Let N be a complete Riemannian manifold with Ricci curvature bounded from below and let f be a C^2 -function bounded from above on N , then for all $\varepsilon > 0$, there exists a point $x \in N$ at which

- (i) $\sup f - \varepsilon < f(x)$,
- (ii) $\|\nabla f(x)\| < \varepsilon$,
- (iii) $\Delta f(x) < \varepsilon$.

3. PROOF OF THE MAIN THEOREM

In this section, the method of the proof used by Ximin in [7] is applied to a totally real minimal submanifold immersed in quaternion space form.

From (2.6) and (2.7), we obtain

$$(3.1) \quad \frac{1}{2} \Delta S \geq S \left(\frac{c}{4}(n+1) - \frac{3}{2} S \right).$$

We know that $S = \frac{c}{4}n(n+1) - \tau$. By the condition of the theorem, we conclude that S is bounded. We define $f = S$ and $F = (f+a)^{\frac{1}{2}}$ (where $a > 0$ is any positive constant number). F is bounded. We have

$$(3.2) \quad dF = \frac{1}{2}(f+a)^{-\frac{1}{2}} df,$$

$$(3.3) \quad \begin{aligned} \Delta F &= \frac{1}{2} \left(-\frac{1}{2}(f+a)^{-\frac{3}{2}} \|df\|^2 + (f+a)^{-\frac{1}{2}} \Delta f \right) \\ &= \frac{1}{2} (-2\|dF\|^2 + \Delta f)(f+a)^{-\frac{1}{2}}, \end{aligned}$$

i.e.,

$$(3.4) \quad \Delta F = \frac{1}{2F}(-2\|dF\|^2 + \Delta f).$$

Hence, $F\Delta F = -\|dF\|^2 + \frac{1}{2}\Delta f$ or $\frac{1}{2}\Delta f = F\Delta F + \|dF\|^2$. Applying Lemma 2 to F , we have for all $\varepsilon > 0$, there exists a point $x \in N$ such that at x

$$(3.5) \quad \|dF(x)\| \leq \varepsilon,$$

$$(3.6) \quad \Delta F(x) < \varepsilon,$$

$$(3.7) \quad F(x) > \sup F - \varepsilon.$$

From (3.5), (3.6) and (3.7), we have

$$(3.8) \quad \frac{1}{2}\Delta f < \varepsilon^2 + F\varepsilon = \varepsilon(\varepsilon + F).$$

We take a sequence $\{\varepsilon_m\}$ such that $\varepsilon_m \rightarrow 0$ ($m \rightarrow \infty$) and for all m , there exists a point $x_m \in N$ such that (3.5), (3.6) and (3.7) hold. Therefore, $\varepsilon_m(\varepsilon_m + F(x_m)) \rightarrow 0$ ($m \rightarrow \infty$) (because F is bounded).

From (3.7), we have $F(x_m) > \sup F - \varepsilon_m$, because $\{F(x_m)\}$ is a bounded sequence. So we get $F(x_m) \rightarrow F_0$ (if necessary, we can choose a subsequence). Hence, $F_0 \geq \sup F$. So we have

$$(3.9) \quad F_0 = \sup F.$$

From the definition of F , we get

$$(3.10) \quad f(x_m) \rightarrow f = \sup f.$$

(3.1) and (3.8) imply that

$$(3.11) \quad f\left(\frac{c}{4}(n+1) - \frac{3}{2}S\right) \leq \frac{1}{2}\Delta f \leq \varepsilon(\varepsilon + F),$$

and

$$(3.12) \quad f(x_m)\left(\frac{c}{4}(n+1) - \frac{3}{2}f(x_m)\right) < \varepsilon_m^2 + \varepsilon_m F(x_m) \leq \varepsilon_m^2 + \varepsilon_m F_0.$$

Let $m \rightarrow \infty$, then $\varepsilon_m \rightarrow 0$ and $f(x_m) \rightarrow f_0$. Hence,

$$(3.13) \quad f_0\left(\frac{c}{4}(n+1) - \frac{3}{2}f_0\right) \leq 0.$$

- (i) If $f_0 = 0$, we have $f = S \equiv 0$. Hence, M is totally geodesic.
(ii) If $f_0 > 0$, we have $\frac{c}{4}(n+1) - \frac{3}{2}f_0 \leq 0$ and

$$f_0 \geq \frac{c}{6}(n+1),$$

that is, $\sup S \geq \frac{c}{6}(n+1)$. Therefore, $\inf \tau \leq \frac{1}{12}(3n-2)(n+1)c$. This completes the proof.

Acknowledgement. The author wishes to express his sincere thanks to the referee for the careful reading and very helpful comments on the earlier versions of this manuscript.

References

- [1] *M. Bektaş and M. Ergüt:* Compact space-like submanifolds with parallel mean curvature vector of a Pseudo-Riemannian space. *Acta Univ. Palack. Olomuc. Fac. Rerum Natur. Math.* 38 (1999), 17–24.
- [2] *B. Y. Chen and K. Ogiue:* On totally real submanifolds. *Trans. Amer. Math. Soc.* 193 (1974), 257–266.
- [3] *B. Y. Chen and C. S. Houh:* Totally real submanifolds of a quaternion projective space. *Ann. Math. Pura Appl.* 120 (1979), 185–199.
- [4] *Y. B. Shen:* Totally real minimal submanifolds in a quaternion projective space. *Chinese Ann. Math. Ser. B* 14 (1993), 297–306.
- [5] *S. S. Chern, M. Carmo and S. Kobayashi:* Minimal submanifolds of a sphere with second fundamental form of constant length, functional analysis and related fields. *Proc. Conf. for M. Stone, Univ. Chicago, 1968. Vol III. Springer-Verlag, New York, 1970, pp. 59–75.*
- [6] *H. Sun:* Totally real pseudo-umbilical submanifolds of a quaternion space form. *Glasgow Math. J.* 40 (1998), 109–115.
- [7] *L. Ximin:* Totally real submanifolds in a complex projective space. *Internat. J. Math. Math. Sci.* 22 (1999), 205–208.
- [8] *L. Ximin:* Totally real submanifolds in $HP^m(1)$ with isotropic second fundamental form. *Saitama Math. J.* 16 (1998), 23–29.
- [9] *L. Ximin:* Totally real minimal submanifolds in a quaternion projective space. *Proc. Japan. Acad. Ser. A-Math. Sci.* 72 (1996), 238–239.

Author's address: Department of Mathematics, Firat University, 23119 Elazığ, Türkiye,
e-mail: mbektas@firat.edu.tr.