

Hishyar Kh. Abdullah

A note on the oscillation of second order differential equations

*Czechoslovak Mathematical Journal*, Vol. 54 (2004), No. 4, 949–954

Persistent URL: <http://dml.cz/dmlcz/127943>

## Terms of use:

© Institute of Mathematics AS CR, 2004

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON THE OSCILLATION OF SECOND ORDER  
DIFFERENTIAL EQUATIONS

HISHYAR KH. ABDULLAH, Sharjah

(Received January 15, 2002)

*Abstract.* We give a sufficient condition for the oscillation of linear homogeneous second order differential equation  $y'' + p(x)y' + q(x)y = 0$ , where  $p(x), q(x) \in C[\alpha, \infty)$  and  $\alpha$  is positive real number.

*Keywords:* oscillatory, second order differential equations

*MSC 2000:* 34A30, 34C10

1. INTRODUCTION

In this paper a sufficient condition is given for the oscillation of linear second order differential equation with variable coefficients

$$(1.1) \quad y'' + p(x)y' + q(x)y = 0,$$

where  $p(x)$  and  $q(x)$  are continuous functions on the interval  $[\alpha, \infty)$ , where  $\alpha$  is real number. Here we are concerned with sufficient conditions on the coefficient  $p(x)$  so that the equation (1.1) is oscillatory.

The study of oscillation of second order differential equations is of great interest. Many criteria have been found which involve the behavior of the integral of a combination of the coefficients. This approach has been motivated by some authors (for example see [1], [5], [6], [7], [9]).

**Definition 1.** A solution  $y(x)$  of the differential equation (1.1) is said to be nontrivial if  $y(x) \neq 0$  for at least one  $x \in [\alpha, \infty)$ .

**Definition 2.** A nontrivial solution  $y(x)$  of (1.1) is said to be oscillatory if it has arbitrarily large zeros on  $[\alpha, \infty)$ , otherwise it is said to be “nonoscillatory”.

## 2. MAIN RESULTS

We prove the following theorems.

**Theorem 1.** *If  $p(x) < 0$  on  $[\alpha, \infty)$  is such that*

$$(2.1) \quad \lim_{x \rightarrow \infty} \left[ \frac{1}{4} \int_{\alpha}^x (4q(s) - (p(s))^2) ds \right] = \infty,$$

*then any solution of the differential equation (1.1) is oscillatory on  $[\alpha, \infty)$ .*

**Proof.** Suppose that the differential equation (1.1) is non oscillatory, then there exists a non trivial solution of (1.1) that has no zeros on  $[\beta, \infty)$  for  $\beta > \alpha$ .

Let  $w(x)$  be the function defined by  $w(x) = -y^{-1}(x)y'(x)$  for  $x \in [\beta, \infty)$ .

Then  $w(x)$  is well defined and satisfies the Riccati equation

$$w'(x) = w^2(x) - p(x)w(x) + q(x), \quad \text{on } [\beta, \infty).$$

Integrating both sides of this equation from  $\beta$  to  $x$  we get

$$\begin{aligned} w(x) &= w(\beta) + \int_{\beta}^x [(w(s))^2 - p(s)w(s) + q(s)] ds \\ &= w(\beta) + \int_{\beta}^x \left( w(s) - \frac{p(s)}{2} \right)^2 ds + \frac{1}{4} \int_{\beta}^x (4q(s) - (p(s))^2) ds. \end{aligned}$$

Now the hypothesis (2.1) implies that there exists  $\gamma > \beta$  such that

$$w(x) > \int_{\gamma}^x \left( w(s) - \frac{p(s)}{2} \right)^2 ds \quad \text{on } [\gamma, \infty).$$

Define

$$(2.2) \quad G(x) = \int_{\gamma}^x \left( w(s) - \frac{p(s)}{2} \right)^2 ds,$$

then  $w(x) > G(x) > 0$  on  $[\gamma, \infty)$ .

Differentiating (2.2) we get

$$G'(x) = \left( w(x) - \frac{p(x)}{2} \right)^2 > w^2(x),$$

since  $p(x) < 0$ , therefore  $G'(x) > G^2(x)$ , thus

$$1 < \frac{G'(x)}{G^2(x)},$$

this inequality holds for  $x > \gamma$ .

Integrating both sides of this inequality from  $\gamma$  to  $x$  we get

$$\int_{\gamma}^x ds < \frac{1}{G(\gamma)} - \frac{1}{G(x)}.$$

Therefore since  $G(x) > 0$ , we conclude that

$$\lim_{x \rightarrow \infty} \int_{\gamma}^x ds < \frac{1}{G(\gamma)}.$$

But this is not true. Thus the differential equation (1.1) is oscillatory, and this completes the proof.  $\square$

**Theorem 2.** *If  $p(x) < 0$  on  $[\alpha, \infty)$  and there exists a non vanishing function  $g(x) \in C^1[\alpha, \infty)$ ;  $g(x) > 0$ , such that*

$$(2.3) \quad \lim_{x \rightarrow \infty} \int_{\alpha}^x (g(s))^{-1} ds = \infty$$

and

$$(2.4) \quad \lim_{x \rightarrow \infty} \left[ -\frac{1}{4} \int_{\alpha}^x \left\{ p(s)^2 g(s) + \frac{((g(s))')^2}{g(s)} - 2p(s)(g(s))' - 4g(s)q(s) \right\} ds + \frac{1}{2}(g(x))' \right] = \infty,$$

then any solution of the differential equation (1.1) is oscillatory.

**P r o o f.** Suppose that there exists a nonoscillatory solution  $y(x)$  of the differential equation (1.1), i.e.  $y(x)$  has no zeros on  $[\alpha, \infty)$ .

Define for  $x > \alpha$

$$(2.5) \quad w(x) = -g(x)y'(x)y^{-1}(x),$$

where  $g(x)$  is a non vanishing function belonging to  $C^1[\alpha, \infty)$  and  $1/g(x) > 0$  on  $[\alpha, \infty)$ .

Differentiating (2.5) with respect to  $x$ , on the interval  $[\alpha, \infty)$ ,  $w(x)$  satisfies the Riccati equation we see that.

$$w'(x) = \frac{1}{g(x)} [w^2(x) - g(x)p(x)w(x) + g'(x)w(x)] + g(x)q(x).$$

Now for  $x \in [\alpha, \infty)$ , defining

$$(2.6) \quad H(x) = w(x) + \frac{1}{2}g'(x),$$

we will have

$$\begin{aligned} w'(x) &= \frac{1}{g(x)} \left[ \left( H(x) - \frac{p(x)g(x)}{2} \right)^2 - \left( \frac{p(x)g(x)}{2} \right)^2 - \frac{1}{4}(g'(x))^2 \right. \\ &\quad \left. + \frac{1}{2}p(x)g(x)g'(x) \right] + g(x)q(x) \\ &= \frac{1}{g(x)} \left( H(x) - \frac{p(x)g(x)}{2} \right)^2 - \frac{p^2(x)g(x)}{4} - \frac{1}{4} \frac{(g'(x))^2}{g(x)} + \frac{1}{2}p(x)g'(x) \\ &= \frac{1}{g(x)} \left( H(x) - \frac{p(x)g(x)}{2} \right)^2 \\ &\quad - \frac{1}{4} \left[ p^2(x)g(x) + \frac{(g'(x))^2}{g(x)} - 2p(x)g'(x) - 4g(x)q(x) \right]. \end{aligned}$$

Integrating both sides of the above equation from  $\alpha$  to  $x$  we get

$$\begin{aligned} w(x) &= w(\alpha) + \int_{\alpha}^x g^{-1}(s) \left( H(s) - \frac{1}{2}p(s)g(s) \right)^2 ds \\ &\quad - \frac{1}{4} \int_{\alpha}^x [p^2(s)g(s) + g^{-1}(s)(g'(s))^2 - 2p(s)g'(s) - 4g(s)q(s)] ds. \end{aligned}$$

Then by using (2.6) we have

$$\begin{aligned} H(x) &= w(\alpha) + \int_{\alpha}^x g^{-1}(s) \left( H(s) - \frac{1}{2}p(s)g(s) \right)^2 ds \\ &\quad - \frac{1}{4} \int_{\alpha}^x [p^2(s)g(s) + g^{-1}(s)(g'(s))^2 - 2p(s)g'(s) - 4g(s)q(s)] ds + \frac{1}{2}g'(x). \end{aligned}$$

The hypothesis of Theorem 2 implies that there exists  $\beta > \alpha$  such that

$$H(x) > \int_{\beta}^x g^{-1}(s) \left( H(s) - \frac{1}{2}p(s)g(s) \right)^2 ds$$

holds for  $x > \beta$ . Define a function  $Q(x)$  for  $x > \beta$  by

$$(2.7) \quad Q(x) = \int_{\beta}^x g^{-1}(s) \left( H(s) - \frac{1}{2}p(s)g(s) \right)^2 ds.$$

Since  $p(x) < 0$ ,  $H(x) > Q(x) > 0$ . Differentiating (2.7) we have

$$\begin{aligned} Q'(x) &= g^{-1}(x) \left( H(x) - \frac{1}{2} p(x) g(x) \right)^2 \\ &\geq g^{-1}(x) \left( Q(x) - \frac{1}{2} p(x) g(x) \right)^2 \\ &> g^{-1}(x) Q^2(x), \end{aligned}$$

therefore

$$g^{-1}(x) < \frac{Q'(x)}{Q^2(x)}.$$

Integrating both sides of this inequality with respect to  $x$  (with  $x$  replaced by  $s$ ) from  $\beta$  to  $x$  we get for  $x > \beta$

$$\int_{\beta}^x g^{-1}(s) \, ds < \frac{1}{Q(\beta)} - \frac{1}{Q(x)};$$

since  $Q(x) > 0$ , therefore

$$\int_{\beta}^x g^{-1}(s) \, ds < \frac{1}{Q(\beta)}$$

which contradicts the hypothesis of the theorem. Hence the differential equation (1.1) is oscillatory.  $\square$

### 3. EXAMPLE

Following is an illustrative example showing the applicability of both theorems. Consider the second order differential equation

$$y''(x) - \frac{2}{x} y'(x) + \left( 1 + \frac{2}{x^2} \right) y(x) = 0, \quad \text{for } x > 0.$$

For this differential equation we have

$$p(x) = \frac{-2}{x} < 0, \quad \text{and} \quad q(x) = \left( 1 + \frac{2}{x^2} \right).$$

To show the applicability of Theorem 1, the hypothesis is satisfied as follows

$$\begin{aligned} \lim_{x \rightarrow \infty} \int_{\alpha}^x \left( q(s) - \left( \frac{p(s)}{2} \right)^2 \right) ds &= \lim_{x \rightarrow \infty} \int_{\alpha}^x \left( 1 + \frac{2}{s^2} - \frac{1}{4} \frac{4}{s^2} \right) ds \\ &= \lim_{x \rightarrow \infty} \int_{\alpha}^x \left( 1 + \frac{1}{s^2} \right) ds = \lim_{x \rightarrow \infty} \left[ s - \frac{1}{s} \right]_{\alpha}^x = \infty. \end{aligned}$$

Therefore the theorem implies that the differential equation is oscillatory.

This fact is directly verified by noting that the solution of the differential equation is given by

$$y(x) = x(k_1 \cos x + k_2 \sin x).$$

To show the applicability of Theorem 2, choose  $g(x) = x$ . It is clear that the hypothesis (2.4) is satisfied hence

$$\begin{aligned} \lim_{x \rightarrow \infty} & \left[ -\frac{1}{4} \int_{\alpha}^x \left\{ (p(s))^2 g(s) + \frac{((g(s))')^2}{g(s)} - 2p(s)(g(s))' - 4g(s)q(s) \right\} ds + \frac{1}{2}(g(x))' \right] \\ &= \lim_{x \rightarrow \infty} \left\{ -\frac{1}{4} \left[ \int_{\alpha}^x \frac{4}{s} + \frac{4}{s} + \frac{1}{s} - 4s - \frac{8}{s} \right] ds + \frac{1}{2} \right\} \\ &= \lim_{x \rightarrow \infty} \left\{ \left[ \frac{s^2}{2} - \frac{1}{4} \ln s \right]_{\alpha}^x + \frac{1}{2} \right\} = \infty. \end{aligned}$$

Hence Theorem 2 is applicable.

### Acknowledgement.

I would like to extend my thanks to the University of Sharjah for its support.

### References

- [1] *A. Sauer*: A note of zero-sequences of solutions of  $f'' + Af = 0$ . Amer. Math. Soc. *125* (1997), 1143–1147.
- [2] *G. J. Butler, I. H. Erbe and A. B. Mingarelli*: Riccati techniques and variational principles in oscillation theory for linear systems. Trans. Amer. Math. Soc. *303* (1987), 263–282.
- [3] *H. Erbe, Qinghai Kong and Shigui Ruan*: Kamenev type theorems for 2nd order matrix differential systems. Proc. Amer. Math. Soc. *117* (1993), 957–962.
- [4] *E. Hille*: Non-oscillation theorems. Trans. Amer. Math. Soc. *64* (1948), 234–252.
- [5] *I. Kamenev*: Integral criterion for oscillation of linear differential equations of second order. Zametki *23* (1978), 136–138.
- [6] *J. W. Macki and J. S. W. Wong*: Oscillation theorems for linear second order differential equations. Proc. Amer. Math. Soc. *20* (1969), 67–72.
- [7] *A. B. Mingarelli*: On a conjecture for oscillation of second order ordinary differential systems. Proc. Amer. Math. Soc. *82* (1981), 592–598.
- [8] *Ch. G. Philos and I. K. Purnaras*: Oscillations in superlinear differential equations of second order. J. Math. Anal. Appl. *165* (1992), 1–11.
- [9] *D. Willett*: On the oscillatory behavior of the solution of second order linear differential equations. Ann. Polon. Math. *21* (1969), 175–194.
- [10] *J. Yan*: A note on an oscillation criterion for an equation with damped term. Proc. Amer. Math. Soc. *90* (1984), 277–280.

*Author's address*: Dept. of Basic Sciences, College of Arts & Science, University of Sharjah, P.O.Box 27272, Sharjah, U.A.E., e-mail: [hishyar@sharjah.ac.ae](mailto:hishyar@sharjah.ac.ae).