

Danica Jakubíková-Studenovská
On a representation of monounary algebras

Czechoslovak Mathematical Journal, Vol. 55 (2005), No. 1, 157–164

Persistent URL: <http://dml.cz/dmlcz/127966>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A REPRESENTATION OF MONOUNARY ALGEBRAS

DANICA JAKUBÍKOVÁ-STUDENOVSKÁ, Košice

(Received April 25, 2002)

Abstract. In this note we deal with a question concerning monounary algebras which is analogous to an open problem for partially ordered sets proposed by Duffus and Rival.

Keywords: monounary algebra, connectedness, retract, retract irreducibility, representation

MSC 2000: 08A60

1. INTRODUCTION

Duffus and Rival [1] studied a certain form of representability of partially ordered sets. The representation under consideration was defined by means of retracts. In [1] it was remarked that “the following important problem remains unsolved:

(*) Does every partially ordered set have a representation $\{P_i : i \in I\}$ such that P_i for each $i \in I$ is irreducible?”

(The detailed definitions of the notions of representation and irreducibility are recalled in Section 2 below.)

We remark that a monounary algebra can be viewed as a particular case of a quasi-ordered set. Namely, a monounary algebra is defined to be an algebraic structure $\mathcal{A} = (A, f)$, where A is a non-empty set and f is a unary operation on A . To each monounary algebra \mathcal{A} there corresponds a quasi-ordered set $Q = (A, \leq)$, where the relation \leq is defined as follows: if $a, b \in A$, then $a \leq b$, whenever $f^n(a) = b$ for some $n \in \mathbb{N} \cup \{0\}$. Conversely, the quasi-ordered set $Q = (A, \leq)$ uniquely defines a monounary algebra $\mathcal{A} = (A, f)$.

Retracts and retract irreducibility of monounary algebras were studied in the author’s papers [2]–[7]. Let \mathcal{U} be the class of all monounary algebras and let \mathcal{U}_c be

Support of Slovak VEGA Grant 1/7468/20 is acknowledged.

the class of all connected monounary algebras. For $\mathcal{A} \in \mathcal{U}$ let $R(\mathcal{A})$ be the system of all isomorphic copies of all retracts of \mathcal{A} .

In the present paper we deal with a question analogous to (*) concerning a representation of a monounary algebra \mathcal{A} in a class \mathcal{K} for the case when $\mathcal{K} \in \{\mathcal{U}, \mathcal{U}_c, R(\mathcal{A})\}$; let us denote this question by (**). We prove that the answer to (**) is “No”.

2. ON THE QUESTION (**)

We start by recalling some definition.

First we recall some definitions for partially ordered sets.

Let P be a partially ordered set and let $R(P)$ be the system of all partially ordered sets Q such that Q is isomorphic to some retract of P . We say that P is irreducible, if, whenever $P_i \in R(P)$ for $i \in I$ and $P \in R\left(\prod_{i \in I} P_i\right)$, then there is $j \in I$ such that $P \in R(P_j)$. If $P \in R\left(\prod_{i \in I} P_i\right)$ and $P_i \in R(P)$ for each $i \in I$, then the system $\{P_i : i \in I\}$ is called a representation of P . (Cf. Duffus and Rival [1].)

We will use these definitions of representation and of irreducibility for monounary algebras.

Let $\mathcal{A} = (A, f) \in \mathcal{U}$. A nonempty subset M of A is said to be a retract of \mathcal{A} if there is a mapping h of A onto M such that h is an endomorphism of \mathcal{A} and $h(x) = x$ for each $x \in M$. The mapping h is called a retraction endomorphism corresponding to the retract M . Further, let $R(\mathcal{A})$ be the system of all monounary algebras \mathcal{B} such that \mathcal{B} is isomorphic to (M, f) for some retract M of \mathcal{A} .

Let \mathcal{K} be a system of monounary algebras. In [4] there was introduced the following definition: an element \mathcal{A} of \mathcal{U} is said to be retract irreducible in \mathcal{K} , if, whenever $\mathcal{B}_i \in \mathcal{K}$ for $i \in I$ and $\mathcal{A} \in R\left(\prod_{i \in I} \mathcal{B}_i\right)$, then there is $j \in I$ such that $\mathcal{A} \in R(\mathcal{B}_j)$.

In [2] and [3] there were described all $\mathcal{A} \in \mathcal{U}_c$ which are retract irreducible in \mathcal{U}_c and in [4] all $\mathcal{A} \in \mathcal{U}_c$ which are retract irreducible in \mathcal{U} . Further, in [6] and [7] there were found all $\mathcal{A} \in \mathcal{U}_c$ such that \mathcal{A} is retract irreducible in $R(\mathcal{A})$ (they were denoted as irreducible in the sense of Duffus and Rival, or, shortly, DR-irreducible). All $\mathcal{A} \in \mathcal{U}$ which are retract irreducible in \mathcal{U} were described in [5].

Analogously as for partially ordered sets we define the following notion. Let $\mathcal{A} \in \mathcal{U}$, $\mathcal{K} \subseteq \mathcal{U}$. A system $\{\mathcal{B}_i : i \in I\} \subseteq \mathcal{K}$ will be called a representation of \mathcal{A} in \mathcal{K} , if $\mathcal{A} \in R\left(\prod_{i \in I} \mathcal{B}_i\right)$.

We will consider the following question for a class $\mathcal{K} \subseteq \mathcal{U}$: (**) Does every monounary algebra \mathcal{A} have a representation $\{\mathcal{B}_i: i \in I\}$ in \mathcal{K} such that \mathcal{B}_i for each $i \in I$ is retract irreducible in \mathcal{K} ?

The aim of this paper is to prove

Theorem. *There exists a connected monounary algebra \mathcal{A} such that if $\mathcal{K} \in \{\mathcal{U}, \mathcal{U}_c, R(\mathcal{A})\}$, then \mathcal{A} possesses no representation $\{\mathcal{B}_i: i \in I\}$ of \mathcal{A} in \mathcal{K} such that for each $i \in I$, \mathcal{B}_i is retract irreducible in \mathcal{K} .*

3. THE CLASS $\mathcal{K} = R(\mathcal{A})$

In the following notation suppose that distinct symbols mean distinct elements.

3.1. Notation. For $n \in \mathbb{N}$ let

$$A_n = \{j_n: j \in \{1, \dots, n\}\}.$$

Put

$$A = \mathbb{N} \cup \bigcup_{n \in \mathbb{N}} A_n.$$

Further let

$$f(n) = n + 1 \quad \text{for each } n \in \mathbb{N},$$

$$f(j_n) = \begin{cases} (j + 1)_n & \text{for each } n \in \mathbb{N}, j \in \{1, \dots, n - 1\}, \\ 1 & \text{for each } n \in \mathbb{N}, j = n. \end{cases}$$

Denote $\mathcal{A} = (A, f)$ (cf. Fig. 1).




Fig. 1

Then we obviously have

3.2. Lemma. \mathcal{A} is a connected monounary algebra.

3.3. Lemma. Let M be a retract of \mathcal{A} . Then $1 \in M$ and $\text{card}(f^{-1}(1) \cap M) > 2$.

Proof. It is obvious that if M is a retract of \mathcal{A} , then $\mathbb{N} \subseteq M$. Suppose that $\text{card}(f^{-1}(1) \cap M) \leq 2$. Then there is $n \in \mathbb{N}$ such that $k_k \notin M$ for each $k \in \mathbb{N}$, $k \geq n$. Hence $A_k \cap M = \emptyset$ for each $k \in \mathbb{N}$, $k \geq n$. Let h be a retraction endomorphism corresponding to M . Denote $z = h(1_n)$. Then

$$z \in M \subseteq \mathbb{N} \cup A_1 \cup \dots \cup A_{n-1},$$

thus

$$(1) \quad f^n(z) \in \mathbb{N} - \{1\}.$$

Further, $1 \in M$, hence

$$f^n(z) = f^n(h(1_n)) = h(f^n(1_n)) = h(1) = 1,$$

a contradiction to (1). □

3.4. Corollary. If M is a retract of \mathcal{A} , then (M, f) is not retract irreducible in the class $R(\mathcal{A})$.

Proof. According to [6, 2.9] and [7, 4.1], we obtain that if (M, f) is retract irreducible in $R(\mathcal{A})$, then $\text{card} f^{-1}(x) < 2$ for each $x \in M$. Hence 3.3 yields the required assertion. □

3.5. Proposition. Let $\{B_i: i \in I\}$ be a representation of \mathcal{A} in the class $R(\mathcal{A})$. Then B_i fails to be retract irreducible in $R(\mathcal{A})$ for each $i \in I$.

Proof. Let $i \in I$. Then there exists a retract M of \mathcal{A} such that

$$(1) \quad \mathcal{B}_i \cong (M, f).$$

By 3.4 and (1), B_i is not retract irreducible in the class $R(\mathcal{A})$. □

4. THE CLASS $\mathcal{K} = \mathcal{U}_c$

Let \mathcal{A} be as in Section 3 and suppose that the system $\{\mathcal{B}_i: i \in I\} \subseteq \mathcal{U}_c$ is a representation of \mathcal{A} such that if $i \in I$, then \mathcal{B}_i is retract irreducible in \mathcal{U}_c . Then [2, (R)] and [3, (R1)] imply

4.1. Lemma. *If $i \in I$, then one of the following conditions is satisfied:*

- (i) \mathcal{B}_i is a cycle with p^m elements, where p is a prime and $m \in \mathbb{N}$,
- (ii) $\mathcal{B}_i \cong (\mathbb{N}, f)$,
- (iii) \mathcal{B}_i contains a one element cycle $\{c\}$ and if $\{a, b\} \subseteq \mathcal{B}_i$ with $f(a) = f(b)$, then either $a=b$ or $c \in \{a, b\}$.

The system $\{\mathcal{B}_i : i \in I\} \subseteq \mathcal{U}_c$ is a representation of \mathcal{A} , thus there is a retract M of $\prod_{i \in I} \mathcal{B}_i$ such that $\mathcal{A} \cong (M, f)$. Let ν be an isomorphism of \mathcal{A} onto (M, f) .

4.2. Lemma. *If $i \in I$, then (ii) fails to hold.*

Proof. Let $i \in I$ and suppose that (ii) is valid. Then $\mathcal{B}_i = \{c_n : n \in \mathbb{N}\}$ and $f(c_n) = c_{n+1}$ for each $n \in \mathbb{N}$, where $c_k \neq c_l$ for each $k, l \in \mathbb{N}$, $k \neq l$. Denote $t = \nu(1)$ and, if $n \in \mathbb{N}$, $\nu(1_n) = b_n$. Then there is $k \in \mathbb{N}$ such that

$$(1) \quad t(i) = c_k.$$

(The symbol $t(i)$ means the i th coordinate of the element t .)

We have

$$f^k(1_k) = 1,$$

thus

$$f^k(b_k) = f^k(\nu(1_k)) = \nu(f^k(1_k)) = \nu(1) = t,$$

hence (1) implies

$$c_k = t(i) = (f^k(b_k))(i) = f^k(b_k(i)),$$

i.e.,

$$b_k(i) \in f^{-k}(c_k) = \emptyset,$$

a contradiction. □

4.3. Lemma. *If $n \in \mathbb{N}$, then there exist $i_n \in I$, $m_n \in \mathbb{N}$ and a prime p_n such that*

- (a) $p_1^{m_1} < p_2^{m_2} < \dots$,
- (b) \mathcal{B}_{i_n} is a cycle with $p_n^{m_n}$ elements.

Proof. By 4.2, if $i \in I$, then \mathcal{B}_i contains a cycle. If the cardinalities of these cycles are bounded, then each connected component of $\prod_{i \in I} \mathcal{B}_i$ contains a cycle, thus each subalgebra of $\prod_{i \in I} \mathcal{B}_i$ contains a cycle, hence (M, f) and \mathcal{A} , too, contain a cycle, which is a contradiction. Therefore the assertion is valid according to 4.1 and 4.2. □

4.4. Lemma. *There exist distinct elements $b_n \in \prod_{i \in I} \mathcal{B}_i$ for $n \in \mathbb{N}$ such that $f(b_{n+1}) = b_n$ for each $n \in \mathbb{N}$.*

Proof. Assume that, for each $n \in \mathbb{N}$, \mathcal{B}_{i_n} is as in 4.3. By 4.2, if $i \in I$, then \mathcal{B}_i contains a cycle; take an arbitrary element

$$b_1 \in \prod_{i \in I} \mathcal{B}_i$$

such that $b_1(i)$ belongs to the cycle of \mathcal{B}_i . By induction, if $k \in \mathbb{N}$, $k > 1$ and b_j is defined for each $j \in \mathbb{N}$, $j < k$, then let b_k be the (unique) element of $\prod_{i \in I} \mathcal{B}_i$ such that

- (1) $b_k(i)$ belongs to the cycle of \mathcal{B}_i ,
- (2) $f(b_k(i)) = b_{k-1}(i)$.

Then obviously $f(b_{n+1}) = b_n$ for each $n \in \mathbb{N}$.

Suppose that there are $k, l \in \mathbb{N}$, $k < l$ such that $b_k = b_l$. In view of 4.3 (a) there exists $n \in \mathbb{N}$ such that

$$(3) \quad l - k < p_n^{m_n}.$$

We have

$$b_k = f^{l-k}(b_l) = f^{l-k}(b_k),$$

thus

$$(4) \quad b_k(i_n) = (f^{l-k}(b_k))(i_n) = f^{l-k}(b_k(i_n)).$$

The element $b_k(i_n)$ belongs to \mathcal{B}_{i_n} , i.e., to a cycle with $p_n^{m_n}$ elements, therefore (3) and (4) yield a contradiction. \square

4.5. Corollary. *No retract of $\prod_{i \in I} \mathcal{B}_i$ is isomorphic to \mathcal{A} .*

Proof. For $n \in \mathbb{N}$ let b_n be as in 4.4. If Q is a retract of $\prod_{i \in I} \mathcal{B}_i$ and φ is a corresponding retraction endomorphism, then either

- (a) $\varphi(b_1)$ belongs to a cycle, or
- (b) there are distinct elements $q_n \in Q$ for $n \in \mathbb{N}$ such that $\varphi(b_n) = q_n$ for each $n \in \mathbb{N}$.

Let Q be a retract of $\prod_{i \in I} \mathcal{B}_i$, $(Q, f) \cong \mathcal{A}$. Then (a) fails to hold. If (b) is valid, then, for each $n \in \mathbb{N}$,

$$f(q_{n+1}) = f(\varphi(b_{n+1})) = \varphi(f(b_{n+1})) = \varphi(b_n) = q_n,$$

which is again a contradiction to the definition of \mathcal{A} . \square

As a corollary we obtain

4.6. Proposition. \mathcal{A} possesses no representation $\{\mathcal{B}_i: i \in I\}$ of \mathcal{A} in \mathcal{U}_c such that each \mathcal{B}_i for $i \in I$ is retract irreducible in \mathcal{U}_c .

5. THE CLASS $\mathcal{K} = \mathcal{U}$

Let \mathcal{A} be as in the previous sections and suppose that the system $\{\mathcal{B}_i: i \in I\} \subseteq \mathcal{U}$ is a representation of \mathcal{A} such that if $i \in I$, then \mathcal{B}_i is retract irreducible in \mathcal{U} . In view of [5, Thm. 4.5] we obtain

5.1. Lemma. *If $i \in I$, then one of the following conditions is satisfied:*

- (i) \mathcal{B}_i contains a cycle with p^m elements, where p is a prime and $m \in \mathbb{N}$,
- (ii) $\mathcal{B}_i \cong (\mathbb{N}, f)$,
- (iii) \mathcal{B}_i contains a one element cycle $\{c\}$ and if $\{a, b\} \subseteq \mathcal{B}_i$ with $f(a) = f(b)$, then either $a=b$ or $c \in \{a, b\}$.

Analogously as above the following assertions can be proved:

5.2. Lemma. *If $i \in I$, then (ii) fails to hold.*

5.3. Lemma. *If $n \in \mathbb{N}$, then there exist $i_n \in I$, $m_n \in \mathbb{N}$ and a prime p_n such that*

- (a) $p_1^{m_1} < p_2^{m_2} < \dots$,
- (b) \mathcal{B}_{i_n} contains a cycle with $p_n^{m_n}$ elements.

5.4. Lemma. *There exist distinct elements $b_n \in \prod_{i \in I} \mathcal{B}_i$ for $n \in \mathbb{N}$ such that $f(b_{n+1}) = b_n$ for each $n \in \mathbb{N}$.*

5.5. Lemma. *No retract of $\prod_{i \in I} \mathcal{B}_i$ is isomorphic to \mathcal{A} .*

5.6. Proposition. \mathcal{A} possesses no representation $\{\mathcal{B}_i: i \in I\}$ of \mathcal{A} in \mathcal{U} such that \mathcal{B}_i for each $i \in I$ is retract irreducible in \mathcal{U} .

References

- [1] *D. Duffus and I. Rival*: A structure theory for ordered sets. *Discrete Math.* 35 (1981), 53–118.
- [2] *D. Jakubíková-Studenovská*: Retract irreducibility of connected monounary algebras I. *Czechoslovak Math. J.* 46(121) (1996), 291–308.
- [3] *D. Jakubíková-Studenovská*: Retract irreducibility of connected monounary algebras II. *Czechoslovak Math. J.* 47(122) (1997), 113–126.
- [4] *D. Jakubíková-Studenovská*: Two types of retract irreducibility of connected monounary algebras. *Math. Bohem.* 2(121) (1996), 143–150.
- [5] *D. Jakubíková-Studenovská*: Retract irreducibility of monounary algebras. *Czechoslovak Math. J.* 49(124) (1999), 363–390.
- [6] *D. Jakubíková-Studenovská*: DR-irreducibility of monounary algebras with a cycle. *Czechoslovak Math. J.* 50(125) (2000), 681–698.
- [7] *D. Jakubíková-Studenovská*: DR-irreducibility of monounary algebras. *Czechoslovak Math. J.* 50(125) (2000), 705–720.

Author's address: P. J. Šafárik University, Faculty of Science, Institute of Mathematics, Jesenná 5, SK-041 54 Košice, Slovak Republic, e-mail: studenovska@science.upjs.sk.