

Ali Ahmed Nough

On convergence theory in fuzzy topological spaces and its applications

Czechoslovak Mathematical Journal, Vol. 55 (2005), No. 2, 295–316

Persistent URL: <http://dml.cz/dmlcz/127979>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON CONVERGENCE THEORY IN FUZZY TOPOLOGICAL SPACES AND ITS APPLICATIONS

ALI AHMED NOUH, Mansoura

(Received May 29, 2002)

Abstract. In this paper we introduce and study new concepts of convergence and adherent points for fuzzy filters and fuzzy nets in the light of the Q -relation and the Q -neighborhood of fuzzy points due to Pu and Liu [28]. As applications of these concepts we give several new characterizations of the closure of fuzzy sets, fuzzy Hausdorff spaces, fuzzy continuous mappings and strong Q -compactness. We show that there is a relation between the convergence of fuzzy filters and the convergence of fuzzy nets similar to the one which exists between the convergence of filters and the convergence of nets in topological spaces.

Keywords: fuzzy points, Q -neighborhoods, fuzzy filters, fuzzy nets, limit, adherent and Q -adherent points of fuzzy filters and fuzzy nets, fuzzy continuity, strong Q -compactness

MSC 2000: 54A20, 54A40, 54C08, 54H12

1. INTRODUCTION

The notion of convergence is one of the basic notion in analysis. There are two different convergence theories used in general topology that lead to equivalent results. One of them is based on the notion of a net from 1922 due to Moore and Smith [25]; another one, which goes back to the work of Cartan [5] in 1937 and Bourbaki [4] in 1940, is based on the notion of a filter. In 1955 Bartle [1] studied the relation between filters and nets convergence theories, and he proved that they are equivalent. In 1979 Lowen [20] introduced and studied the theory of convergence for fuzzy filters (prefilters) in fuzzy topological spaces and applied the results to describe fuzzy compactness and fuzzy continuity. This was followed by an extensive study of the convergence theory of fuzzy filters in fuzzy topological spaces by several authors [7], [9]–[12], [21], [22], [26], [27], [31] from different standpoints. However, the concepts of Q -relation and Q -neighborhood of fuzzy points have not been used in these approaches. In 1980 Pu and Liu [28], by means of the concept of a Q -neighborhood

of a fuzzy point in a fuzzy topological space, gave the notion of convergence for fuzzy nets. The relationship between the two types of convergence was studied by Lowen in [21]. This was also followed by an extensive study of the convergence theory in fuzzy topological spaces by several authors [7], [8], [13], [21]–[24], [28]–[30] from different standpoints. In this paper, we will give new concepts of convergence and adherence for fuzzy filters (prefilters in the sense of Lowen [20]) and fuzzy nets in fuzzy topological spaces. These concepts depend on the notions of Q -relation and Q -neighborhood of a fuzzy point given by Liu and Pu [28], as these concepts seem to be extremely suitable for the fuzzy situation. Then several characterizations of some fuzzy topological concepts, namely: the closure of a fuzzy set, fuzzy Hausdorff spaces, fuzzy continuous mappings and strong Q -compactness are given by means of convergence of fuzzy filters and fuzzy nets. Finally, we show that there is a relation between the convergence of fuzzy nets and the convergence of fuzzy filters similar to the one which exists between the convergence of nets and filters in topological spaces.

2. PRELIMINARIES

Let X be an arbitrary nonempty set. A fuzzy set in X is a mapping from X to the closed unit interval $I = [0, 1]$, that is, an element of I^X . A fuzzy point (singleton) x_α is a fuzzy set in X defined by $x_\alpha(x) = \alpha$ and $x_\alpha(y) = 0$ for all $y \neq x$, whose support is the single point x and whose value is $\alpha \in (0, 1]$. We denote by $\text{FP}(X)$ the collection of all fuzzy points in X . A family $\mathcal{A} \subseteq I^X$ is said to have finite intersection property (FIP, for short) if any finite intersection of elements of \mathcal{A} is nonempty.

Throughout this paper X, Y, Z , etc. denote ordinary sets while $\mu, \eta, \lambda, \varrho$, etc. denote fuzzy sets defined on an ordinary set. For $A \subseteq X$, by 1_A we mean the characteristic mapping of X to $\{0, 1\}$. Also, we use F to denote fuzzy and fts to denote a fuzzy topological space. The fuzzy set theoretical and fuzzy topological concepts used in this paper are standard and can be found in Zadeh [32], Klir and Yuan [18], Chang [6], Belohlavek [2], Lowen [19] and Pu and Liu [28]. However, we feel it necessary to fix some notions and recall a few concepts from [2], [6], [17], [18], [20], [28], [30], [32].

Definition 2.1 [18], [32]. Let $\mu, \eta \in I^X$. We define the following fuzzy sets:

- (i) $\mu \wedge \eta \in I^X$, by $(\mu \wedge \eta)(x) = \min\{\mu(x), \eta(x)\}$ for each $x \in X$ (intersection).
- (ii) $\mu \vee \eta \in I^X$, by $(\mu \vee \eta)(x) = \max\{\mu(x), \eta(x)\}$ for each $x \in X$ (union).
- (iii) $\mu^c \in I^X$, by $\mu^c(x) = 1 - \mu(x)$ for each $x \in X$ (complement).

Definition 2.2 [17]. Let $\mu \in I^X$. We define the following crisp subsets:

- (i) $\mu_{w\alpha} = \{x \in X : \mu(x) \geq \alpha\}$, the weak α -cut of μ , $\alpha \in (0, 1]$.
- (ii) $\text{supp}(\mu) = \{x \in X : \mu(x) > 0\}$, the support of μ .

Definition 2.3 [6]. Let $\tau \subseteq I^X$ satisfy the following three conditions:

- (i) $X, \emptyset \in \tau$.
- (ii) If $\mu_1, \mu_2 \in \tau$, then $\mu_1 \wedge \mu_2 \in \tau$.
- (iii) If $\{\mu_j : j \in J\} \subseteq \tau$, then $\bigvee_{j \in J} \mu_j \in \tau$.

Then τ is called a fuzzy topology on X and (X, τ) is called a fuzzy topological space (fts, for short). The elements of τ are called open fuzzy sets. A fuzzy set is called a closed fuzzy set if $\mu^c \in \tau$. We denote by τ' the collection of all closed fuzzy sets in X .

One may easily verify the following example.

Example 2.4. Let (X, T) be a topological space, $\mu \in I^X$ and $A \subseteq X$. Then:

- (i) The classes $\tau_1 = \{X, \emptyset\}$, $\tau_2 = I^X$, $\tau_3 = \{\mu \in I^X : \text{supp}(\mu) \in T\}$ and $\tau_4 = \{1_A : A \in T\}$ are fuzzy topologies on X .
- (ii) If X is an infinite set, then the class $\tau_\infty = \{\mu \in I^X : \text{supp}(\mu^c) \text{ is finite set}\} \vee \{\emptyset\}$ is a fuzzy topology on X ; the so-called co-finite fuzzy topology on X .

Definition 2.5 [28]. Let (X, τ) be a fts and $\mu, \varrho \in I^X$. Then:

- (i) μ is called a quasi-coincident with ϱ , denoted by $\mu q \varrho$, if there exists $x \in X$ such that $\mu(x) + \varrho(x) > 1$. If μ is not quasi-coincident with ϱ , then we write $\mu \bar{q} \varrho$.
- (ii) μ is called Q -neighborhood of a fuzzy point $x_\alpha \in \text{FP}(X)$ if there exists an open fuzzy set $\eta \in \tau$ such that $x_\alpha q \eta$ and $\eta \leq \mu$. The class of all open Q -neighborhoods of x_α is denoted by $N_{x_\alpha}^Q$.
- (iii) ϱ is called a Q_α -neighborhood of a fuzzy set μ iff ϱ is a Q -neighborhood of each fuzzy point $x_\alpha \leq \mu$. The class of all open Q_α -neighborhoods of μ will be denoted by $N_\mu^{Q_\alpha}$.

Definition 2.6 [2], [28]. Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and $\mu \in I^X$. The closure of μ , denoted by $\text{cl}(\mu)$, is defined by: $x_\alpha \leq \text{cl}(\mu)$ iff $(\forall \eta \in N_{x_\alpha}^Q)(\eta q \mu)$. The fuzzy set μ is closed if $\mu = \text{cl}(\mu)$.

Theorem 2.7 [28]. Let (X, τ) be a fts and $\mu \in I^X$. Then:

- (i) μ is open iff $(\forall x_\alpha q \mu)(\exists \eta \in N_{x_\alpha}^Q)(\eta \leq \mu)$.
- (ii) For each $\eta \in \tau$, $\eta q \mu$ iff $\eta q \text{cl}(\mu)$.

Definition 2.8 [28]. A fts (X, τ) is called a fuzzy Hausdorff space (FT_2 , for short) iff $(\forall x_\varepsilon, y_\nu \in \text{FP}(X), x \neq y)(\exists \mu \in N_{x_\varepsilon}^Q)(\exists \eta \in N_{y_\nu}^Q)(\mu \wedge \eta = \emptyset)$.

3. NEW CONVERGENCE THEORY OF FUZZY FILTERS

In this section we will introduce and study the notions of convergence, adherence and Q -adherence for fuzzy filters and filter bases in fuzzy topological spaces by means of the concept of a Q -neighborhood of a fuzzy point given by Liu and Pu [28]. This will enable us to give some results about fuzzy Hausdorff spaces [18], [28], closure of a fuzzy set [18], [28], fuzzy continuity [14], [28] and about strong Q -compactness [33], similar to the ones which hold in topological spaces.

Definition 3.1 [20]. A fuzzy filter on X is a nonempty subset $\mathcal{F} \subseteq I^X$ such that

- (i) $\emptyset \notin \mathcal{F}$.
- (ii) If $\mu \in \mathcal{F}$ and $\mu \leq \eta$, then $\eta \in \mathcal{F}$.
- (iii) If $\mu_1, \mu_2 \in \mathcal{F}$, then $\mu_1 \wedge \mu_2 \in \mathcal{F}$.

The class of all fuzzy filters on X will be denoted by $\text{FL}(I^X)$.

Definition 3.2 [20]. A fuzzy filter base on X is a nonempty subset $\beta \subseteq I^X$ such that:

- (i) $\emptyset \notin \beta$.
- (ii) If $\mu_1, \mu_2 \in \beta$, then $\exists \mu_3 \in \beta$ such that $\mu_3 \leq \mu_1 \wedge \mu_2$.

The class of all fuzzy filter bases on X will be denoted by $\text{FLB}(I^X)$.

The fuzzy filter \mathcal{F} generated by β is defined by $\mathcal{F} = \{\mu \in I^X : \eta \leq \mu \text{ for some } \eta \in \beta\}$ and is denoted by $\langle \beta \rangle$. A collection β of subsets of \mathcal{F} is a base for \mathcal{F} if for each $\mu \in \mathcal{F}$ there is $\eta \in \beta$ such that $\eta \leq \mu$.

Definition 3.3 [20]. Let $\mathcal{F}, \mathcal{H} \in \text{FL}(I^X)$ (or $\text{FLB}(I^X)$). Then we say that \mathcal{F} is finer than \mathcal{H} , written as $\mathcal{H} < \mathcal{F}$, if $(\forall \mu \in \mathcal{H})(\exists \varrho \in \mathcal{F})(\varrho \leq \mu)$.

Definition 3.4 [20]. Let X be a nonempty set. Then:

- (i) A fuzzy filter \mathcal{F} on X is called a prime if for all $\mu, \eta \in I^X$, $\mu \vee \eta \in \mathcal{F}$ implies that either $\mu \in \mathcal{F}$ or $\eta \in \mathcal{F}$.
- (ii) A fuzzy filter \mathcal{F} on X is called a maximal fuzzy filter on X iff \mathcal{F} is finer than every fuzzy filter comparable with it.
- (iii) A fuzzy filter base β on X is called a maximal fuzzy filter base on X if it is a base for a maximal fuzzy filter on X .
- (iv) A subfamily ξ of fuzzy filter \mathcal{F} on X is said to be a subbase for \mathcal{F} if the family of all finite intersections of members of ξ is a base for \mathcal{F} . We say that ξ generates \mathcal{F} .

If $\mathcal{P}(\mathcal{F})$ denotes the family of all prime fuzzy filters finer than \mathcal{F} , then it is easy to verify that $\mathcal{F} = \bigcap_{\mathcal{P} \in \mathcal{P}(\mathcal{F})} \mathcal{P}$.

The class of all maximal fuzzy filters and maximal fuzzy filter bases on X will be denoted by $\text{MFL}(I^X)$ and $\text{MFLB}(I^X)$, respectively.

Theorem 3.5. *If \mathcal{C} is a nonempty family of fuzzy subsets of X having the FIP, then there exists a fuzzy filter base on X containing \mathcal{C} .*

Proof. Similar to the proof of the corresponding one in the crisp case [15]. \square

Theorem 3.6. *Let ξ be a family of nonempty fuzzy subsets of X which has FIP. Then there exists a fuzzy filter \mathcal{F} on X having ξ as a subbase iff ξ has FIP.*

Proof. Follows immediately from Definition 3.4 (iv) and Theorem 3.5. \square

Theorem 3.7 [31]. *For an $\mathcal{F} \in \text{MFL}(I^X)$, the following statements are true:*

- (i) \mathcal{F} is a prime fuzzy filter.
- (ii) For every $\mu \in I^X$, either $\mu \in \mathcal{F}$ or $1 - \mu \in \mathcal{F}$.
- (iii) Let $\mu \in I^X$. If $\mu \notin \mathcal{F}$, then there is $\varrho \in \mathcal{F}$ such that $\mu \wedge \varrho = \emptyset$.

Theorem 3.8 [26]. *Let $f: X \rightarrow Y$ be a mapping.*

- (i) If β is a fuzzy filter base on X , then so is $f(\beta) = \{f(\lambda): \lambda \in \beta\}$ on Y .
- (ii) If β is a fuzzy filter base on Y and f is onto, then $f^{-1}(\beta) = \{f^{-1}(\lambda): \lambda \in \beta\}$ is a fuzzy filter base on X .

For further details on fuzzy filters and related concepts, see [20], [26].

Definition 3.9. A fuzzy filter (filter base) \mathcal{F} on X is called:

- (i) an upper fuzzy filter (filter base) on X if for every $\lambda \in \mathcal{F}$, $\lambda(x) > \frac{1}{2}$ for some $x \in X$;
- (ii) on $\mu \in I^X$ if $\lambda \wedge \mu \neq \emptyset$ for each $\lambda \in \mathcal{F}$.

Definition 3.10. A fuzzy filter (filter base) \mathcal{F} in a fts (X, τ) is said to be:

- (i) convergent to a fuzzy point x_α , denoted by $\mathcal{F} \rightarrow x_\alpha$, iff $(\forall \eta \in N_{x_\alpha}^Q) (\exists \varrho \in \mathcal{F})(\varrho \leq \eta)$. The limit of a fuzzy filter base \mathcal{F} is defined by $\lim(\mathcal{F}) = \bigvee \{x_\alpha \in \text{FP}(X): \mathcal{F} \rightarrow x_\alpha\}$.
- (ii) convergent to a fuzzy set μ , denoted by $\mathcal{F} \rightarrow \mu$, iff for each open Q_α -cover \mathcal{U} of μ (see Definition 3.26 below), there exist a finite subcollection \mathcal{U}_0 of \mathcal{U} and an $\lambda \in \mathcal{F}$ such that $\lambda \leq \bigvee \{\eta: \eta \in \mathcal{U}_0\}$.

Definition 3.11. Let (X, τ) be a fts and \mathcal{F} a fuzzy filter (filter base) on X . A fuzzy point x_α is said to be an adherent (Q -adherent) point for a fuzzy filter (filter base) \mathcal{F} , denoted by $\mathcal{F} \propto x_\alpha$ ($\mathcal{F} \overset{Q}{\propto} x_\alpha$), iff for all $\eta \in N_{x_\alpha}^Q$ and for all $\varrho \in \mathcal{F}$ we have $\varrho \wedge \eta \neq \emptyset$ ($\varrho q \eta$). The adherence and Q -adherence of the fuzzy filter (filter base) \mathcal{F} is defined by $\text{adh}(\mathcal{F}) = \bigvee \{x_\alpha \in \text{FP}(X): \mathcal{F} \propto x_\alpha\}$ and $Q\text{-adh}(\mathcal{F}) = \bigvee \{x_\alpha \in \text{FP}(X): \mathcal{F} \overset{Q}{\propto} x_\alpha\}$, respectively.

The proof of the following result is straightforward and therefore it is omitted.

Theorem 3.12. *Let (X, τ) be a fts and $\mathcal{F} \in \text{FL}(I^X)$ (or $\text{FLB}(I^X)$). Then*

- (i) $N_{x_\alpha}^Q$ is a fuzzy filter (filter base) on X and $N_{x_\alpha}^Q \rightarrow x_\alpha$.
- (ii) $\mathcal{F} \rightarrow x_\alpha$ iff $x_\alpha \leq \lim(\mathcal{F})$ iff $N_{x_\alpha}^Q < \mathcal{F}$.
- (iii) $\mathcal{F} \times x_\alpha$ ($\mathcal{F} \overset{Q}{\times} x_\alpha$) iff $x_\alpha \leq \text{adh}(\mathcal{F})$ ($x_\alpha \leq Q\text{-adh}(\mathcal{F})$).
- (iv) $\lim(\mathcal{F}) \leq \text{adh}(\mathcal{F})$ and $Q\text{-adh}(\mathcal{F}) \leq \text{adh}(\mathcal{F})$ but $\lim(\mathcal{F}) \not\leq Q\text{-adh}(\mathcal{F})$. If \mathcal{F} is an upper fuzzy filter (filter base), then $\lim(\mathcal{F}) \leq Q\text{-adh}(\mathcal{F})$.
- (v) If $\mathcal{F} < \mathcal{F}^*$, then $\lim(\mathcal{F}) \leq \lim(\mathcal{F}^*)$, $\text{adh}(\mathcal{F}^*) \leq \text{adh}(\mathcal{F})$ and $Q\text{-adh}(\mathcal{F}^*) \leq Q\text{-adh}(\mathcal{F})$.
- (vi) If $\mathcal{F} \rightarrow \mu$, then $N_\mu^Q < \mathcal{F}$, where N_μ^Q is a fuzzy filter base on X .
- (vii) If $\mathcal{F} \rightarrow x_\alpha$ ($\mathcal{F} \times x_\alpha$, $\mathcal{F} \overset{Q}{\times} x_\alpha$) and if $\nu \leq \alpha$, then $\mathcal{F} \rightarrow x_\nu$ ($\mathcal{F} \times x_\nu$, $\mathcal{F} \overset{Q}{\times} x_\nu$, respectively).
- (viii) If $\mathcal{F} \rightarrow x_\alpha$ and $x_\alpha \leq \mu$, then $\mathcal{F} \rightarrow \mu$.
- (ix) If $\mathcal{F} < \mathcal{F}^*$ and $\mathcal{F} \rightarrow \mu$, then $\mathcal{F}^* \rightarrow \mu$.
- (x) If β is a base for a fuzzy filter \mathcal{F} on X , then $\beta \rightarrow x_\alpha$ ($\beta \times x_\alpha$, $\beta \overset{Q}{\times} x_\alpha$, $\beta \rightarrow \mu$) iff $\mathcal{F} \rightarrow x_\alpha$ ($\mathcal{F} \times x_\alpha$, $\mathcal{F} \overset{Q}{\times} x_\alpha$, $\mathcal{F} \rightarrow \mu$, respectively).

Theorem 3.13. *Let (X, τ) be a fts. The fuzzy point x_α is an adherent point for a fuzzy filter base β on X iff there exists a fuzzy filter base β^* on X finer than β and converging to x_α .*

Proof. If x_α is an adherent point of β , then $(\forall \eta \in N_{x_\alpha}^Q)(\forall \varrho \in \beta)(\varrho \wedge \eta \neq \emptyset)$. Then the family $\xi = \beta \vee N_{x_\alpha}^Q$ forms a subbase for some fuzzy filter base β^* on X finer than β and $\beta^* \rightarrow x_\alpha$. Conversely, if there exists a fuzzy filter base β^* on X finer than β and $\beta^* \rightarrow x_\alpha$, then by Theorem 3.12 (iv), $\beta^* \times x_\alpha$ and so by Theorem 3.12 (v), $\beta \rightarrow x_\alpha$. \square

Theorem 3.14. *Let (X, τ) be a fts and β an upper fuzzy filter base on X . A fuzzy point x_α is a Q -adherent point for β iff there exists an upper fuzzy filter base β^* on X finer than β and converging to x_α .*

Proof. It is similar to that of Theorem 3.13 taking into consideration that β and β^* are upper fuzzy filter bases. \square

Theorem 3.15. *Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and $\mu \in I^X$. Then $x_\alpha \leq \text{cl}(\mu)$ iff there exists a fuzzy filter \mathcal{F} on X with $\mathcal{F} \rightarrow x_\alpha$ and $\varrho q \mu$ for each $\varrho \in \mathcal{F}$.*

Proof. Let $x_\alpha \leq \text{cl}(\mu)$. Then $\mu q \eta$ for each $\eta \in N_{x_\alpha}^Q$. So it suffices to take $\mathcal{F} = N_{x_\alpha}^Q$. Conversely, let $\mathcal{F} \rightarrow x_\alpha$ and $\varrho q \mu$ for each $\varrho \in \mathcal{F}$. Then $N_{x_\alpha}^Q \leq \mathcal{F}$ and so $x_\alpha \leq \text{cl}(\mu)$. \square

Theorem 3.16. *Let (X, τ) be a fts and $\mu \in I^X$. Then μ is open iff μ is a member of each fuzzy filter on X converging to a fuzzy point quasi-coincident with μ .*

Proof. Suppose that $\mu \in I^X$ is open and let \mathcal{F} be an arbitrary fuzzy filter on X converging to x_α and $x_\alpha q \mu$. Since $\mu \in N_{x_\alpha}^Q$, there exists $\lambda \in \mathcal{F}$ such that $\lambda \leq \mu$. Hence $\mu \in \mathcal{F}$. Conversely, for each $x_\alpha q \mu$ we consider $\mathcal{F} = N_{x_\alpha}^Q$. Then $\mu \in N_{x_\alpha}^Q$ for each $x_\alpha q \mu$ and so $\mu \in \tau$. \square

The set of fuzzy points to which a fuzzy filter converges is, in general, infinite. But if we put certain restrictions on the supports, we can obtain a result concerning the uniqueness of convergence.

Theorem 3.17. *A fts (X, τ) is FT_2 iff no fuzzy filter \mathcal{F} on X converges to two fuzzy points with different supports.*

Proof. Let (X, τ) be an FT_2 -space and \mathcal{F} a fuzzy filter on X such that $\mathcal{F} \rightarrow x_\alpha$ and y_α such that $x \neq y$. Since $\mathcal{F} \rightarrow x_\alpha$, we have $(\forall \mu \in N_{x_\alpha}^Q)(\exists \rho_1 \in \mathcal{F})(\rho_1 \leq \mu)$. Also, since $\mathcal{F} \rightarrow y_\alpha$, we have $(\forall \eta \in N_{y_\alpha}^Q)(\exists \rho_2 \in \mathcal{F})(\rho_2 \leq \eta)$. Since $\rho_1, \rho_2 \in \mathcal{F}$, then $\rho_3 = \rho_1 \wedge \rho_2 \in \mathcal{F}$ such that $\rho_3 \leq \mu \wedge \eta$ and so $\mu \wedge \eta \neq \emptyset$, a contradiction. Conversely, suppose that no fuzzy filter \mathcal{F} on X converges to two fuzzy points with different supports and (X, τ) is not an FT_2 -space, then $(\forall \mu \in N_{x_\alpha}^Q)(\forall \eta \in N_{y_\alpha}^Q)(\mu \wedge \eta \neq \emptyset)$. Then it is easy to verify that the family $\beta = \{\mu \wedge \eta : \mu \in N_{x_\alpha}^Q \text{ and } \eta \in N_{y_\alpha}^Q\}$ is a fuzzy filter base on X converging to both x_α and y_α . Let $\mathcal{F} = \langle \beta \rangle$. Then by Theorem 3.12 (x), \mathcal{F} converges to both x_α and y_α , a contradiction. \square

Theorem 3.18. *Let (X, τ) be a fts and let \mathcal{F} be a fuzzy filter on X . Then:*

- (i) $\lim(\mathcal{F})$, $\text{adh}(\mathcal{F})$ and $Q\text{-adh}(\mathcal{F})$ are closed fuzzy sets.
- (ii) If \mathcal{F} is a maximal fuzzy filter, then $\lim(\mathcal{F}) = \text{adh}(\mathcal{F})$.
- (iii) If \mathcal{F} is a maximal upper fuzzy filter, then $\lim(\mathcal{F}) = Q\text{-adh}(\mathcal{F})$.

Proof. (i) Since $\lim(\mathcal{F}) \leq \text{cl}(\lim(\mathcal{F}))$ it suffices to show that $\text{cl}(\lim(\mathcal{F})) \leq \lim(\mathcal{F})$. Let $x_\alpha \leq \text{cl}(\lim(\mathcal{F}))$ and $\eta \in N_{x_\alpha}^Q$. Then $\eta q \lim(\mathcal{F})$ and so there exists $y \in X$ such that $\eta(y) + \lim(\mathcal{F})(y) > 1$. Put $t = \lim(\mathcal{F})(y)$. Then $\eta \in N_{y_t}^Q$ and $y_t \leq \lim(\mathcal{F})$. Then for each $\rho \in N_{y_t}^Q$, there exists $\lambda \in \mathcal{F}$ such that $\lambda \leq \rho$. Thus $\lambda \leq \eta$ and so $x_\alpha \leq \lim(\mathcal{F})$. Hence $\text{cl}(\lim(\mathcal{F})) \leq \lim(\mathcal{F})$. Thus $\lim(\mathcal{F})$ is a closed fuzzy set. Similarly, $\text{adh}(\mathcal{F})$ and $Q\text{-adh}(\mathcal{F})$ are closed fuzzy sets.

(ii) follows from Theorem 3.12 (iv) and Theorem 3.13.

(iii) follows from Theorem 3.12 (iv) and Theorem 3.14. \square

Theorem 3.19. If $\{\mathcal{F}_j : j \in J\}$ is a family of fuzzy filters on X and $\mathcal{F} = \bigcap_{j \in J} \mathcal{F}_j$,

then:

- (i) $\lim(\mathcal{F}) = \bigwedge_{j \in J} \lim(\mathcal{F}_j)$.
- (ii) $\mathcal{F} \rightarrow \mu$ iff $\mathcal{F}_j \rightarrow \mu$ for each $j \in J$.

Proof. (i) Since $\mathcal{F} < \mathcal{F}_j$ for all $j \in J$, we have $\lim(\mathcal{F}) \leq \lim(\mathcal{F}_j)$ for all $j \in J$ and so $\lim(\mathcal{F}) \leq \bigwedge_{j \in J} \lim(\mathcal{F}_j)$. Conversely, if $x_\alpha \leq \bigwedge_{j \in J} \lim(\mathcal{F}_j)$, then $x_\alpha \leq \lim(\mathcal{F}_j)$ for all $j \in J$, then $(\forall \eta \in N_{x_\alpha}^Q)(\eta \in \mathcal{F}_j)$ for all $j \in J$. So $\eta \in \mathcal{F}$ and hence $x_\alpha \leq \lim(\mathcal{F})$. Thus $\bigwedge_{j \in J} \lim(\mathcal{F}_j) \leq \lim(\mathcal{F})$.

(ii) Since $\mathcal{F} < \mathcal{F}_j$ for all $j \in J$, then by Theorem 3.12 (ix), $\mathcal{F} \rightarrow \mu$ implies $\mathcal{F}_j \rightarrow \mu$ for all $j \in J$. Conversely, if $\mathcal{F}_j \rightarrow \mu$ for each $j \in J$, then for each open Q_α -cover \mathcal{U} of μ there exist a finite subset \mathcal{U}_0 of \mathcal{U} and a $\lambda_j \in \mathcal{F}_j$ such that $\lambda_j \leq \bigvee\{\eta : \eta \in \mathcal{U}_0\}$ for each $j \in J$. So $\bigwedge_{j \in J} \lambda_j \leq \bigvee\{\eta : \eta \in \mathcal{U}_0\}$. Put $\lambda = \bigwedge_{j \in J} \lambda_j$. Then $\lambda \in \mathcal{F}$ and $\lambda \leq \bigvee\{\eta : \eta \in \mathcal{U}_0\}$. Hence $\mathcal{F} \rightarrow \mu$. \square

Corollary 3.20. For a fuzzy filter \mathcal{F} on X , we have $\mathcal{F} \rightarrow x_\alpha$ ($\mathcal{F} \rightarrow \mu$) iff $\mathcal{P} \rightarrow x_\alpha$ ($\mathcal{P} \rightarrow \mu$) for each $\mathcal{P} \in P(\mathcal{F})$.

It is well known that continuity can be completely characterized by means of the convergence of filter bases. In the following, we characterize the fuzzy continuity between fuzzy topological spaces by means of convergent fuzzy filter bases.

Definition 3.21 [14], [28]. A mapping $f : (X, \tau) \rightarrow (Y, \Delta)$ is called F -continuous if $(\forall x_\alpha \in \text{FP}(X))(\forall \eta \in N_{f(x_\alpha)}^Q)(\exists \mu \in N_{x_\alpha}^Q)(f(\mu) \leq \eta)$.

Theorem 3.22. Let $f : (X, \tau) \rightarrow (Y, \Delta)$ be a mapping. Then the following are equivalent:

- (i) f is F -continuous.
- (ii) For each $x_\alpha \in \text{FP}(X)$, $f(N_{x_\alpha}^Q) \rightarrow f(x_\alpha)$.
- (iii) For each fuzzy filter base β on X , $\beta \rightarrow \mu$ implies $f(\beta) \rightarrow f(\mu)$.
- (iv) For each fuzzy filter base β on X , $\beta \rightarrow x_\alpha$ implies $f(\beta) \rightarrow f(x_\alpha)$ for each $x_\alpha \in \text{FP}(X)$.
- (v) For each prime fuzzy filter \mathcal{P} on X , $\mathcal{P} \rightarrow \mu$ implies $f(\mathcal{P}) \rightarrow f(\mu)$.
- (vi) For each prime fuzzy filter \mathcal{P} on X , $\mathcal{P} \rightarrow x_\alpha$ implies $f(\mathcal{P}) \rightarrow f(x_\alpha)$ for each $x_\alpha \in \text{FP}(X)$.
- (vii) For each fuzzy filter base β on X , $f(\lim(\beta)) \leq \lim(f(\beta))$.
- (viii) For each prime fuzzy filter \mathcal{P} on X , $f(\lim(\mathcal{P})) \leq \lim(f(\mathcal{P}))$.

Proof. (i) \iff (ii): The statement that f is F -continuous is $(\forall x_\alpha \in \text{FP}(X))(\forall \eta \in N_{f(x_\alpha)}^Q)(\exists \mu \in N_{x_\alpha}^Q)(f(\mu) \leq \eta)$ and this is exactly the statement that the fuzzy filter base $f(N_{x_\alpha}^Q) \rightarrow f(x_\alpha)$.

(i) \implies (iii): Let $\mu \in I^X$ and let β be a fuzzy filter base on X such that $\beta \rightarrow \mu$. Let $\mathcal{U} = \{\eta_j : j \in J\}$ be an open Q_α -cover of $f(\mu)$. Then $(\forall x \in X \text{ with } \mu(x) \geq \alpha)(\exists j \in J)(\eta_j \in N_{f(x_\alpha)}^Q)$. By F -continuity of f , $(\exists \sigma_j \in N_{x_\alpha}^Q)(f(\sigma_j) \leq \eta_j)$. Then the family $\{\sigma_j : j \in J\}$ is an open Q_α -cover of μ . Since $\beta \rightarrow \mu$, there exist a finite subset J_0 of J and a $\lambda \in \beta$ such that $\lambda \leq \bigvee \{\sigma_j : j \in J_0\}$. Then $f(\lambda) \leq f(\bigvee \{\sigma_j : j \in J_0\}) \leq \bigvee \{f(\sigma_j) : j \in J_0\} \leq \bigvee \{\eta_j : j \in J_0\}$. Thus $f(\beta) \rightarrow f(\mu)$.

(iii) \iff (iv) and (v) \iff (vi). Immediate from the fact that for each $\mu \in I^X$, $\mu = \bigvee \{x_\alpha : x_\alpha \leq \mu\}$ and $f(\mu) = \bigvee \{f(x_\alpha) : x_\alpha \leq \mu\}$.

(iv) \iff (vii) and (vi) \iff (viii): Follow immediately from the definition.

(iii) \implies (vi): Obvious.

(vi) \implies (iii): Let β be a fuzzy filter base on X with $\beta \rightarrow \mu$ and let $\mathcal{U} = \{\eta_j : j \in J\}$ be an open Q_α -cover of $f(\mu)$. Put $\mathcal{F} = \langle \beta \rangle$. Then \mathcal{F} is a fuzzy filter on X with $\mathcal{F} \rightarrow \mu$. By Corollary 3.20, for each $\mathcal{P} \in \mathcal{P}(\mathcal{F})$ we have $\mathcal{P} \rightarrow \mu$ and so by (vi), $f(\mathcal{P}) \rightarrow f(\mu)$. Then there exist a finite subset J_0 of J and a $\lambda \in \mathcal{P}$ such that $f(\lambda) \leq \bigvee \{\eta_j : j \in J_0\}$. Since $\lambda \in \mathcal{P}$ for each $\mathcal{P} \in \mathcal{P}(\mathcal{F})$, we have $\lambda \in \mathcal{F}$ and so $f(\lambda) \in f(\mathcal{F})$. Thus $f(\mathcal{F}) \rightarrow f(\mu)$.

(iv) \implies (i): Let $x_\alpha \in \text{FP}(X)$ and $\eta \in N_{f(x_\alpha)}^Q$. Since $\beta = N_{x_\alpha}^Q \rightarrow x_\alpha$, we have $f(\beta) \rightarrow f(x_\alpha)$ by (iv) and so $(\exists \varrho \in \beta)(f(\varrho) \leq \eta)$. Hence f is F -continuous. \square

If (X, τ) and (Y, Δ) are topological spaces and $f: (X, \tau) \rightarrow (Y, \Delta)$, then f is continuous iff for any filter \mathcal{F} on X , we have $f(\text{adh}(\mathcal{B})) \subseteq \text{adh}(f(\mathcal{F}))$. This property is lost in the fuzzy case as the following theorem shows.

Theorem 3.23. *If a mapping $f: (X, \tau) \rightarrow (Y, \Delta)$ is F -continuous, then for every $\beta \in \text{FLB}(I^X)$ we have $f(\text{adh}(\beta)) \leq \text{adh}(f(\beta))$ ($f(Q\text{-adh}(\beta)) \leq Q\text{-adh}(f(\beta))$).*

Proof. Let β be a fuzzy filter base on X , $x_\alpha \leq Q\text{-adh}(\beta)$, $\varrho \in \beta$ and $\eta \in N_{f(x_\alpha)}^Q$. By hypothesis, there exists $\mu \in N_{x_\alpha}^Q$ such that $f(\mu) \leq \eta$. Since $\mu q \varrho$, we have $\eta q f(\varrho)$. So $f(x_\alpha) \leq Q\text{-adh}(f(\beta))$. This shows that $f(Q\text{-adh}(\beta)) \leq Q\text{-adh}(f(\beta))$. The proof for the part in front of the parentheses is similar. \square

Definition 3.24 [19]. Let $\{(X_j, \tau_j) : j \in J\}$ be a family of fuzzy topological spaces, $X = \prod_{j \in J} X_j$ and let τ be the fuzzy topology on X generated by the subbase $\{P_j^{-1}(\eta_j) : \eta_j \in \tau_j, j \in J\}$, where $P_j: X \rightarrow X_j$ is the projection mapping for each $j \in J$. The pair (X, τ) is called the fuzzy product space of the family $\{(X_j, \tau_j) : j \in J\}$.

Theorem 3.25. Let (X, τ) be the fuzzy product space of a family $\{(X_j, \tau_j) : j \in J\}$ of fuzzy topological spaces, β a fuzzy filter base on X , $\mu \in I^X$, $x_\alpha \in \text{FP}(X)$ and let $P_j : X \rightarrow X_j$ be the projection map. Then:

- (i) $\beta \rightarrow x_\alpha$ iff $P_j(\beta) \rightarrow P_j(x_\alpha)$ for each $j \in J$.
- (ii) $\beta \rightarrow \mu$ iff $P_j(\beta) \rightarrow P_j(\mu)$ for each $j \in J$.
- (iii) If $\beta \propto x_\alpha$, then $P_j(\beta) \propto P_j(x_\alpha)$ for each $j \in J$.
- (iv) If β is upper and $\beta \overset{Q}{\propto} x_\alpha$, then $P_j(\beta) \overset{Q}{\propto} P_j(x_\alpha)$ for each $j \in J$.

Proof. (i) Since each P_j is F -continuous, by Theorem 3.22 we have that $\beta \rightarrow x_\alpha$ implies that $p_j(\mathcal{B}) \rightarrow P_j(x_\alpha)$ for each $j \in J$. Conversely, suppose that $P_j(\beta) \rightarrow P_j(x_\alpha)$ for each $j \in J$. Let $\mu \in N_{x_\alpha}^Q$ in (X, τ) . Then there exist j_1, j_2, \dots, j_n in J and $\mu_n \in N_{x_\alpha}^Q$ in (X_{j_n}, τ_{j_n}) such that $\varrho = \bigwedge_{k=1}^n P_{j_k}^{-1}(\mu_k) \leq \mu$ and $x_\alpha q \varrho$. Since each $P_j(\beta) \rightarrow P_j(x_\alpha)$, $\mu_k \in P_{j_k}(\beta)$. Let $\varrho_k \in \mathcal{B}$ be such that $P_{j_k}(\varrho_k) = \mu_k$. Then $\varrho_1 \wedge \varrho_2 \dots \wedge \varrho_n \leq \varrho \leq \mu$ and hence $\mu \in \beta$. This proves that $\beta \rightarrow x_\alpha$.

(ii) Since each P_j is F -continuous, by Theorem 3.22 we have that $\beta \rightarrow \mu$ implies $P_j(\beta) \rightarrow P_j(\mu)$. Conversely, suppose that $P_j(\beta) \rightarrow P_j(\mu)$ and let $\mathcal{U} = \{\eta_j : j \in J\}$ be an open Q_α -cover of μ . Then $(\forall x \in X \text{ with } \mu(x) \geq \alpha)(\exists j_0 \in J)(x_\alpha q \eta_{j_0})$. Since $\eta_{j_0} \in \tau$, there exist j_1, j_2, \dots, j_n in J and $\sigma_i \in \tau_{j_i}$ such that $\sigma_0 = \bigwedge_{i=1}^n P_{j_i}^{-1}(\sigma_i) \leq \eta_{j_0}$ and $\sigma_0 q x_\alpha$. For each i we have $P_{j_i}(x_\alpha) q \sigma_i$ and thus there exists $\lambda_i \in \beta$ with $P_{j_i}(\lambda_i) \leq \sigma_i$. Let $\lambda \in \beta$ with $\lambda \leq \bigwedge_{i=1}^n \lambda_i$. Since $\lambda_i \leq P_{j_i}^{-1}(\sigma_i)$, we have $\lambda \leq \sigma_0 \leq \eta_{j_0}$. Thus $\lambda \leq \bigvee \{\eta_{j_i} : i = 0, 1, 2, \dots, n\}$ and this proves that $\beta \rightarrow \mu$.

(iii) follows from (i) and Theorem 3.13.

(iv) follows from (i) and Theorem 3.14. □

The concept of Q_α -compactness (strong Q -compactness) due to Zhongfu [33] provides the theory of fuzzy compactness which is applicable to arbitrary fuzzy subsets. So, Kerre, Noun and Kandil in [16], [17] introduced and studied the concept of $\varphi_{1,2}$ - F - Q_α -compactness to unify and generalize several characterizations and properties of Q_α -compactness due to Zhonghu [33] and their weaker and stronger forms. In this section we give new characterizations and properties of strong Q -compactness in terms of convergence of the fuzzy filter base and the fuzzy filter.

Definition 3.26 [33]. Let (X, τ) be a fts, $\alpha \in (0, 1]$ and $\mu \in I^X$. Then:

- (i) The family $\mathcal{U} = \{\eta_j : j \in J\} \subseteq \tau$ is called an open Q_α -cover of μ iff $(\forall x \in X \text{ with } \mu(x) \geq \alpha)(\exists j \in J)(x_\alpha q \eta_j)$.
- (ii) A subfamily \mathcal{U}_0 of a Q_α -cover \mathcal{U} of μ , which is also a Q_α -cover of μ , is called a Q_α -subcover of μ .
- (iii) A fuzzy set μ is called Q_α -compact if each open Q_α -cover of μ has a finite Q_α -subcover of μ .

- (iii) A fuzzy set $\mu \in I^X$ of a fts (X, τ) is said to be strong Q -compact if μ is Q_α -compact for each $\alpha \in (0, 1]$.
- (iv) A fts (X, τ) is called Q_α -compact (strong Q -compact) iff X is Q_α -compact (strong Q -compact).

Theorem 3.27. *Let (X, τ) be a fts and $\mu \in I^X$. Then μ is strong Q -compact iff for each family $\{\mu_j: j \in J\} \subseteq \tau'$ such that $\bigwedge_{j \in J} \mu_j(x) < \alpha$ for each $x \in \mu_{w\alpha}$, there exists a finite subset J_0 of J such that $\bigwedge_{j \in J_0} \mu_j(x) < \alpha$ for each $x \in \mu_{w\alpha}$.*

Proof. Let $\{\mu_j: j \in J\} \subseteq \tau'$ be such that $\bigwedge_{j \in J} \mu_j(x) < \alpha$ for each $x \in \mu_{w\alpha}$. Then $(\forall x \in X, \mu(x) \geq \alpha) \left(x_\alpha q \bigvee_{j \in J} \mu_j^c \right)$ and so the family $\{\mu_j^c: j \in J\}$ is an open Q_α -cover of μ . Since μ is strong Q -compact, there exists a finite subset J_0 of J such that the family $\{\mu_j^c: j \in J_0\}$ is a Q_α -subcover of μ . Then $(\forall x \in X, \mu(x) \geq \alpha) \left(x_\alpha q \bigvee_{j \in J_0} \mu_j^c \right)$. Thus $\bigwedge_{j \in J_0} \mu_j(x) < \alpha$ for each $x \in \mu_{w\alpha}$. Conversely, let $\{\eta_j: j \in J\}$ be an open Q_α -cover of μ . Then $(\forall x \in X, \mu(x) \geq \alpha) (\exists j \in J) (x_\alpha q \eta_j)$ and so $\bigwedge_{j \in J} \eta_j^c(x) < \alpha$ for each $x \in \mu_{w\alpha}$. Since $\eta_j^c \in \tau'$, by hypothesis there exists a finite subset J_0 of J such that $\bigwedge_{j \in J_0} \eta_j^c(x) < \alpha$ for each $x \in \mu_{w\alpha}$. It follows that $\{\eta_j: j \in J_0\}$ is a Q_α -subcover of μ . Thus μ is Q_α -compact for each $\alpha \in (0, 1]$. Thus μ is strong Q -compact. \square

Corollary 3.28. *A fts (X, τ) is strong Q -compact iff for each family $\{\mu_j: j \in J\} \subseteq \tau'$ such that $\bigwedge_{j \in J} \mu_j(x) < \alpha$ for each $x \in X$ there exists a finite subset J_0 of J such that $\bigwedge_{j \in J_0} \mu_j(x) < \alpha$ for each $x \in X$.*

Theorem 3.29. *Let (X, τ) be a fts and $\mu \in I^X$. If μ is strong Q -compact, then every upper fuzzy filter base (fuzzy filter base) β on μ is Q -adherent (adherent) to some x_α in μ .*

Proof. The proofs for both parts are very similar; so, we only present the proof for the part not in the parentheses. Let $\mu \in I^X$ be strong Q -compact and suppose that $\beta = \{\eta_j: j \in J\}$ is an upper fuzzy filter base on μ having no Q -adherent point in μ . Let $x_\alpha \leq \mu$. Corresponding to each $j \in J$ there exists $\varrho_{x_\alpha} \in N_{x_\alpha}^Q$ and $\eta_j(x_\alpha) \in \beta$ such that $\varrho_{x_\alpha} \bar{q} \eta_j(x_\alpha)$. Thus, the family $\mathcal{U} = \{\varrho_{x_\alpha}: x \in X, \mu(x) \geq \alpha\}$ is an open Q_α -cover of μ . Since μ is strong Q -compact, for each $\alpha \in (0, 1]$ there exist finitely many members $\varrho_{x_\alpha^1}, \varrho_{x_\alpha^2}, \dots, \varrho_{x_\alpha^n}$ from \mathcal{U} for $x_\alpha^1, x_\alpha^2, \dots, x_\alpha^n \leq \mu$ such that the family $\mathcal{U}_0 = \{\varrho_{x_\alpha^k}: k = 1, 2, \dots, n\}$ is a Q_α -subcover of μ . Since β is a fuzzy

filter base on μ , there exists $\eta_{j_0} \in \beta$ such that $\eta_{j_0} \leq \bigwedge_{k=1}^n \eta_j(x_\alpha^k)$ and $\eta_{j_0} \wedge \mu \neq \emptyset$ and so there exists $y_\nu \leq \eta_{j_0} \wedge \mu$. Since the family $\{\varrho_{x_\alpha^k} : k = 1, 2, \dots, n\}$ is a Q_α -cover of μ , we have $y_\nu q \varrho_{x_\alpha^k}$ for some $1 \leq k \leq n$ and so $\eta_{j_0} q \varrho_{x_\alpha^k}$ which implies $\varrho_{x_\alpha^k} q \eta_j(x_\alpha^k)$, a contradiction. \square

Corollary 3.30. *If a fts (X, τ) is strong Q -compact, then every upper fuzzy filter base (fuzzy filter base) β on X is Q -adherent (adherent) to some x_α .*

Theorem 3.31. *Let (X, τ) be a fts and $\mu \in I^X$. If μ is strong Q -compact, then every maximal upper fuzzy filter base (maximal fuzzy filter base) β on μ converges to some x_α in μ .*

Proof. The proofs for both parts are very similar; so, we only present the proof for the part not in the parentheses. Let β be a maximal upper fuzzy filter base on μ . By Theorem 3.29, we have $\beta \overset{Q}{\propto} x_\alpha$ in μ and then by Theorem 3.18 (iii), we have $\beta \rightarrow x_\alpha$ in μ . \square

Corollary 3.32. *If a fts (X, τ) is strong Q -compact, then every maximal upper fuzzy filter base (maximal fuzzy filter base) β on X converges to some x_α .*

As an application of the above results we can give a proof of the fuzzy Tychonoff Theorem (Theorem 3.16 in [17]) without use of the fuzzy Alexander's subbase theorem (Theorem 3.12 in [17]).

Theorem 3.33. *Let $\{(X_j, \tau_j) : j \in J\}$ be a family of fts's, let $(X, \tau) = \left(\prod_{j \in J} X_j, \prod_{j \in J} \tau_j\right)$ be a fuzzy product space and $\mu = \prod_{j \in J} \mu_j \in I^X$. Then μ_j is a strong Q -compact fuzzy set in (X_j, τ_j) for each $j \in J$ iff μ is a strong Q -compact fuzzy set in (X, τ) .*

Proof. Let $P_j : X \rightarrow X_j$ be the projection mapping and $\mu \in I^X$ a strong Q -compact fuzzy set in (X, τ) . Since P_j is F -continuous, hence by Theorem 3.8 in [16], $P_j(\mu) = \mu_j \in I^{X_j}$ is a strong Q -compact fuzzy set in (X_j, τ_j) for each $j \in J$. Conversely, suppose that each $\mu_j \in I^{X_j}$ is strong Q -compact and let β be a fuzzy filter base on μ . Then $p_j(\beta) = \beta_j$ is a fuzzy filter base on μ_j for each $j \in J$. By Theorem 3.27, $\beta_j \rightarrow x_\alpha^j$ and $x_\alpha^j \leq \mu_j$ for each $j \in J$. Then by Theorem 3.25 (i), $\beta \rightarrow x_\alpha = \langle x_\alpha^j : j \in J \rangle$ and $x_\alpha \leq \mu$. Hence μ is strong Q -compact. \square

4. NEW CONVERGENCE THEORY OF FUZZY NETS

In 1980, Pu and Liu introduced the notion of fuzzy nets and a new concept of the so-called Q-neighborhood was given, which could reflect the features of the neighborhood structure in fuzzy topological spaces. By this new neighborhood structure the Moore-Smith convergence theory was established splendidly [28].

In this section we will give a new notion of convergence for fuzzy nets different from the one given by Pu and Liu in [28]. For more details on the difference we refer to the results in Section 5, which are valid only for our new notions. Our notion is such that the relationship between fuzzy nets and fuzzy filters and their convergence is the same as the relation between filters and nets and their convergence in topological spaces.

Definition 4.1 [28]. A mapping $\mathcal{S}: D \rightarrow \text{FP}(X)$ is called a fuzzy net in X and is denoted by $\{S(n); n \in D\}$, where D is a directed set. If $S(n) = x_{\alpha_n}^n$ for each $n \in D$, where $x \in X$, $n \in D$ and $\alpha_n \in (0, 1]$ then the fuzzy net \mathcal{S} is denoted as $\{x_{\alpha_n}^n; n \in D\}$ or simply $\{x_{\alpha_n}^n\}$.

Definition 4.2 [28]. A fuzzy net $\mathcal{T} = \{y_{\alpha_m}^m; m \in E\}$ in X is called a fuzzy subnet of a fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ iff there is a mapping $f: E \rightarrow D$ such that:

- (i) $\mathcal{T} = \mathcal{S} \circ f$, that is, $T_i = \mathcal{S}_{f(i)}$ for each $i \in E$.
- (ii) For each $n \in D$ there exists some $m \in E$ such that, if $p \in E$ with $p \geq m$, then $f(p) \geq n$.

We shall denote a fuzzy subnet of a fuzzy net $\{x_{\alpha_n}^n; n \in D\}$ by $\{x_{\alpha_{f(m)}}^{f(m)}; m \in E\}$, the justification for which is clear from the definition.

Definition 4.3. Let (X, τ) be a fts and let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X and $\mu \in I^X$. Then \mathcal{S} is said to be :

- (i) in μ iff $(\forall n \in D)(x_{\alpha_n}^n \leq \mu)$.
- (ii) eventually with μ iff $(\exists m \in D)(\forall n \in D, n \geq m)(x_{\alpha_n}^n \leq \mu)$.
- (iii) frequently with μ iff $(\forall m \in D)(\exists n \in D, n \geq m)(x_{\alpha_n}^n \leq \mu)$.
- (iv) universal iff $\forall \eta, \varrho \in I^X$, if \mathcal{S} is eventually with $\eta \vee \varrho$, then it is eventually with η or ϱ .

Lemma 4.4. *Let (X, τ) be a fts and $\alpha \in [0, 1)$, then the net $\{x^n\}$ is a universal net in $(X, \iota_\alpha(\tau))$ iff the fuzzy net $\{x_{1-\alpha}^n\}$ is universal in (X, τ) .*

Proof. It is similar to that of Lemma 1.1 in [8] and hence omitted.

Definition 4.5. Let (X, τ) be a fts, $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ a fuzzy net in X and $x_\alpha \in \text{FP}(X)$. Then \mathcal{S} is said to be:

- (i) convergent to x_α , denoted by $\mathcal{S} \rightarrow x_\alpha$, if $(\forall \eta \in N_{x_\alpha}^Q)(\exists m \in D)(\forall n \in D \text{ with } n \geq m)(x_{\alpha_n}^n \leq \eta)$.
- (ii) Q -convergent to x_α [28], denoted by $\mathcal{S} \xrightarrow{Q} x_\alpha$, if $(\forall \eta \in N_{x_\alpha}^Q)(\exists m \in D)(\forall n \in D \text{ with } n \geq m)(x_{\alpha_n}^n q\eta)$.

The limit and Q -limit of the fuzzy net \mathcal{S} is defined by $\lim(S) = \bigvee \{x_\alpha \in \text{FP}(X) : S \rightarrow x_\alpha\}$ and $Q\text{-}\lim(S) = \bigvee \{x_\alpha \in \text{FP}(X) : \mathcal{S} \xrightarrow{Q} x_\alpha\}$, respectively.

Lemma 4.6. *Let (X, τ) be a fts and $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ a fuzzy net in X . Then*

- (i) $x_\alpha \leq \lim(\mathcal{S})$ ($x_\alpha \leq Q\text{-}\lim(\mathcal{S})$) iff $\mathcal{S} \rightarrow x_\alpha$ ($\mathcal{S} \xrightarrow{Q} x_\alpha$).
- (ii) $\lim(\mathcal{S})$ and $Q\text{-}\lim(\mathcal{S})$ are closed fuzzy sets.

Proof. (i) follows immediately from Definition 4.5.

(ii) Since $\lim(\mathcal{S}) \leq \text{cl}(\lim(\mathcal{S}))$ it suffices to show that $\text{cl}(\lim(\mathcal{S})) \leq \lim(\mathcal{S})$. Let $x_\alpha \leq \text{cl}(\lim(\mathcal{S}))$ and $\eta \in N_{x_\alpha}^Q$. Then $\eta q \lim(\mathcal{S})$ and so there exists $y \in X$ such that $\eta(y) + \lim(\mathcal{S})(y) > 1$. Put $t = \lim(\mathcal{S})(y)$. Then $y_t q \eta$ and $y_t \leq \lim(\mathcal{S})$ and so $(\forall \varrho \in N_{y_t}^Q)(\exists m \in D)(\forall n \in D, n \geq m)(x_{\alpha_n}^n \leq \varrho)$, which implies $x_{\alpha_n}^n \leq \eta$. Thus $x_\alpha \leq \lim(\mathcal{S})$. Hence $\text{cl}(\lim(\mathcal{S})) \leq \lim(\mathcal{S})$. Thus $\lim(\mathcal{S})$ is a closed fuzzy set. The proof for $Q\text{-}\lim(\mathcal{S})$ is similar. \square

Theorem 4.7. *Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and $\mu \in I^X$. Then $x_\alpha \leq \text{cl}(\mu)$ iff there exists a fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ convergent to x_α such that $x_{\alpha_n}^n q\mu$ for all n .*

Proof. Let $\mathcal{S} \rightarrow x_\alpha$ and $x_{\alpha_n}^n q\mu$ for all n . If $\eta \in N_{x_\alpha}^Q$, then $x_{\alpha_n}^n \leq \eta$ for some n and so $\eta q\mu$. Thus $x_\alpha \leq \text{cl}(\mu)$. Conversely, let $x_\alpha \leq \text{cl}(\mu)$. Then $(\forall \eta \in N_{x_\alpha}^Q)(\eta q\mu)$. That is, there exist $y \in \text{supp}(\eta)$ and a real number t_η with $0 < t_\eta \leq \eta(y)$ such that $y_{t_\eta}^\eta \leq \eta$ and $y_{t_\eta}^\eta q\mu$. Let $D = N_{x_\alpha}^Q$. Then (D, \leq) is directed under the inclusion relation. Then $\mathcal{S} = \{y_{t_\eta}^\eta \leq \eta : y_{t_\eta}^\eta q\mu, \eta \in D\}$ is a fuzzy net such that $\mathcal{S} \rightarrow x_\alpha$. \square

Lemma 4.8. *Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$. If $\mathcal{S} \rightarrow x_\alpha$, then every fuzzy subnet of \mathcal{S} is also convergent to x_α .*

Proof. Let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X such that $\mathcal{S} \rightarrow x_\alpha$ and let $T = \{y_{t_m}^m; m \in E\}$ be a fuzzy subnet of \mathcal{S} . Let $\eta \in N_{x_\alpha}^Q$. Then there exists $m \in D$ such that for every $n \in D, n \geq m$, we have $x_{\alpha_n}^n \leq \eta$. By definition of T , for this m there exists $l \in E$ such that for each $p \in E, p \geq l$, we have $f(p) \geq m$, where $f: E \rightarrow D$. Now, $y_{t_p}^p = x_{\alpha_{f(p)}}^{f(p)}$. Then $y_{t_p}^p \leq \eta$ for all $p \geq l$ and so $\mathcal{T} \rightarrow x_\alpha$. \square

Theorem 4.9. *The fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ is convergent to x_α iff every universal fuzzy subnet of \mathcal{S} is convergent to x_α .*

Proof. If $\mathcal{S} \rightarrow x_\alpha$, then by Lemma 4.8 every universal fuzzy subnet of \mathcal{S} is convergent to x_α . Conversely, assume that the condition is satisfied and (by way of contradiction) that \mathcal{S} is not convergent to x_α . Then $(\exists \rho \in N_{x_\alpha}^Q)(\forall m \in D)(\exists n \in D, n \geq m)(\rho(x^n) < x_{\alpha_n}^n(x^n))$. So we may assume that $\rho(x^n) < x_{\alpha_n}^n(x^n)$ for all $n \in D$. But then, although there are universal subnets of \mathcal{S} , no such subnet could converge to x_α . \square

Definition 4.10. Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X . Then x_α is called:

- (i) an adherent point of a fuzzy net \mathcal{S} , denoted by $\mathcal{S} \propto x_\alpha$, if $(\forall \eta \in N_{x_\alpha}^Q)(\forall n \in D)(\exists m \in D, m \geq n)(x_{\alpha_m}^m \leq \eta)$.
- (ii) a Q -adherent point of a fuzzy net \mathcal{S} [28], denoted by $\mathcal{S} \overset{Q}{\propto} x_\alpha$, if $(\forall \eta \in N_{x_\alpha}^Q)(\forall n \in D)(\exists m \in D, m \geq n)(x_{\alpha_m}^m \overset{Q}{\leq} \eta)$.

The adherence and Q -adherence of the fuzzy net \mathcal{S} is defined by $\text{adh}(\mathcal{S}) = \bigvee \{x_\alpha \in \text{FP}(X) : \mathcal{S} \propto x_\alpha\}$ and $Q\text{-adh}(\mathcal{S}) = \bigvee \{x_\alpha \in \text{FP}(X) : \mathcal{S} \overset{Q}{\propto} x_\alpha\}$, respectively.

We now prove the expected relationship between adherent (Q -adherent) points and fuzzy subnets.

Lemma 4.11. *Let (X, τ) be a fts and $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ a fuzzy net in X . Then:*

- (i) $x_\alpha \leq \text{adh}(\mathcal{S})$ ($x_\alpha \leq Q\text{-adh}(\mathcal{S})$) iff $\mathcal{S} \propto x_\alpha$ ($\mathcal{S} \overset{Q}{\propto} x_\alpha$).
- (ii) $\text{lim}(\mathcal{S}) \leq \text{adh}(\mathcal{S})$ and $Q\text{-lim}(\mathcal{S}) \leq Q\text{-adh}(\mathcal{S})$.
- (iii) $\text{adh}(\mathcal{S})$ and $Q\text{-adh}(\mathcal{S})$ are closed fuzzy sets.

Proof. It is similar to that of Lemma 4.6. \square

Theorem 4.12. *A fuzzy point x_α is an adherent (Q -adherent) point of a fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ iff there exists a fuzzy subnet \mathcal{T} of \mathcal{S} which converges (Q -converges) to x_α .*

Proof. The proofs for both parts are very similar; so, we only present the proof for the part not in the parentheses. Let $x_\alpha \leq \text{adh}(\mathcal{S})$. Then for any $\mu \in N_{x_\alpha}^Q$, there exists an element $x_{\alpha_n}^n$ of the fuzzy net \mathcal{S} such that $x_{\alpha_n}^n \leq \mu$. Let $E = \{(n, \mu) : n \in D, \mu \in N_{x_\alpha}^Q \text{ and } x_{\alpha_n}^n \leq \mu\}$. Then (E, \gg) is a directed set where $(m, \mu) \gg (n, \rho)$ iff $m \geq n$ in D and $\mu \leq \rho$ in $N_{x_\alpha}^Q$. Then $\mathcal{T} : E \rightarrow \text{FP}(X)$ given by $\mathcal{T}(m, \mu) = x_{\alpha_m}^m$ can be checked by a fuzzy subnet of \mathcal{S} . To show that $\mathcal{T} \rightarrow x_\alpha$, let $\eta \in N_{x_\alpha}^Q$. Then

there exists $n \in D$ such that $(n, \eta) \in E$ and then $x_{\alpha_n}^n \leq \eta$. Thus, for any $(m, \mu) \in E$ such that $(m, \mu) \gg (n, \eta)$, we have $\mathcal{T}(m, \mu) = x_{\alpha_m}^m \leq \mu \leq \eta$. Hence $\mathcal{T} \rightarrow x_\alpha$. The converse is clear. \square

In the following, we characterize fuzzy continuity between fuzzy topological spaces by means of convergent fuzzy nets.

Theorem 4.13. *Let $f: (X, \tau) \rightarrow (Y, \Delta)$ be a mapping. Then the following are equivalent:*

- (i) f is F -continuous.
- (ii) $\mathcal{S} \rightarrow x_\alpha$ implies $f(\mathcal{S}) \rightarrow f(x_\alpha)$ for each $x_\alpha \in \text{FP}(X)$.
- (iii) For each universal fuzzy net \mathcal{S} in X , $\mathcal{S} \rightarrow x_\alpha$ implies $f(\mathcal{S}) \rightarrow f(x_\alpha)$ for each $x_\alpha \in \text{FP}(X)$.
- (iv) For each fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ in X , $f(\lim(\mathcal{S})) \leq \lim(f(\mathcal{S}))$.
- (v) For each universal fuzzy net \mathcal{S} in X , $f(\lim(\mathcal{S})) \leq \lim(f(\mathcal{S}))$.
- (vi) For each fuzzy net \mathcal{S} in X , $f(\text{adh}(\mathcal{S})) \leq \text{adh}(f(\mathcal{S}))$.

Proof. (i) \implies (ii). Let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X such that $\mathcal{S} \rightarrow x_\alpha$. If $\eta \in N_{f(x_\alpha)}^Q$, then by (i), $f^{-1}(\eta) \in N_{x_\alpha}^Q$. Since $\mathcal{S} \rightarrow x_\alpha$, we have $(\exists m \in D)(\forall n \in D, n \geq m)(x_{\alpha_n}^n \leq f^{-1}(\eta))$ which implies that $f(x_{\alpha_n}^n) \leq \eta$. Thus $f(\mathcal{S}) \rightarrow f(x_\alpha)$.

(ii) \implies (i). Let $\eta \in I^Y$ be open and $x_\alpha \in \text{FP}(X)$ be such that $x_\alpha q f^{-1}(\eta)$. Then $f(x_\alpha) q \eta$ and so $\eta \in N_{f(x_\alpha)}^Q$. Suppose that for each $\mu \in N_{x_\alpha}^Q$, we have $\mu \not\leq f^{-1}(\eta)$. Then $\mu(x^\mu) > f^{-1}(\eta)(x^\mu)$ for some $x^\mu \in X$. Take $x_{\alpha(\mu)}^\mu$ to be a fuzzy point with $x_{\alpha(\mu)}^\mu(x^\mu) = \mu(x^\mu)$. Then $x_{\alpha(\mu)}^\mu \leq \mu$ and $\alpha(\mu) > f^{-1}(\eta)(x^\mu) = \eta(f(x^\mu))$. Let $D = (N_{x_\alpha}^Q, \leq)$. Then $\mathcal{S} = \{x_{\alpha(\mu)}^\mu; \mu \in N_{x_\alpha}^Q\}$ is a fuzzy net in X . It is easy to verify that \mathcal{S} converges to x_α but $f(\mathcal{S})$ does not converge to $f(x_\alpha)$, since $\eta \in N_{f(x_\alpha)}^Q$ and $f(x_{\alpha(\mu)}^\mu) \not\leq \eta$. This contradiction shows that there exists some $\mu \in N_{x_\alpha}^Q$ with $\mu \leq f^{-1}(\eta)$. Thus $f^{-1}(\eta)$ is an open Q -neighborhood of x_α and so $f^{-1}(\eta)$ is an open fuzzy set. Thus f is F -continuous.

(ii) \implies (vi): Let $x_\alpha \leq \text{adh}(\mathcal{S})$. By Theorem 4.11, there exists a fuzzy subnet T of \mathcal{S} with $T \rightarrow x_\alpha$. Then $f(T)$ is a fuzzy subnet of $f(\mathcal{S})$ and by (ii), $f(T) \rightarrow f(x_\alpha)$ which by Theorem 4.12 implies that $f(x_\alpha) \leq \text{adh}(f(\mathcal{S}))$. Thus $f(\text{adh}(\mathcal{S})) \leq \text{adh}(f(\mathcal{S}))$.

(vi) \implies (ii): Suppose that there exists a fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ with $\mathcal{S} \rightarrow x_\alpha$ while $f(\mathcal{S})$ does not converge to $f(x_\alpha)$. So $(\exists \eta \in N_{f(x_\alpha)}^Q)(\forall m \in D)(\exists n \in D, n \geq m)(f(x_{\alpha_n}^n) \not\leq \eta)$. So we may assume that $f(x_{\alpha_n}^n) \not\leq \eta$ for all $n \in D$. Since $x_\alpha \leq \lim(\mathcal{S}) \leq \text{adh}(\mathcal{S})$, we have $f(x_\alpha) \leq \text{adh}(f(\mathcal{S}))$ by (vi). Thus by Theorem 4.12, there exists a fuzzy subnet of $f(\mathcal{S})$ which is convergent to $f(x_\alpha)$, which is impossible since $f(x_{\alpha_n}^n) \not\leq \eta$ for all $n \in D$. Thus $f(\mathcal{S})$ must be convergent to $f(x_\alpha)$.

(iii) \implies (iv). It is obvious.

(iv) \implies (iii). Let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X and suppose that $\mathcal{S} \rightarrow x_\alpha$ while $f(\mathcal{S})$ does not converge to $f(x_\alpha)$. Then $(\exists \eta \in N_{f(x_\alpha)}^Q)(\forall m \in D)(\exists n \in D, n \geq m)(f(x_{\alpha_n}^n) \not\leq \eta)$. So we may assume that $f(x_{\alpha_n}^n) \not\leq \eta$ for all $n \in D$. If T is a universal fuzzy subnet of \mathcal{S} , then $T \rightarrow x_\alpha$ but $f(T)$ is not convergent to $f(x_\alpha)$, a contradiction with (iv). \square

Theorem 4.14. *Let $f: (X, \tau) \rightarrow (Y, \Delta)$ be a mapping. Then the following are equivalent:*

- (i) f is F -continuous.
- (ii) $\mathcal{S} \xrightarrow{Q} x_\alpha$ implies $f(\mathcal{S}) \xrightarrow{Q} f(x_\alpha)$ for each $x_\alpha \in FP(X)$.
- (iii) For each universal fuzzy net \mathcal{S} in X , $\mathcal{S} \xrightarrow{Q} x_\alpha$ implies $f(\mathcal{S}) \xrightarrow{Q} f(x_\alpha)$ for each $x_\alpha \in FP(X)$.
- (iv) For each fuzzy net $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ in X , $f(Q\text{-lim}(\mathcal{S})) \leq Q\text{-lim}(f(\mathcal{S}))$.
- (v) For each universal fuzzy net \mathcal{S} in X , $f(Q\text{-lim}(\mathcal{S})) \leq Q\text{-lim}(f(\mathcal{S}))$.
- (vi) For each fuzzy net \mathcal{S} in X , $f(Q\text{-adh}(\mathcal{S})) \leq Q\text{-adh}(f(\mathcal{S}))$.

Proof. We omit the proof which follows a line similar to that of Theorem 4.13 with certain straightforward modifications. \square

Theorem 4.15. *Let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in the fuzzy product space $(X, \tau) = \left(\prod_{j \in J} X_j, \prod_{j \in J} \tau_j\right)$, $x_\alpha \in FP(X)$ and let $P_j: X \rightarrow X_j$ be the projection map for each $j \in J$. Then:*

- (i) $\mathcal{S} \rightarrow x_\alpha$ iff $P_j(\mathcal{S}) = \{P_j(x_{\alpha_n}^n); n \in D\} \rightarrow P_j(x_\alpha)$ in X_j for each $j \in J$.
- (ii) If $\mathcal{S} \propto x_\alpha$, then $P_j(\mathcal{S}) = \{P_j(x_{\alpha_n}^n); n \in D\} \propto P_j(x_\alpha)$ in X_j for each $j \in J$.
- (iii) If $\mathcal{S} \overset{Q}{\propto} x_\alpha$, then $P_j(\mathcal{S}) = \{P_j(x_{\alpha_n}^n); n \in D\} \overset{Q}{\propto} P_j(x_\alpha)$ in X_j for each $j \in J$.

Proof. (i) Since P_j is F -continuous for each $j \in J$, then by Theorem 4.13, $\mathcal{S} \rightarrow x_\alpha$ implies $P_j(\mathcal{S}) \rightarrow P_j(x_\alpha)$ for each $j \in J$. Conversely, suppose that $P_j(\mathcal{S}) \rightarrow P_j(x_\alpha)$ for each $j \in J$. Let J_0 be a finite subset of J and $\beta_{x_\alpha}^Q = \left\{ \bigwedge_{j \in J_0} P_j^{-1}(\mu_j) : x_\alpha \wedge \bigwedge_{j \in J_0} P_j^{-1}(\mu_j), \mu_j \in \tau_j \right\}$. Then $\beta_{x_\alpha}^Q$ is a local base at x_α [28]. Let $\mu = \bigwedge_{j \in J_0} P_j^{-1}(\mu_j) \in \beta_{x_\alpha}^Q$. Then $(\forall j \in J)(\exists n_j \in D, j \in J_0)(\forall n \in D, n \geq n_j)(P_j(x_{\alpha_n}^n) \leq \mu_j)$. Let $n_0 \geq n_j$ for each $j \in J_0$. Then, if $n \geq n_0$, we have $P_j(x_{\alpha_n}^n) \leq \mu_j$ for each $j \in J$ and so $x_{\alpha_n}^n \leq \bigwedge_{j \in J_0} P_j^{-1}(\mu_j) = \mu$. Thus, $\mathcal{S} \rightarrow x_\alpha$.

(ii) and (iii) follow from (i) and Theorem 4.12. \square

In the following results we give new characterizations and properties of strong Q -compactness in terms of convergence of fuzzy nets.

Theorem 4.16. *Let (X, τ) be a fts, $\alpha \in (0, 1]$ and $\mu \in I^X$. Then μ is strong Q -compact iff each fuzzy net \mathcal{S} in μ has a Q -adherent point in μ .*

P r o o f. Let (X, τ) be a fts and let $\mu \in I^X$ be strong Q -compact and if possible, let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in μ without any Q -adherent point in μ . Then $(\forall x_\alpha \leq \mu)(\exists \eta_{x_\alpha} \in N_{x_\alpha}^Q)(\exists n_{x_\alpha} \in D)(\forall m \in D, m \geq n_{x_\alpha})(x_{\alpha_m}^m \bar{q}\eta_{x_\alpha})$. Then the family $\mathcal{U} = \{\eta_{x_\alpha} : x \in X, \mu(x) \geq \alpha\}$ is an open Q_α -cover of μ . Since μ is strong Q -compact, for each $\alpha \in (0, 1]$ there exist finitely many members $\eta_{x_\alpha^1}, \eta_{x_\alpha^2}, \dots, \eta_{x_\alpha^k}$ from \mathcal{U} for $x_\alpha^1, x_\alpha^2, \dots, x_\alpha^k \leq \mu$ such that the family $\mathcal{U}_0 = \{\eta_{x_\alpha^i} : i = 1, 2, \dots, k\}$ is a Q_α -subcover of μ . Let the corresponding elements in D be $n_{x_\alpha^1}, n_{x_\alpha^2}, \dots, n_{x_\alpha^k}$. Because D is a directed set, there exists $m \in D$ such that $m \geq n_{x_\alpha^i}$ for $i = 1, 2, \dots, k$. Then $x_{\alpha_m}^m q\eta_{x_\alpha^i}$ for some $1 \leq i \leq k$, a contradiction. Thus \mathcal{S} must have a Q -adherent point x_α (say) in μ . Conversely, let $\mathcal{A} = \{\mu_j : j \in J\} \subseteq \tau'$ such that for each finite subset J_0 of J we have $\bigwedge_{j \in J_0} \mu_j(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$. Let $D = \left\{ \bigwedge_{j \in J_0} \mu_j : \mu_j \in \mathcal{A}, j \in J \right\}$. Then $\mathcal{A} \subseteq D$. For each $\lambda_j \in D$, let us choose a fuzzy point $x_\alpha^{\lambda_j}$ and consider the fuzzy net $\mathcal{S} = \{x_\alpha^{\lambda_j} : \lambda_j \in D\}$ with the directed set (D, \geq) where for $\lambda_1, \lambda_2 \in D$, $\lambda_1 \geq \lambda_2$ iff $\lambda_1 \leq \lambda_2$ in X . By hypothesis, there exists $x_\alpha \leq \mu \wedge Q\text{-adh}(\mathcal{S})$. Let $\eta \in N_{x_\alpha}^Q$ and $\mu_j \in \mathcal{A}$. Since $\mu_j \in D$, there is $\lambda \in D$ with $\lambda \geq \mu_j$ (that is, $\lambda \leq \mu_j$) such that $\lambda q\eta$ and hence $\mu_j q\eta$. Thus $x_\alpha \leq \text{cl}(\mu_j) = \mu_j$ for each $j \in J$ and so $\bigwedge_{j \in J} \mu_j(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$, a contradiction. Consequently, by Theorem 3.27, we have that μ is strong Q -compact. \square

Corollary 4.17. *A fts (X, τ) is strong Q -compact iff each fuzzy net in X has a Q -adherent point.*

Theorem 4.18. *Let (X, τ) be a fts and $\mu \in I^X$. Then μ is strong Q -compact iff each fuzzy net in μ has a fuzzy subnet in μ Q -convergent to some fuzzy point in μ .*

P r o o f. Immediate consequence of Theorems 4.12 and 4.16. \square

Corollary 4.19. *A fts (X, τ) is strong Q -compact iff each fuzzy net has a fuzzy subnet convergent to some fuzzy point.*

Definition 4.20. Let (X, τ) be a fts, (D, \geq) a directed set. A fuzzy point x_α is called an adherent point of the fuzzy net $\mathcal{S} = \{\lambda_n; n \in D\}$ of fuzzy sets in X iff $(\forall n \in D)(\forall \eta \in N_{x_\alpha}^Q)(\exists m \in D, m \geq n)(\eta q\lambda_m)$.

Theorem 4.21. Let (X, τ) be a fts and $\mu \in I^X$. Then μ is strong Q -compact iff every fuzzy net of fuzzy sets in μ has an adherent point with value α in μ .

Proof. Let $\mathcal{S} = \{\mu_n; n \in D\}$ be a fuzzy net in μ (that is, $\mu_n \wedge \mu \neq \emptyset$ for each $n \in D$) of fuzzy sets in X and μ is strong Q -compact. For each $n \in D$, let $\lambda_n = \text{cl}(\bigvee\{\mu_m : m \in D, m \geq n\})$. Then $\mathcal{A} = \{\lambda_n : n \in D\}$ is a family of closed fuzzy sets with $\bigwedge_{n \in D_0} \lambda_n(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$ and every finite subset D_0 of D . By Theorem 3.27, we have $\bigwedge_{n \in D} \lambda_n(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$. Let $n \in D$ and $\eta \in N_{x_\alpha}^Q$. Then $\eta q \left(\bigvee_{m \geq n} \mu_m \right)$. Then $(\exists m \in D, m \geq n)(\eta q \mu_m)$. Thus $x_\alpha \leq \text{adh}(S)$. Conversely, let $\mathcal{A} = \{\mu_j : j \in J\}$ be a family of closed fuzzy sets with $\bigwedge_{j \in J_0} \mu_j(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$ and for each finite subset J_0 of J . Let $D = \left\{ \bigwedge_{j \in J} \mu_j : j \in J_0 \right\}$. Then (D, \leq) is a directed set, where \leq denotes the inclusion relation on D . For each $\lambda \in D$, we have $\lambda(x) \geq \alpha$ for some $x \in \mu_{w\alpha}$. Then the fuzzy net $\mathcal{S} = \{\lambda : \lambda \in D\}$ of fuzzy sets in μ has an adherent point x_α (say) in μ . Let $\mu_j \in \mathcal{A}$ and $\eta \in N_{x_\alpha}^Q$. Since $\mu_j \in D$, we have $(\exists \lambda \in D)(\lambda \leq \mu_j \text{ and } \lambda q \eta)$. Then $\mu_j q \eta$ and so $(\forall j \in J)(x_\alpha \leq \text{cl}(\mu_j) = \mu_j)$. Thus $\bigwedge_{j \in J} \mu_j(y) \geq \alpha$ for some $y \in \mu_{w\alpha}$. Consequently, by Theorem 3.27, we have that μ is strong Q -compact. \square

Corollary 4.22. A fts (X, τ) is strong Q -compact iff every fuzzy net of fuzzy sets in X has an adherent point.

5. THE RELATION BETWEEN FUZZY FILTER AND FUZZY NET CONVERGENCE

Like in convergence theory in general topology, we can associate with each fuzzy net in X a fuzzy filter on X and conversely. In this section we define the concepts of fuzzy filter base generated by a fuzzy net and fuzzy net based on a fuzzy filter. Then we study the relation between their convergence.

Definition 5.1. Let $\mathcal{S} = \{x_{\alpha_n}^n; n \in D\}$ be a fuzzy net in X . For each $n \in D$, let $\mu_n = \bigvee\{x_{\alpha_m}^m : m \in D, m \geq n\}$. Then the family $\beta_{\mathcal{S}} = \{\mu_n : n \in D\}$ forms a fuzzy filter base on X , called the fuzzy filter base generated by the fuzzy net \mathcal{S} .

Theorem 5.2. Let (X, τ) be a fts, $x_\alpha \in \text{FP}(X)$ and let \mathcal{S} be a fuzzy net in X . Then

- (i) $\mathcal{S} \rightarrow x_\alpha$ iff $\beta_{\mathcal{S}} \rightarrow x_\alpha$.
- (ii) $\mathcal{S} \propto x_\alpha$ iff $\beta_{\mathcal{S}} \propto x_\alpha$.

Proof. (i) Since $\beta_{\mathcal{S}} = \{\mu \in I^X : (\exists n \in D)(\mu = \bigvee\{x_{\alpha_m}^m : m \in D, m \geq n\})\}$, we have $\beta_{\mathcal{S}} \rightarrow x_{\alpha}$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\exists \mu \in \beta_{\mathcal{S}})(\mu \leq \eta)$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\exists n \in D)(\mu = \bigvee_{m \geq n} x_{\alpha_m}^m \leq \eta)$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\exists n \in D)(\forall m \in D, m \geq n)(x_{\alpha_m}^m \leq \eta)$ iff $\mathcal{S} \rightarrow x_{\alpha}$.

(ii) Since $\beta_{\mathcal{S}} \propto x_{\alpha}$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\forall \mu \in \beta_{\mathcal{S}})(\mu \wedge \eta \neq \emptyset)$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\forall n \in D)(\bigvee_{m \geq n} x_{\alpha_m}^m \wedge \eta \neq \emptyset)$ iff $(\forall \eta \in N_{x_{\alpha}}^Q)(\forall n \in D)(\exists m \in D, m \geq n)(x_{\alpha_m}^m \leq \eta)$ iff $\mathcal{S} \propto x_{\alpha}$. \square

Definition 5.3. Let β be a fuzzy filter base on X , $x_{\alpha} \in \text{FP}(X)$ and let $D_{\beta} = \{(\lambda, x_{\alpha}) : x_{\alpha} \leq \lambda \in \beta\}$. Let us define \geq in D_{β} by $(x_{\alpha}^1, \lambda_1) \geq (x_{\alpha}^2, \lambda_2)$ iff $\lambda_1 \leq \lambda_2$. Then (D_{β}, \geq) is a directed set. Define $\mathcal{S}_{\beta} : D_{\beta} \rightarrow \text{FP}(X)$ by $\mathcal{S}_{\beta}(x_{\alpha}, \lambda) = x_{\alpha}$. Then \mathcal{S}_{β} is a fuzzy net in X , called the fuzzy net based on the fuzzy filter base β .

Theorem 5.4. Let (X, τ) be a fts, $x_{\alpha} \in \text{FP}(X)$ and let β be a fuzzy filter base on X . Then:

- (i) $\beta \rightarrow x_{\alpha}$ iff $\mathcal{S}_{\beta} \rightarrow x_{\alpha}$.
- (ii) $\mathcal{S} \propto x_{\alpha}$ iff $\mathcal{S}_{\beta} \propto x_{\alpha}$.

Proof. (i) Let $\beta \rightarrow x_{\alpha}$ and let $\mathcal{S}_{\beta} : D_{\beta} \rightarrow \text{FP}(X)$ be the fuzzy net based on β , where $D_{\beta} = \{(x_{\alpha}, \lambda) : x_{\alpha} \leq \lambda \in \beta\}$ and $\mathcal{S}_{\beta}(x_{\alpha}, \lambda) = x_{\alpha}$. If $\mu \in N_{x_{\alpha}}^Q$, then there exists $\lambda \in \beta$ such that $\lambda \leq \mu$. Choose $y_t \leq \lambda$ such that $(y_t, \lambda) \in D_{\beta}$. If $(z_r, \lambda_1) \in D_{\beta}$ is such that $(z_r, \lambda_1) \geq (y_t, \lambda)$, then $\mathcal{S}_{\beta}(z_r, \lambda_1) = z_r \leq \lambda_1$. Since $\lambda_1 \leq \lambda \leq \mu$, we have $z_r \leq \mu$. Hence $\mathcal{S}_{\beta}(z_r, \lambda_1) \leq \mu$. Thus $\mathcal{S}_{\beta} \rightarrow x_{\alpha}$.

Conversely, let $\mathcal{S}_{\beta} \rightarrow x_{\alpha}$ and let $\mu \in N_{x_{\alpha}}^Q$. Then there is $(z_r, \lambda_1) \in D_{\beta}$ such that for each $(y_t, \lambda) \in D_{\beta}$ with $(y_t, \lambda) \geq (z_r, \lambda_1)$ we have $\mathcal{S}_{\beta}(y_t, \lambda) = y_t \leq \mu$. For each $w_{\nu} \leq \lambda_1$ we have $(w_{\nu}, \lambda_1) \geq (z_r, \lambda_1)$ and hence $\mathcal{S}_{\beta}(w_{\nu}, \lambda_1) = w_{\nu} \leq \mu$. Hence $\lambda_1 \leq \mu$. Thus $\beta \rightarrow x_{\alpha}$.

(ii) Let $\beta \propto x_{\alpha}$ and let $\mathcal{S}_{\beta} : D_{\beta} \rightarrow \text{FP}(X)$ be the fuzzy net based on β , where $D_{\beta} = \{(x_{\alpha}, \lambda) : x_{\alpha} \leq \lambda \in \beta\}$ and $\mathcal{S}_{\beta}(x_{\alpha}, \lambda) = x_{\alpha}$. Let $\mu \in N_{x_{\alpha}}^Q$ and $(y_t, \lambda) \in D_{\beta}$. Then $\lambda \wedge \mu \neq \emptyset$. If $z_r \leq \lambda \wedge \mu$, then $(z_r, \lambda) \in D_{\beta}$ is such that $(z_r, \lambda) \geq (y_t, \lambda)$. Hence $\mathcal{S}_{\beta}(z_r, \lambda) = z_r \leq \mu$. Thus $\mathcal{S}_{\beta} \propto x_{\alpha}$. Conversely, let $\mathcal{S}_{\beta} \propto x_{\alpha}$ and let $\mu \in N_{x_{\alpha}}^Q$ and $\lambda \in \beta$. If $y_t \leq \lambda$, then $(y_t, \lambda) \in D_{\beta}$ and hence there is $(z_r, \lambda_1) \in D_{\beta}$ with $(z_r, \lambda_1) \geq (y_t, \lambda)$, and we have $\mathcal{S}_{\beta}(z_r, \lambda_1) = z_r \leq \mu$. Since $\lambda_1 \leq \lambda$ and $z_r \leq \lambda_1 \wedge \mu$, we have $z_r \leq \lambda \wedge \mu$ and so $\lambda \wedge \mu \neq \emptyset$. Thus $\beta \propto x_{\alpha}$. \square

In the above Theorems 5.2 and 5.4, the concepts of convergence and adherence of a fuzzy net in our sense can not be replaced by Q -convergence and Q -adherence of a fuzzy net due to Pu and Liu [28].

References

- [1] *R. G. Bartle*: Nets and filters in topology. Amer. Math. Monthly *62* (1955), 551–557.
- [2] *R. Bělohávek*: Fuzzy Relational Systems: Foundations and Principles. Kluwer, New York, 2002.
- [3] *G. Birkhoff*: Moore-Smith convergence in general topology. Ann. Math. *38* (1937), 39–56.
- [4] *N. Bourbaki*: Topologie Generale, Ch. 1. Actualites Sci. Indust., Paris 858, 1940.
- [5] *H. Cartan*: Théorie des filtres. C. R. Acad. Sci. Paris *205* (1937), 595–598.
- [6] *C. L. Chang*: Fuzzy topological spaces. J. Math. Anal. Appl. *24* (1968), 182–190.
- [7] *Hu Cheng-Ming*: Theory of convergence in fuzzy topological spaces. J. Fuzzy Math. *1* (1993), 1–12.
- [8] *A. Choubey and A. K. Srivastava*: On α -compact fuzzy topological spaces. J. Fuzzy Math. *1* (1993), 321–326.
- [9] *P. Eklund and W. Gähler*: Fuzzy filter functors and convergence. Applications of Category Theory to Fuzzy Subsets. Kluwer Academic Publishers, Dordrecht-Boston-London, 1992, pp. 109–136.
- [10] *W. Gähler*: Convergence. Fuzzy Sets and Systems *73* (1995), 97–129.
- [11] *W. Gähler*: The general fuzzy filter approach to fuzzy topology. Part I. Fuzzy Sets and Systems *76* (1995), 205–224.
- [12] *W. Gähler*: The general fuzzy filter approach to fuzzy topology. Part II. Fuzzy Sets and Systems *76* (1995), 225–246.
- [13] *Wang Guojun*: A new fuzzy compactness defined by fuzzy nets. J. Math. Anal. Appl. *94* (1983), 1–23.
- [14] *A. Kandil, E. Kerre, A. Nouh and M. E. El-Shafei*: Generalized mappings between fuzzy topological spaces. Math. Pannon. *31* (1992), 59–71.
- [15] *J. L. Kelley*: General Topology. Van Nostrand, New York, 1955.
- [16] *E. E. Kerre, A. A. Nouh and A. Kandil*: Generalized compactness in fuzzy topological spaces. Math. Vesnik *43* (1991), 29–40.
- [17] *E. E. Kerre, A. A. Nouh and A. Kandil*: Operations on the class of all fuzzy sets on a universe endowed with a fuzzy topology. J. Math. Anal. Appl. *180* (1993), 325–341.
- [18] *G. J. Klir and B. Yuan*: Fuzzy Sets and Fuzzy Logic: Theory and Applications. Perentice Hall, 1995, pp. 574.
- [19] *R. Lowen*: Initial and final fuzzy topologies and the fuzzy Tychonoff theorem. J. Math. Anal. Appl. *58* (1977), 11–21.
- [20] *R. Lowen*: Convergence in fuzzy topological spaces. Gen. Topol. Appl. *10* (1979), 147–160.
- [21] *R. Lowen*: The relation between filter and net convergence in fuzzy topological spaces. Fuzzy Math. *3* (1983), 41–52.
- [22] *M. Macho Stadler and M. A. De Prada Vicente*: Fuzzy t -net theory. Fuzzy Sets and Systems *37* (1990), 225–235.
- [23] *M. Macho Stadler and M. A. De Prada Vicente*: On N -convergence of fuzzy nets. Fuzzy Sets and Systems *51* (1992), 203–217.
- [24] *M. Macho Stadler and M. A. De Prada Vicente*: t^* -fuzzy topological concepts. Portugaliae Math. *49* (1992), 85–108.
- [25] *E. H. Moore and H. L. Smith*: A general theory of limits. Amer. J. Math. *44* (1922), 102–121.
- [26] *M. A. De Prada Vicente and M. S. Aranguren*: Fuzzy filters. J. Math. Anal. Appl. *129* (1988), 560–568.
- [27] *M. A. De Prada Vicente and M. M. Stadler*: t -prefilter theory. Fuzzy Sets and Systems *38* (1990), 115–124.

- [28] *Pu Pao-Ming and Liu Ying-Ming*: Fuzzy topology I. *J. Math. Anal. Appl.* 76 (1980), 571–599.
- [29] *Pu Pao-Ming and Liu Ying-Ming*: Fuzzy topology II. *J. Math. Anal. Appl.* 77 (1980), 20–37.
- [30] *R. V. Sarma and N. Ajmal*: Fuzzy nets and their application. *Fuzzy Sets and Systems* 51 (1992), 41–51.
- [31] *Lee Bu Yong, Park Jin Han and Park Bae Hun*: Fuzzy convergence structures. *Fuzzy Sets and Systems* 56 (1993), 309–315.
- [32] *L. A. Zadeh*: Fuzzy sets. *Information and Control* 8 (1965), 338–353.
- [33] *Li Zhongfu*: Compactness in fuzzy topological spaces. *Kuxue Tongbao* 29 (1984), 582–585.

Author's address: Mathematics Department, Faculty of Science, Mansoura University, Mansoura 35516, Egypt, e-mail: aanouh@hotmail.com.