

Rade Doroslovački; Jovanka Pantović; Gradimir Vojvodić
One interval in the lattice of partial hyperclones

Czechoslovak Mathematical Journal, Vol. 55 (2005), No. 3, 719–724

Persistent URL: <http://dml.cz/dmlcz/128016>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ONE INTERVAL IN THE LATTICE OF PARTIAL HYPERCLONES

RADE DOROSLOVAČKI, JOVANKA PANTOVIĆ
and GRADIMIR VOJVODIĆ, Novi Sad

(Received November 3, 2002)

Abstract. In this paper the structure of the interval $[O_A, Hp_A]$ in the lattice of partial hyperclones is determined, where O_A is the clone of all total operations and Hp_A is the clone of all partial hyperoperations on A .

Keywords: clone, hyperoperation, hyperalgebra, hyperclone

MSC 2000: 08A40

1. PRELIMINARIES

Let A be a nonempty set. For a positive integer n , a function from A^n to the family $P(A)$ of all subsets of A is called a *partial n -hyperoperation on A* . Denote by $Hp_A^{(n)}$ the set of all partial n -hyperoperations on A and by Hp_A the set of all partial hyperoperations on A , i.e. $Hp_A = \bigcup_{n \geq 0} Hp_A^{(n)}$. A map f from A^n to $P(A) \setminus \{\emptyset\}$ is called a *hyperoperation* [5], and the set of all hyperoperations is denoted by H ($H \subseteq Hp_A$).

Every n -ary operation f from A^n to A can be viewed as a special partial hyperoperation (if we do not make difference between an element $a \in A$ and the corresponding one element subset $\{a\}$ of A). In the same sense, partial operations f from $\text{dom}(f)$ to A , where $\text{dom}(f) \subseteq A^n$, are also special partial hyperoperations (if $(x_1, \dots, x_n) \notin \text{dom}(f)$, for $(x_1, \dots, x_n) \in A^n$ and $f \in Hp_A^{(n)}$ we can put $f(x_1, \dots, x_n) = \emptyset$). Namely, $f \in Hp_A$ with $|f(x)| \leq 1$ for each $x \in A^n$ is de facto a partial operation on A (if A is a set, then $|A|$ is the cardinality of A). The set of all operations and the set of all partial operations on A are denoted by O_A and P_A , respectively.

For a positive integer n and for $1 \leq i \leq n$, e_i^n is a *partial n -hyperprojection* if $e_i^n(x_1, \dots, x_n) = \{x_i\}$ for all $x_1, \dots, x_n \in A$. The set of all hyperprojections is denoted by J_A .

For positive integers n and m , $f \in Hp_A^{(n)}$ and $g_1, \dots, g_n \in Hp_A^{(m)}$, the *composition of f and g_1, \dots, g_n* , denoted by $f(g_1, \dots, g_n) \in Hp_A^{(m)}$, is defined by $f(g_1, \dots, g_n)(x_1, \dots, x_m) = \bigcup \{f(y_1, \dots, y_n) : y_i \in g_i(x_1, \dots, x_m), 1 \leq i \leq n\}$ for each $(x_1, \dots, x_m) \in A^m$.

The set $C \subseteq Hp_A$ is a *clone of partial hyperoperations* on A or a *partial hyperclone* if C is composition closed and C contains all partial n -hyperprojections for each positive integer n .

For $F \subseteq Hp_A$, $\langle F \rangle$ stands for the clone of partial hyperoperations generated by F . The set F of partial hyperoperations is *complete* if $\langle F \rangle = Hp_A$.

A partial hyperclone C_1 on A is *covered by* a partial hyperclone C_2 if $C_1 \subset C \subset C_2$ holds for no partial hyperclone C . A maximal partial hyperclone on A is a partial hyperclone covered by Hp_A .

We say that an operation $f \in O_A^{(n)}$ *depends on* its i -th variable if there are $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in A$ such that $h \in O_A^{(1)}$, defined by $h(x) := f(a_1, \dots, a_{i-1}, x, a_{i+1}, \dots, a_n)$ for every $x \in A$, is non-constant. An n -ary operation $f \in O_A^{(n)}$ on A is *essential* if it depends on at least two variables and $\text{im } f = A$.

2. RESULTS

Let

$$\begin{aligned} H_A &= \bigcup_{n \in \mathbb{N}} \{f \in Hp_A^{(n)} : |f(\mathbf{x})| \geq 1 \text{ for every } \mathbf{x} \in A^n\}, \\ M &= \left\langle \bigcup_{n \in \mathbb{N}} \{f \in Hp_A^{(n)} : |f(\mathbf{x})| < 1 \text{ for every } \mathbf{x} \in A^n\} \right\rangle, \\ &= \bigcup_{n \in \mathbb{N}} \{f : A^n \rightarrow \{\emptyset\}\} \cup J_A, \\ O_A &= \bigcup_{n \in \mathbb{N}} \{f \in Hp_A^{(n)} : |f(\mathbf{x})| = 1 \text{ for every } \mathbf{x} \in A^n\} \end{aligned}$$

and

$$P_A = \bigcup_{n \in \mathbb{N}} \{f \in Hp_A^{(n)} : |f(\mathbf{x})| \leq 1 \text{ for every } \mathbf{x} \in A^n\}.$$

It is clear that these sets are clones of partial hyperoperations.

The next lemma follows from [3].

Lemma 2.1. *The clone O_A is covered by $O_A \cup M$ and the clone $O_A \cup M$ is covered by P_A .*

Lemma 2.2. *If $f \in H_A \setminus O_A$, then $\langle O_A \cup f \rangle = H_A$, i.e. the clone O_A is covered by H_A .*

Proof. If for an arbitrary $n \geq 1$, $f \in (H_A)^{(n)} \setminus O_A$, then there is at least one n -tuple $\mathbf{a} = (a_1, \dots, a_n) \in A^n$ such that $f(\mathbf{a}) = \{c_0, \dots, c_{p-1}\}$ and $p \geq 2$.

The statement $\langle O_A \cup \{f\} \rangle \subseteq H_A$ is obvious because H_A is a partial hyperclone and $f \in H_A$ and $O_A \subseteq H_A$. Now, we shall prove that the statement $\langle O_A \cup \{f\} \rangle \supseteq H_A$ is also correct. Let h be an arbitrary m -ary hyperoperation from H_A . Let us define maps $f_1, \dots, f_n \in O_A^{(m)}$ and $g \in O_A^{(l+m)}$ in the following way. If $h(y_1, \dots, y_m) = \{d_0, d_1, \dots, d_{q-1}\}$ for some $q \geq 1$, then $(f_1(y_1, \dots, y_m), \dots, f_n(y_1, \dots, y_m)) = (\{a_1\}, \dots, \{a_n\})$ and

$$\begin{aligned} g(y_1, \dots, y_m, c_0, \dots, c_0, c_0) &= \{d_0\} \\ g(y_1, \dots, y_m, c_0, \dots, c_0, c_1) &= \{d_1\} \\ &\vdots \\ g(y_1, \dots, y_m, c_{p-1}, \dots, c_{p-1}, c_{p-1}) &= \{d_{q-1}\} \end{aligned}$$

where $l \in \mathbb{N}$ is the number such that $p^{l-1} < \max_{(y_1, \dots, y_m) \in A^m} |h(y_1, \dots, y_m)| \leq p^l$.

Precisely, $g(y_1, \dots, y_m, c_{i_1}, \dots, c_{i_{l-1}}, c_{i_l}) = \{d_i\}$ where

$$i = \begin{cases} i_1 p^{l-1} + i_2 p^{l-2} + \dots + i_l p^0 & \text{if } i_1 p^{l-1} + i_2 p^{l-2} + \dots + i_l p^0 \leq q-1, \\ q-1 & \text{else.} \end{cases}$$

Now, we can prove that $h = g(e_1^m, \dots, e_m^m, f(f_1, \dots, f_n), \dots, f(f_1, \dots, f_n))$, which implies $h \in \langle O_A \cup f \rangle$. For $h(y_1, \dots, y_m) = \{d_0, d_1, \dots, d_{q-1}\}$, we have $g(e_1^m, \dots, e_m^m, f(f_1, \dots, f_n), \dots, f(f_1, \dots, f_n))(y_1, \dots, y_m) = g(\{y_1\}, \dots, \{y_m\}, \{c_0, \dots, c_{p-1}\}, \dots, \{c_0, \dots, c_{p-1}\}) = g(y_1, \dots, y_m, c_0, \dots, c_0) \cup g(y_1, \dots, y_m, c_0, \dots, c_0, c_1) \cup \dots \cup g(y_1, \dots, y_m, c_{p-1}, \dots, c_{p-1}, c_{p-1}) = \{d_0, d_1, \dots, d_{q-1}\} = h(y_1, \dots, y_m)$. \square

Lemma 2.3. *If $f \in M$, then $\langle H_A \cup \{f\} \rangle = H_A \cup M$, i.e. the clone H_A is covered by $H_A \cup M$.*

Proof. It is obvious that $\langle H_A \cup \{f\} \rangle \subseteq H_A \cup M$.

We only have to prove that for an arbitrary $h \in M$, $h \in \langle H_A \cup \{f\} \rangle$ holds. It is easy to see that $h = f(g_1, \dots, g_n)$ because $f(g_1, \dots, g_n)(y_1, \dots, y_m) = \emptyset$ for each $g_1, \dots, g_n \in H_A^{(m)}$. \square

Corollary 2.1. *If $f \in H_A \setminus O_A$, then $\langle O_A \cup M \cup \{f\} \rangle = H_A \cup M$, i.e. the clone $O_A \cup M$ is covered by $H_A \cup M$.*

Lemma 2.4. *If $f \in Hp_A \setminus (P_A \cup H_A)$, then $\langle O_A \cup \{f\} \rangle = Hp_A$.*

Proof. It is obvious that $\langle O_A \cup \{f\} \rangle \subseteq Hp_A$. It remains to prove that $Hp_A \subseteq \langle O_A \cup \{f\} \rangle$.

Since $f \in Hp_A \setminus (P_A \cup H_A)$, there is $\mathbf{a} = (a_1, \dots, a_n) \in A^n$ such that $f(\mathbf{a}) = \emptyset$ and $\mathbf{b} = (b_1, \dots, b_n) \in A^n$ such that $|f(\mathbf{b})| \geq 2$. We can suppose that $f(\mathbf{b}) = f(b_1, \dots, b_n) = \{c_0, \dots, c_{p-1}\}$ and $p \geq 2$.

Let h be an arbitrary map from $Hp_A^{(m)}$.

Let us define $f_1, \dots, f_n \in O_A^{(m)}$ and $e \in O_A^{(m+l)}$ as follows. If $h(\mathbf{y}) = \emptyset$, where $\mathbf{y} = (y_1, \dots, y_m) \in A^n$, then $(f_1(\mathbf{y}), \dots, f_n(\mathbf{y})) = (\{a_1\}, \dots, \{a_n\})$. If $h(\mathbf{y}) = \{d_0, d_1, \dots, d_{q-1}\}$ for some $q \geq 1$, then $(f_1(\mathbf{y}), \dots, f_n(\mathbf{y})) = (\{b_1\}, \dots, \{b_n\})$ and $e(y_1, \dots, y_m, c_{i_1}, \dots, c_{i_{l-1}}, c_{i_l}) = \{d_i\}$, where $l \in \mathbb{N}$ satisfies $p^{l-1} < \max_{(y_1, \dots, y_m) \in A^m} |h(\mathbf{y})| \leq p^l$ and

$$i = \begin{cases} i_1 p^{l-1} + i_2 p^{l-2} + \dots + i_l p^0 & \text{if } i_1 p^{l-1} + i_2 p^{l-2} + \dots + i_l p^0 \leq q - 1, \\ q - 1 & \text{else.} \end{cases}$$

Now, we can prove that $h = e(e_1^m, \dots, e_m^m, f(f_1, \dots, f_n), \dots, f(f_1, \dots, f_n))$, i.e. h belongs to $\langle O_A \cup \{f\} \rangle$.

For $h(y_1, \dots, y_m) = \emptyset$ the statement is obvious and for $h(y_1, \dots, y_m) = \{d_0, d_1, \dots, d_{q-1}\}$, for some $q \geq 1$, we have $e(e_1^m, \dots, e_m^m, f(f_1, \dots, f_n), \dots, f(f_1, \dots, f_n))(y_1, \dots, y_m) = e(\{y_1\}, \dots, \{y_m\}, f(\mathbf{a}), \dots, f(\mathbf{a})) = e(\{y_1\}, \dots, \{y_m\}, \{c_0, \dots, c_{p-1}\}, \dots, \{c_0, \dots, c_{p-1}\}) = e(y_1, \dots, y_m, c_0, \dots, c_0) \cup e(y_1, \dots, y_m, c_0, \dots, c_0, c_1) \cup \dots \cup e(y_1, \dots, y_m, c_{p-1}, \dots, c_{p-1}, c_{p-1}) = \{d_0, d_1, \dots, d_{q-1}\} = h(y_1, \dots, y_m)$. \square

Lemma 2.5. *If $f \in Hp_A \setminus (H_A \cup M)$, then $\langle H_A \cup M \cup \{f\} \rangle = Hp_A$, i.e. the clone $H_A \cup M$ is covered by Hp_A .*

Proof. If $f \in P_A$, the statement is a consequence of Lemma 2.1, or else (if $f \notin P_A$) of Lemma 2.4. \square

Lemma 2.6. *If $f \in Hp_A \setminus P_A$, then $\langle P_A \cup \{f\} \rangle = Hp_A$, i.e. P_A is covered by Hp_A .*

Proof. If $f \in H_A$, then the statement follows from Lemma 2.2. Otherwise, if $f \notin H_A$, then it follows from Lemma 2.4. \square

Theorem 2.1. *The structure of the interval*

$$[O_A, Hp_A] = \{O_A, H_A, O_A \cup M, H_A \cup M, P_A, Hp_A\}$$

is described in the following figure.




Figure. The interval $[O_A, Hp_A]$.

Theorem 2.2 (Slupecki-type criterion). *Let A be finite. If $F \subseteq Hp_A$ satisfies the following three conditions: (i) F contains an essential operation, (ii) F generates all unary operations and (iii) F contains the partial hyperoperation $f \in Hp_A \setminus (P_A \cup H_A)$, then F is complete.*

Proof. By Slupecki criterion, (i) and (ii) imply $O_A \subseteq \langle F \rangle$. From (iii) and Lemma 2.4 we obtain $Hp_A = \langle O_A \cup \{f\} \rangle \subseteq \langle F \rangle \subseteq Hp_A$, i.e. $\langle F \rangle = Hp_A$. \square

Corollary 2.2. *Let A be finite and $f \in Hp_A$. Then $\{f\}$ is complete iff $f \in Hp_A \setminus (P_A \cup H_A)$ and $\langle \{f\} \rangle$ contains all unary operations and at least one essential operation.*

References

- [1] *S. Burris and H. P. Sankappanavar*: A Course in Universal Algebra. Graduate Texts in Mathematics, Vol 78. Springer-Verlag, New York-Heidelberg-Berlin, 1981.
- [2] *Th. Drescher and R. Pöschel*: Multiclones and relations. Multi-Val. Logic 7 (2001), 313–337.
- [3] *L. Haddad, I. G. Rosenberg and D. Schweigert*: A maximal partial clone and a Slupecki-type criterion. Acta Sci. Math. 54 (1990), 89–98.
- [4] *R. Pöschel and L. A. Kalužnin*: Funktionen und Relationenalgebren. Ein Kapitel der diskreten Mathematik. Deutscher Verlag der Wiss., Berlin, 1979 (in German); Birkhäuser Verlag, Basel u. Stuttgart (Math. Reihe Bd. 67).
- [5] *I. G. Rosenberg*: An algebraic approach to hyperalgebras. Proceedings of 26th ISMVL, Santiago de Compostela, May 28–31, 1996. IEEE, 1996, pp. 203–207.

- [6] *I. G. Rosenberg*: Multiple-valued hyperstructures. Proceedings of 28th ISMVL, May 27–29, 1998. IEEE, 1998, pp. 326–333.

Authors' addresses: R. Doroslovački, J. Pantović, Faculty of Engineering, University of Novi Sad, Trg Dositeja Obradovića, 21000 Novi Sad, Serbia and Montenegro, e-mails: ftn_dora@eunet.yu, pantovic@uns.ns.ac.yu; G. Vojvodić, Dept. of Mathematics and Informatics, University of Novi Sad, Trg Dositeja Obradovića, 21000 Novi Sad, Serbia and Montenegro, e-mail: vojvodic@unsim.ns.ac.yu.