

Anatolij Dvurečenskij
Holland's theorem for pseudo-effect algebras

Czechoslovak Mathematical Journal, Vol. 56 (2006), No. 1, 47–59

Persistent URL: <http://dml.cz/dmlcz/128053>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

HOLLAND'S THEOREM FOR PSEUDO-EFFECT ALGEBRAS

ANATOLIJ DVUREČENSKIJ, Bratislava

(Received May 6, 2003)

Abstract. We give two variations of the Holland representation theorem for ℓ -groups and of its generalization of Glass for directed interpolation po-groups as groups of automorphisms of a linearly ordered set or of an antilattice, respectively. We show that every pseudo-effect algebra with some kind of the Riesz decomposition property as well as any pseudo MV -algebra can be represented as a pseudo-effect algebra or as a pseudo MV -algebra of automorphisms of some antilattice or of some linearly ordered set.

Keywords: pseudo-effect algebra, pseudo MV -algebra, antilattice, prime ideal, automorphism, unital po-group, unital ℓ -group

MSC 2000: 06F20, 03G12, 03B50

1. INTRODUCTION

A fundamental result of Holland [9] says that every ℓ -group G is an ℓ -subgroup of the ℓ -group $A(\Omega)$, the set of all automorphisms of a linearly ordered set Ω . This result was extended to directed interpolation po-groups¹ by Glass [7, Thm. 54] showing that G is isomorphic to a po-subgroup of the po-group $A(\Omega)$, the set of all automorphisms of an antilattice Ω .

Recently, partial algebraic structures, called pseudo-effect algebras and pseudo MV -algebras (as total algebraic structures), were introduced in [4], [5] and in [6], respectively. They can serve as models of quantum structures as well as of non-commutative logic, [8]. Under some natural conditions, it was proved, [4], [5] and [1], that they are precisely the intervals in unital po-groups or in unital ℓ -groups. Using these properties, we give an analogue of the Holland theorem showing that such a

This paper has been supported by the grant 2/7193/20 SAV, Bratislava, Slovakia.

¹ A po-group G is an *interpolation group* if, for $g_1, g_2 \leq h_1, h_2$, there exists an element $s \in G$ such that $g_1, g_2 \leq s \leq h_1, h_2$, $g_1, g_2, h_1, h_2 \in G^+$.

pseudo-effect algebra is isomorphic to a pseudo-effect algebra of automorphisms of a \wedge -antilattice Ω . As a corollary, we show that every pseudo MV -algebra is isomorphic to a pseudo MV -algebra of automorphisms of a linearly ordered set Ω .

Such a representation is useful since it gives a visualization of some pseudo-effect algebras as a set of automorphisms.

The paper is organized as follows. Pseudo-effect algebras and pseudo MV -algebras are presented in Section 2. Ideals P and mainly prime ideals of a pseudo-effect algebra E , and their characterizations via \wedge -antilattice properties of cosets E/P are studied in Section 3. A connection among prime ideals and prime subgroups of unital po-groups is shown in Section 4. Finally, the main results, the Holland theorems for pseudo-effect algebras and pseudo MV -algebras, are presented in Section 5.

2. PSEUDO-EFFECT ALGEBRAS

A partial algebra $(E; +, 0, 1)$, where $+$ is a partial binary operation and 0 and 1 are constants, is called a *pseudo-effect algebra*, [4], [5], if, for all $a, b, c \in E$, the following holds

- (i) $a + b$ and $(a + b) + c$ exist if and only if $b + c$ and $a + (b + c)$ exist, and in this case $(a + b) + c = a + (b + c)$;
- (ii) there is exactly one $d \in E$ and exactly one $e \in E$ such that $a + d = e + a = 1$;
- (iii) if $a + b$ exists, there are elements $d, e \in E$ such that $a + b = d + a = b + e$;
- (iv) if $1 + a$ or $a + 1$ exists, then $a = 0$.

If we define $a \leq b$ if and only if there exists an element $c \in E$ such that $a + c = b$, then \leq is a partial ordering on E such that $0 \leq a \leq 1$ for any $a \in E$. It is possible to show that $a \leq b$ if and only if $b = a + c = d + a$ for some $c, d \in E$. We write $c = a / b$ and $d = b \setminus a$. Then

$$(b \setminus a) + a = a + (a / b) = b,$$

and we write $a^- = 1 \setminus a$ and $a^\sim = a / 1$ for any $a \in E$.

For basic properties of pseudo-effect algebras see [4], [5]. We recall that if $+$ is commutative, E is said to be an *effect algebra*, for properties of effect algebras see [3].

For example, if (G, u) is a unital (not necessary Abelian) po-group with a strong unit u (in fact it is sufficient to take a positive element u in G),² and

$$\Gamma(G, u) := \{g \in G : 0 \leq g \leq u\},$$

then $(\Gamma(G, u); +, 0, u)$ is a pseudo-effect algebra if we restrict the group addition $+$ to $\Gamma(G, u)$.

² We say that a positive element u of a po-group G is a *strong unit* if, for any $g \in G$, there is an integer $n \geq 1$ such that $g \leq nu$.

According to [4], we introduce for pseudo-effect algebras the following forms of the Riesz decomposition properties:

- (a) For $a, b \in E$, we write $a \mathbf{com} b$ to mean that for all $a_1 \leq a$ and $b_1 \leq b$, a_1 and b_1 commute.
- (b) We say that E fulfils the *Riesz interpolation property*, (RIP) for short, if for any $a_1, a_2, b_1, b_2 \in E$ such that $a_1, a_2 \leq b_1, b_2$ there is a $c \in E$ such that $a_1, a_2 \leq c \leq b_1, b_2$.
- (c) We say that E fulfils the *weak Riesz decomposition property*, (RDP₀) for short, if for any $a, b_1, b_2 \in E$ such that $a \leq b_1 + b_2$ there are $d_1, d_2 \in E$ such that $d_1 \leq b_1, d_2 \leq b_2$ and $a = d_1 + d_2$.
- (d) We say that E fulfils the *Riesz decomposition property*, (RDP) for short, if for any $a_1, a_2, b_1, b_2 \in E$ such that $a_1 + a_2 = b_1 + b_2$ there are $d_1, d_2, d_3, d_4 \in E$ such that $d_1 + d_2 = a_1, d_3 + d_4 = a_2, d_1 + d_3 = b_1, d_2 + d_4 = b_2$.
- (e) We say that E fulfils the *commutational Riesz decomposition property*, (RDP₁) for short, if for any $a_1, a_2, b_1, b_2 \in E$ such that $a_1 + a_2 = b_1 + b_2$ there are $d_1, d_2, d_3, d_4 \in E$ such that
 - (i) $d_1 + d_2 = a_1, d_3 + d_4 = a_2, d_1 + d_3 = b_1, d_2 + d_4 = b_2$, and
 - (ii) $d_2 \mathbf{com} d_3$.
- (f) We say that E fulfils the *strong Riesz decomposition property*, (RDP₂) for short, if for any $a_1, a_2, b_1, b_2 \in E$ such that $a_1 + a_2 = b_1 + b_2$ there are $d_1, d_2, d_3, d_4 \in E$ such that
 - (i) $d_1 + d_2 = a_1, d_3 + d_4 = a_2, d_1 + d_3 = b_1, d_2 + d_4 = b_2$, and
 - (ii) $d_2 \wedge d_3 = 0$.

We introduce analogous notions for po-groups. Let G be a po-group and for $a, b \in G^+$, we write $a \mathbf{com} b$ iff, for all $a_1, b_1 \in G^+$ such that $a_1 \leq a$ and $b_1 \leq b$, we have $a_1 + b_1 = b_1 + a_1$.

Let $(G; +, 0, \leq)$ be a directed po-group. According to [4], [5], we say that G fulfils (RIP), (RDP₀), (RDP), (RDP₁), and (RDP₂), respectively, if analogous properties as those for pseudo-effect algebras hold also for the positive cone G^+ of G .

A mapping $h: E \rightarrow F$, where E and F are pseudo-effect algebras, is said to be a *homomorphism* if (i) $h(0) = 0$ and $h(1) = 1$, and (ii) $h(a + b) = h(a) + h(b)$ whenever $a + b$ is defined in E . If h is injective and surjective such that also h^{-1} is a homomorphism, then h is said to be an *isomorphism*, and E and F are *isomorphic*. It is clear that a one-to-one homomorphism f from E onto F is an isomorphism iff $f(a) \leq f(b)$ implies $a \leq b$.

According to [6], a *pseudo MV-algebra* is an algebra $(M; \oplus, -, \sim, 0, 1)$ of type $(2, 1, 1, 0, 0)$ such that the following axioms hold for all $x, y, z \in M$ with an additional

binary operation \odot defined via

$$y \odot x = (x^- \oplus y^-)^\sim$$

$$(A1) \quad x \oplus (y \oplus z) = (x \oplus y) \oplus z;$$

$$(A2) \quad x \oplus 0 = 0 \oplus x = x;$$

$$(A3) \quad x \oplus 1 = 1 \oplus x = 1;$$

$$(A4) \quad 1^\sim = 0; 1^- = 0;$$

$$(A5) \quad (x^- \oplus y^-)^\sim = (x^\sim \oplus y^\sim)^-;$$

$$(A6) \quad x \oplus x^\sim \odot y = y \oplus y^\sim \odot x = x \odot y^- \oplus y = y \odot x^- \oplus x;^3$$

$$(A7) \quad x \odot (x^- \oplus y) = (x \oplus y^\sim) \odot y;$$

$$(A8) \quad (x^-)^\sim = x.$$

If we define $x \leq y$ iff $x^- \oplus y = 1$, then \leq is a partial order such that M is a distributive lattice with $x \vee y = x \oplus x^\sim \odot y$ and $x \wedge y = x \odot (x^- \oplus y)$. For basic properties of pseudo MV -algebras see [6].

If we define a partial binary operation $+$ on M via: $x + y$ is defined iff $x \leq y^-$, and in this case $x + y := x \oplus y$, then $(M; +, 0, 1)$ is a pseudo-effect algebra, and a pseudo-effect algebra E can be converted into a pseudo MV -algebra such that the $+$ derived from \oplus and the original $+$ coincide iff E satisfies (RDP₂) [5].

For example, if u is a strong unit of a (not necessarily Abelian) ℓ -group G ,

$$\Gamma(G, u) := [0, u]$$

and

$$x \oplus y := (x + y) \wedge u,$$

$$x^- := u - x,$$

$$x^\sim := -x + u,$$

$$x \odot y := (x - u + y) \vee 0,$$

then $(\Gamma(G, u); \oplus, ^-, ^\sim, 0, u)$ is a pseudo MV -algebra [6].

The basic representation theorem for pseudo effect-algebras is the following result [4], [5], and for pseudo MV -algebras see also [1].

³ \odot has a higher priority than \oplus .

Theorem 2.1. For a pseudo-effect algebra E fulfilling (RDP_1) , there is a unique (up to isomorphism of unital po-groups) unital po-group (G, u) fulfilling (RDP_1) such that $E \cong \Gamma(G, u)$.

If M is a pseudo MV-algebra, there is a unique (up to isomorphism of unital ℓ -groups) unital ℓ -group (G, u) such that $M \cong \Gamma(G, u)$.

3. IDEALS OF PSEUDO-EFFECT ALGEBRAS

A non-empty subset I of a pseudo-effect algebra E is said to be an *ideal* of E if (i) $x + y \in I$ whenever $x, y \in I$ and if $x + y$ is defined in E , and (ii) if $x \leq y$ for $x \in E$ and $y \in I$, then $x \in I$. Then E as well as $\{0\}$ are ideals of E . We denote by $\mathcal{I}(E)$ the set of all ideals of E .

Let $a \in E$, then by $I_0(a)$ we denote the ideal of E generated by a . If E satisfies (RDP_0) , then by [2, Prop. 3.1],

$$I_0(a) = \{x \in E : x = a_1 + \dots + a_n, a_i \leq a, i = 1, \dots, n, n \geq 1\}.$$

An ideal I of E is (i) *normal* if $a + I = I + a$,⁴ (ii) *maximal* if I is a proper subset of E and it is not included in any proper ideal of E as a proper subset, and (iii) *prime* if $I_0(a) \cap I_0(b) \subseteq I$ implies $a \in I$ or $b \in I$. We denote by $\mathcal{N}(E)$, $\mathcal{M}(E)$, and $\mathcal{P}(E)$ the set of all normal ideals, maximal ideals, and prime ideals, respectively, of E . Using the Zorn lemma, we see that $\mathcal{M}(E)$ is non-void. Under some conditions on E , [2], we can prove that $\mathcal{M}(E) \subseteq \mathcal{P}(E)$.

We recall that $\{0\}, E \in \mathcal{N}(E)$ and if f is a homomorphism from a pseudo-effect algebra E into another one F , then

$$\text{Ker}(f) := \{x \in E : f(x) = 0\}$$

is a normal ideal of E .

The following result was proved in [2, Prop. 3.5].

Proposition 3.1.

- (1) An ideal P of a pseudo-effect algebra E is prime if and only if, for all $I, J \in \mathcal{I}(E)$ with $I \cap J \subseteq P$, we have $I \subseteq P$ or $J \subseteq P$.
- (2) If P is prime, then $I \cap J = P$ implies $I = P$ or $J = P$. If E satisfies (RDP) , then an ideal P is prime if and only if, for all $I, J \in \mathcal{I}(E)$ with $I \cap J = P$, we have $I = P$ or $J = P$.

⁴ If A is a non-empty subset of E , then $a + A := \{a + x : x \in A \text{ and } a + x \text{ is defined in } E\}$. In a similar way we define $A + a$.

Let P be an ideal of a pseudo-effect algebra E . For $a, b \in E$, we write

$$a \sim_P b$$

iff there are two elements $e, f \in P$ such that $a \setminus e = b \setminus f$. We note that in Remark 3.6, we define another relation, symmetric to \sim_P , which coincides with \sim_P in the case of a normal ideal P .

Proposition 3.2. *Let E be a pseudo-effect algebra with (RDP). If P is an ideal of E , then \sim_P is an equivalence on E , and on $E/P = \{a/P : a \in P\}$, where $a/P := \{b \in E : b \sim_P a\}$, we can define a partial ordering $a/P \leq b/P$ if and only if there is an element $e \in P$ such that $a \setminus e \leq b$. If $a \wedge b$ is defined in E , then $(a \wedge b)/P = a/P \wedge b/P$.*

In addition, if P is a normal ideal, then E/P can be organized into a pseudo-effect algebra $(E/P; +, 0/P, 1/P)$, where the partial addition $+$ is defined by $a/P + b/P = c/P$ if and only if there are $a_1 \in a/P, b_1 \in b/P$ and $c_1 \in c/P$ such that $a_1 + b_1 = c_1$. Moreover, if P is a normal ideal of an E satisfying (RDP), or (RDP₁), or (RDP₂), then so satisfies E/P .

Proof. (i) \sim_P is an equivalence. It is clear that $a \sim_P a$, and $a \sim_P b$ implies $b \sim_P a$. Assume now $a \sim_P b$ and $b \sim_P c$. There are four elements $e, f, u, v \in P$ such that $a \setminus e = b \setminus f$ and $b \setminus u = c \setminus v$. Therefore, $b = (a \setminus e) + f = (c \setminus v) + u$. Due to (RDP), we can find $c_{11}, c_{12}, c_{21}, c_{22}$ in E such that $a \setminus e = c_{11} + c_{12}$, $c \setminus v = c_{11} + c_{21}$, $f = c_{21} + c_{22}$, and $u = c_{12} + c_{22}$. It is clear that $c_{12}, c_{21}, c_{22} \in P$. Hence, $a = c_{11} + c_{12} + e$ and $c = (b \setminus u) + v = (c_{11} + c_{12} + c_{21} + c_{22}) \setminus (c_{12} + c_{22}) + v = c_{11} + c_{21} + v$. Putting $s = c_{12} + e \in P$ and $t = c_{21} + v \in P$, we have $a \setminus s = c_{11} = c \setminus t$, i.e., $a \sim_P c$.

(ii) We show that \leq is a well-defined relation. Assume $a/P = a_1/P$ and $b/P = b_1/P$ and let $a \setminus e \leq b$ for some $e \in P$. There are $u, v, s, t \in P$ such that $a \setminus u = a_1 \setminus v$ and $b \setminus s = b_1 \setminus t$. Then $a = (a_1 \setminus v) + u$ and $b = (b_1 \setminus t) + s$, and there is an element $x \in E$ such that $(a \setminus e) + x = b$. Then $s = s_1 + s_2 + s_3$, where $s_1 \leq a_1 \setminus v$, $s_2 \leq u$, and $s_3 \leq x$. Hence

$$\begin{aligned} (a \setminus v) + u + x &= (b_1 \setminus t) + s, \\ ((a \setminus v) \setminus s_1) + s_1 + (u \setminus s_2) + s_2 + (x \setminus s_3) + s_3 &= (b_1 \setminus t) + s_1 + s_2 + s_3, \\ (a \setminus (s_1 + v)) + x_1 + x_2 + s_1 + s_2 + s_3 &= (b_1 \setminus t) + s_1 + s_2 + s_3, \end{aligned}$$

where $x_1, x_2 \in E$, which gives $(a \setminus (s_1 + v)) \leq b_1 \setminus t \leq b_1$.

(iii) We now show that \leq is a partial order on E/P . It is clear that $a/P \leq a/P$. Assume $a/P \leq b/P$ and $b/P \leq a/P$. There are two elements $a_1, b_1 \in P$ such that $a \setminus a_1 \leq b$ and $b \setminus b_1 \leq a$. Hence, there exists $x \in E$ such that $(a \setminus a_1) + x = b = (b \setminus b_1) + b_1$. Then $b_1 = b' + b''$, where $b' \leq a \setminus a_1$ and $b'' \leq x$, which gives

$((a \setminus a_1) \setminus b') + b' + (x \setminus b'') + b'' = (b \setminus b_1) + b' + b''$, i.e., $((a \setminus a_1) \setminus b') + x_1 + b' + b'' = (b \setminus b_1) + b' + b''$, where $x_1 \in E$. Hence, $a \setminus (b' + a_1) + x_1 = b \setminus b_1$, and there exists an element $y \in E$ such that

$$(*) \quad (a \setminus (b' + a_1)) + x_1 + y = (b \setminus b_1) + y = a = (a \setminus (b' + a_1)) + (b' + a_1),$$

which yields $x_1 + y = b' + a_1 \in P$, and, consequently, $x_1, y \in P$. Using $(*)$, we have $a \setminus (b' + a_1) = b \setminus (x_1 + b_1)$ which proves $a/P = b/P$.

Finally, assume $a/P \leq b/P$ and $b/P \leq c/P$. There are $a_1, b_1 \in P$ such that $a \setminus a_1 \leq b$ and $b \setminus b_1 \leq c$. Hence, $(a \setminus a_1) + x = b = (b \setminus b_1) + b_1$ for some $x \in E$. Then $b_1 = b' + b''$, where $b' \leq a \setminus a_1$ and $b'' \leq x$. Therefore, $((a \setminus a_1) \setminus b') + b' + (x \setminus b'') + b'' = (b \setminus b_1) + b' + b''$, i.e., $(a \setminus (b' + a_1)) + x_1 \leq b \setminus b_1 \leq c$ for some $x_1 \in E$, which implies $a \setminus (b' + a_1) \leq c$ and, consequently $a/P \leq c/P$.

(iv) It is clear that $(a \wedge b)/P \leq a/P, b/P$. Assume $x/P \leq a/P$ and $x/P \leq b/P$. There are $x_1, x_2 \in x/P$ such that $x_1 \leq a$ and $x_2 \leq b$. Since $x_1 \sim_P x_2$, there are $e, f \in P$ with $x_1 \setminus e = x_2 \setminus f$. Hence, for $x_0 = x_1 \setminus e$, we have $x_0 \in x/P$, and $x_0 \leq x_1, x_0 \leq x_2$. Consequently, $x_0 \leq a, b$ which yields $x_0 \leq a \wedge b$, i.e., $x/P = x_0/P \leq (a \wedge b)/P$.

If P is a normal ideal, the assertion was proved in [2, Prop. 4.1]. \square

We recall that a poset $(E; \leq)$ is (i) an *antilattice* if only comparable elements of E have an infimum or a supremum, (ii) a \wedge -*antilattice* if only comparable elements of E have an infimum. It is clear that any linearly ordered poset is an antilattice. Let E be a pseudo-effect algebra. Then E is an antilattice iff $a \wedge b = 0$ implies $a = 0$ or $b = 0$, while $(a \setminus (a \wedge b)) \wedge (b \setminus (a \wedge b)) = 0$, see [2].

Proposition 3.3. *Let P be an ideal of a pseudo-effect algebra E with (RDP) and let $a \leq b, a, b \in E$. Then $a/P = b/P$ if and only if $a = b \setminus s$ for some $s \in P$.*

Proof. One direction is clear. Assume $a/P = b/P$. There are $e, f \in P$ such that $a \setminus e = b \setminus f$. Then $a = (b \setminus f) + e \leq b = (b \setminus f) + f$ which entails $e \leq f$. Hence $a = (b \setminus f) + e = (b \setminus (e + (e \setminus f))) + e = (b \setminus (e \setminus f)) \setminus e + e = b \setminus (e \setminus f) = b \setminus s$, where $s = e \setminus f \in P$. \square

Proposition 3.4. *Let P be an ideal of a pseudo-effect algebra E with (RDP). Then E/P is a \wedge -antilattice if and only if $x/P \wedge y/P = 0/P$ implies $x/P = 0/P$ or $y/P = 0/P$.*

Proof. One direction is evident. Assume $x/P \wedge y/P = 0/P$ implies $x/P = 0/P$ or $y/P = 0/P$. Suppose $a/P \wedge b/P = c/P$. We claim there exists an element $c_0 \in c/P$ such that $c_0 \leq a, c_0 \leq b$ and $(c_0 \setminus a)/P \wedge (c_0 \setminus b)/P = 0/P$. Indeed, there

are $c_1, c_2 \in c/P$ such that $c_1 \leq a$ and $c_2 \leq b$. Since $c_1 \setminus e = c_2 \setminus f$, for $c_0 := c_1 \setminus e$, we have $c_0/P = a/P \wedge b/P$.

Assume now $x/P \leq (c_0 / a)/P$ and $x/P \leq (c_0 / b)/P$. There are two elements $x_1, x_2 \in x/P$ such that $x_1 \leq c_0 / a$ and $x_2 \leq c_0 / b$. Since $x_1 \sim_P x_2$, there are $e_1, f_1 \in P$ such that $x_0 := x_1 \setminus e_1 = x_2 \setminus f_1 \leq c_0 / a, c_0 / b$. Then $c_0 \leq c_0 + x_0 \leq a, b$, which proves $c_0/P \leq (c_0 + x_0)/P \leq c/P = c_0/P$, i.e., $c_0/P = (c_0 + x_0)/P$. By Proposition 3.3, there is an element $s \in P$ such that $c_0 = (c_0 + x_0) \setminus s$ which yields $c_0 + s = c_0 + x_0$, i.e., $x_0 = s \in P$, and consequently, $(c_0 / a)/P \wedge (c_0 / b)/P = 0/P$.

By the assumptions, $c_0 / a \in P$ or $c_0 / b \in P$. In the first case, there is $t \in P$ such that $c_0 / a = t$, i.e., $a = c_0 + t$ which by Proposition 3.3 gives $a/P = c_0/P = c/P$, i.e., E/P is an \wedge -antilattice. \square

Theorem 3.5. *An ideal P of a pseudo-effect algebra E with (RDP) is prime if and only if E/P is a \wedge -antilattice.*

Proof. Assume that P is prime and let $a/P \wedge b/P = 0/P$. We assert that $I_0(a) \cap I_0(b) \subseteq P$. Take $x \in I_0(a) \cap I_0(b)$. Then $x = a_1 + \dots + a_m = b_1 + \dots + b_n$, where $a_i \leq a$ and $b_j \leq b$ for all i and all j . (RDP) implies that there is a system $\{c_{ij}\}$ of elements of E such that $a_i = \sum_j c_{ij}$ and $b_j = \sum_i c_{ij}$. Since $c_{ij} \leq a, b$, we have $c_{ij}/P = 0/P$, i.e., $c_{ij} \in P$, which yields $a_i \in P$ and $x \in P$. Since P is prime, then $a \in P$ or $b \in P$, i.e., $a/P = 0/P$ or $b/P = 0/P$, which proves by Proposition 3.4 that E/P is a \wedge -antilattice.

Conversely, let E/P be a \wedge -antilattice and assume $I_0(a) \cap I_0(b) \subseteq P$. We assert $a/P \wedge b/P = 0/P$. Assume $x/P \leq a/P$ and $x/P \leq b/P$. As before, there exists an element $x_0 \in x/P$ such that $x_0 \leq a, b$. Hence, $x_0 \in I_0(a) \cap I_0(b) \subseteq P$ which proves $x_0 \in P$, and therefore, $x/P = x_0/P = 0/P$, which implies $a/P = 0/P$ or $b/P = 0/P$, i.e., $a \in P$ or $b \in P$. \square

Remark 3.6. Let P be an ideal of a pseudo-effect algebra E with (RDP). We define a new relation $_P \sim$ on E defined via $a \sim_P b$ iff there are two elements $e, f \in P$ such that $e / a = f / b$. In fact, \sim_P and $_P \sim$ induce two orderings. Then all previous results can be rewritten also for this relation. In addition, if P is normal, then both orderings induced by \sim_P and $_P \sim$ coincide.

4. PRIME AUBGROUPS OF PO-GROUPS

Let G be a directed po-group written additively, and let $\mathcal{C}(G)$ denote the set of all convex directed subgroups of G .

In analogy with pseudo-effect algebras, we say that a directed convex subgroup P of a po-group G is a *prime subgroup* of G if, for all directed convex subgroups I and J of G , $I \cap J \subseteq P$ implies $I \subseteq P$ or $J \subseteq P$. We denote by $\mathcal{P}(G)$ the set of all prime subgroups of a unital po-group G . An equivalent definition is (see [2, Prop. 6.2]): $C \in \mathcal{C}(G)$ is prime iff, for $a, b \in G$, $G_0(a) \cap G_0(b) \subseteq C$ implies $a \in P$ or $b \in P$.

Let $C \in \mathcal{C}(G)$ and define $x/C := \{y \in G: x - y \in C\}$, and $G/C := \{x/C: x \in G\}$. We order the set G/C with the usual order of left cosets of G/C via $x/C \leq y/C$ iff $x \leq y + c$ for some $c \in C$.

The following result has been proved in [7, Lemma 22].

Theorem 4.1. *A convex directed subgroup C of a directed po-group G with (RIP) is prime if and only if G/C is a \wedge -antilattice.*

If a pseudo-effect algebra $E = \Gamma(G, u)$ satisfies (RDP_1) , then there exists a one-to-one correspondence between the sets $\mathcal{I}(E)$ of all ideals or $\mathcal{P}(E)$ of all prime ideals of E and the sets $\mathcal{C}(G)$ and $\mathcal{P}(G)$, respectively, established in [2].

Theorem 4.2. *Let $E = \Gamma(G, u)$, where (G, u) is a unital po-group satisfying (RDP_1) . Let I be an ideal of E . Set*

$$\varphi(I) = \{x \in G: \exists x_i, y_j \in I, x = x_1 + \dots + x_n - y_1 - \dots - y_m\}.$$

Then $\varphi(I)$ is an o -ideal of (G, u) if and only if I is a normal ideal of E . In that case

$$(E/I, u/I) = \Gamma(G/\varphi(I), u/\varphi(I)).$$

In addition, if K is an o -ideal of (G, u) , then its restriction to E , denoted by $\psi(K)$, gives a normal ideal of E , i.e.,

$$\psi(K) := K \cap E \in \mathcal{I}(E), \quad K \in \mathcal{I}(G, u).$$

Moreover, both mappings, φ and ψ , are mutually bijective and preserve the set-theoretical inclusion.

Theorem 4.3. *Let $E = \Gamma(G, u)$, where (G, u) is a unital po-group satisfying (RDP₁). Let I be an ideal of E . Set*

$$\begin{aligned}\delta(I) &= \{x \in G: x = x_1 - y_1 + \dots + x_n - y_n, x_i, y_i \in I\}, \\ \delta_c(I) &= \{h \in G: h = x + p_1 = y - p_2, x, y \in \delta(I), p_1, p_2 \in G^+\}, \\ \delta_0(I) &= \{h_1 - h_2: h_1, h_2 \in \delta_c(I) \cap G^+\}.\end{aligned}$$

Then $\delta(I)$ and $\delta_c(I)$ is the subgroup and the convex subgroup, respectively, of G generated by I , and $\delta_0(I)$ is the largest directed convex subgroup of G that is contained in $\delta_c(I)$.

Let I and J be two ideals of E . Then $I \subseteq J$ if and only if $\delta(I) \subseteq \delta(J)$ if and only if $\delta_c(I) \subseteq \delta_c(J)$ if and only if $\delta_0(I) \subseteq \delta_0(J)$.

Let K be a convex subgroup of (G, u) . Then

$$\gamma(K) := K \cap E$$

is an ideal of E , and $\delta_c(\gamma(K)) \subseteq K$. If K is directed, then $\delta_0(\gamma(K)) = K$, and $\gamma(\delta_0(I)) = I$ for any ideal I of E . In addition, if K_1 and K_2 are two directed convex subgroups of (G, u) , then $\gamma(K_1) \subseteq \gamma(K_2)$ if and only if $K_1 \subseteq K_2$.

If K is a prime subgroup of (G, u) , then $\gamma(K) := K \cap E$ is a prime ideal of E , and if P is a prime ideal of E , then $\delta_0(P)$ is a prime subgroup of (G, u) . In addition, both mappings, γ and δ_0 , are mutually bijective and preserve the set-theoretical inclusion.

Moreover, the mappings γ and δ_0 restricted to normal prime ideals and prime o -ideals are mutually bijective.

We recall that if $a, b \in E$ and if I is an ideal of E , then $a/I \leq b/I$ iff $a/\delta_0(I) \leq b/\delta_0(I)$.

5. HOLLAND THEOREM AND PSEUDO-EFFECT ALGEBRAS

Let (Ω, \leq) be a nonvoid \wedge -antilattice, and let $A(\Omega)$ be the set of all automorphisms $\alpha: \Omega \rightarrow \Omega$ which preserve the partial order \leq . Then $A(\Omega)$ can be converted into a po-group such that the group-addition is the composition of automorphisms, the order on $A(\Omega)$ is defined via $\alpha \leq \beta$ iff $(\omega)\alpha \leq (\omega)\beta$ for all $\omega \in \Omega$, and the neutral element is the identity function on Ω . If α is a positive element from $A(\Omega)$, then $\Gamma(G, \alpha)$ is a pseudo-effect algebra of automorphisms of an \wedge -antilattice set Ω .

Holland [9] proved the basic result that every ℓ -group can be injectively embedded into the ℓ -group $A(\Omega)$ for some linearly ordered set Ω , and Glass [7, Thm. 54] generalized this result to directed po-groups satisfying (RIP) showing that every such a po-group can be embedded into the po-group $A(\Omega)$ for some antilattice Ω .

We show that a similar result can be proved also for pseudo-effect algebras by proving that every pseudo-effect algebra E satisfying (RDP_1) can be embedded into some $\Gamma(A(\Omega), \alpha)$.

Theorem 5.1. *Every pseudo-effect algebra E with (RDP_1) can be represented as a pseudo-effect algebra of automorphisms from $A(\Omega)$ for some \wedge -antilattice set Ω such that all finite infima and suprema existing in E are preserved.*

Proof. Without loss of generality, by Theorem 2.1, we can assume that $E = \Gamma(G, u)$, where (G, u) is a unital po-group satisfying (RDP_1) . The proof will follow the following steps.

Step 1. Let P be a prime ideal of E . According to Theorem 4.3, $\delta_0(P)$ is a prime subgroup of G , and consider the mapping $\varphi_P: E \rightarrow A(\Omega_P)$, where $\Omega_P = G/\delta_0(P)$, defined by $(x/\delta_0(P))\varphi_P(a) := (x+a)/\delta_0(P)$, $a \in E$ ($x \in G$). Then, for $a, b \in E$, (i) $a \leq b$, implies $\varphi_P(a) \leq \varphi_P(b)$, (ii) $\varphi_P(a+b) = \varphi_P(a) \circ \varphi_P(b)$, (iii) $\varphi_P(a \wedge b) = \varphi_P(a) \wedge \varphi_P(b)$ if $a \wedge b$ is defined in E , (iv) $\varphi_P(a \vee b) = \varphi_P(a) \vee \varphi_P(b)$ if $a \vee b$ is defined in E , and (v) $\{a \in E: \varphi_P(a) = 0\} = \bigcap \{-x + \delta_0(P) + x: x \in G\} \cap E = P$. Moreover, we have $E(P) := \varphi_P(E) \subseteq \Gamma(A(\Omega_P), \varphi_P(u))$.

Step 2. Let $g \in G$ and let $g \not\leq 0$ and set $U(g) := \{h \in G: h \geq g\}$, where $E = \Gamma(G, u)$. We denote by $A(g)$ an ideal of E which is maximal with respect to the property $U(g) \cap A(g) = \emptyset$. Since $0 \notin U(g)$, $A(g)$ exists due to the Zorn lemma. We assert $A(g)$ is a prime ideal of E . Let $I \cap J = A(g)$, where I and J are ideals of E . Assume (ad absurdum hypothesis) $A(g)$ that is a proper subset of I as well as of J . Take $a \in I \cap U(g)$ and $b \in J \cap U(g)$. We have $0, g \leq a, b$. By (RIP) holding in (G, u) , there is an element $c \in G$ such that $0, g \leq c \leq a, b$. Since $0 \leq c \leq a$, we have $c \in E$, and $g \leq c \in I \cap J = A(g)$ which gives $c \in U(g) \cap A(g)$, a contradiction.

Step 3. We define the Cartesian product $E_0 = \prod \{A(\Omega_g): g \in G, g \not\leq 0\}$ of the system of \wedge -antilattices $\{A(\Omega_g)\}_g$, where $\Omega_g = G/\delta_0(A(g))$, and we order E_0 by coordinates. Define a mapping $f: E \rightarrow E_0$ by $f(a) = \{\varphi_g(a)\}_g$ ($a \in E$), where $\varphi_g := \varphi_{A(g)}$, and let us set $C_g = \delta_0(A(g))$.

We claim that f is injective. Assume $f(a) = f(b)$. Then $(x+a)/C_g = (x+b)/C_g$ for all $x \in G$ and $g \not\leq 0$. In particular, for $x = 0$ this gives $a/C_g = b/C_g$. Hence, $a - b = c_g$ for some $c_g \in A(g)$ ($a - b$ is taken in the group G), consequently, $a - b \in \bigcap_{g \not\leq 0} C_g = \{0\}$. This proves that f is an injective homomorphism of E onto $f(E) \subseteq E_0$.

Assume $f(a) \leq f(b)$. If $g = -b + a \not\leq 0$, then $(x+a)/C_g \leq (x+b)/C_g$ for all $x \in G$ and $g \not\leq 0$. Consequently, this holds also for $x = 0$, i.e., $a/C_g \leq b/C_g$ which means $a \leq b + c'_g$ for some $c'_g \in A(g)$. Therefore, $-b + a \leq c'_g$, and $c'_g \in A(g) \cap U(g)$,

a contradiction according to Step 2. The set $f(E)$ can be converted into a pseudo-effect algebra, i.e., $(f(E); \circ, f(0), f(1))$ is a pseudo-effect algebra isomorphic to E , where \circ is the composition of automorphisms defined by coordinates.

According to Step 1, f preserves all finite infima and suprema existing in E .

Step 4. Totally order the nonnegative elements of G , say $\{g_t : t \in T\}$, where T is a linearly ordered set. Set $\Omega_t := G/C_{g_t}$, and without loss of generality we can assume $\Omega_s \cap \Omega_t = \emptyset$ for all $s, t \in T$ such that $s \neq t$. Let $\Omega = \bigcup_{t \in T} \Omega_t$, and define a partial order \preceq on Ω by $\omega_1 \preceq \omega_2$ iff $\omega_1 \in \Omega_s$ and $\omega_2 \in \Omega_t$ and $s < t$ or $s = t$ and $\omega_1 \leq \omega_2$ in Ω_s . Then Ω is a \wedge -antilattice with respect to \preceq .

Define a mapping $f_0: E \rightarrow A(\Omega)$ via: let $\omega \in \Omega$, and $\omega \in \Omega_t$ for a unique $t \in T$. Let $(\omega)f_0(a) = (\omega)(\varphi_{g_t})(a) \in \Omega_t$, where φ_{g_t} is defined in Step 1 and Step 3. Hence, if $a \in E$, then $f_0(a) \mid \Omega_t$ maps Ω_t onto Ω_t for all $t \in T$. Similarly as in Step 3, f_0 is injective from E onto $f_0(E)$, and $f_0(E)$ is a pseudo-effect algebra of automorphisms of Ω (indeed, f_0 practically coincides with the function f defined in Step 3), which finishes the proof. \square

As a direct consequence of Theorem 5.1, we show that every pseudo MV -algebra is isomorphic to a pseudo MV -algebra of automorphisms of a linearly ordered set Ω .

Corollary 5.2. *Every pseudo MV -algebra M can be represented as a pseudo MV -algebra of automorphisms from $A(\Omega)$ for some linearly ordered set Ω .*

Proof. Since a pseudo MV -algebra is a distributive lattice, an ideal of a pseudo MV -algebra M (considered as a pseudo-effect algebra) is prime iff M/P is a linearly ordered set. Consequently, $M = \Gamma(G, u)$ for some unital ℓ -group (G, u) , M satisfies (RDP_2) , hence also (RDP_1) . Hence, the set Ω from the proof of Theorem 5.1 is linearly ordered, which by Theorem 5.1 gives the assertion in question. \square

References

- [1] A. Dvurečenskij: Pseudo MV -algebras are intervals in ℓ -groups. J. Austral. Math. Soc. 72 (2002), 427–445. [Zbl 1027.06014](#)
- [2] A. Dvurečenskij: Ideals of pseudo-effect algebras and their applications. Tatra Mt. Math. Publ. 27 (2003), 45–65. [Zbl pre02172903](#)
- [3] A. Dvurečenskij, S. Pulmannová: New Trends in Quantum Structures. Kluwer Acad. Publ., Dordrecht, Ister Science, Bratislava, 2000. [Zbl 0987.81005](#)
- [4] A. Dvurečenskij, T. Vetterlein: Pseudoeffect algebras. I. Basic properties. Inter. J. Theor. Phys. 40 (2001), 685–701. [Zbl 0994.81008](#)
- [5] A. Dvurečenskij, T. Vetterlein: Pseudoeffect algebras. II. Group representations. Inter. J. Theor. Phys. 40 (2001), 703–726. [Zbl 0994.81009](#)
- [6] G. Georgescu, A. Iorgulescu: Pseudo- MV algebras. Multi. Val. Logic 6 (2001), 95–135. [Zbl 1014.06008](#)

- [7] *A. M. W. Glass*: Polars and their applications in directed interpolation groups. *Trans. Amer. Math. Soc.* *166* (1972), 1–25. [Zbl 0235.06004](#)
- [8] *P. Hájek*: Observations on non-commutative fuzzy logic. *Soft Computing* *8* (2003), 38–43. [Zbl pre02184852](#)
- [9] *C. Holland*: The lattice-ordered group of automorphism of an ordered set. *Michigan Math. J.* *10* (1963), 399–408. [Zbl 0116.02102](#)

Author's address: Mathematical Institute, Slovak Academy of Sciences, Štefánikova 49, SK-814 73 Bratislava, Slovakia, e-mail: dvurecen@mat.savba.sk.