

Bojan Kuzma

Noninvertibility preservers on Banach algebras

Czechoslovak Mathematical Journal, Vol. 56 (2006), No. 3, 919–921

Persistent URL: <http://dml.cz/dmlcz/128116>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NONINVERTIBILITY PRESERVERS ON BANACH ALGEBRAS

BOJAN KUZMA, Ljubljana

(Received April 26, 2004)

Abstract. It is proved that a linear surjection $\Phi: \mathcal{A} \rightarrow \mathcal{B}$, which preserves noninvertibility between semisimple, unital, complex Banach algebras, is automatically injective.

Keywords: linear preserver, noninvertible element, semisimple Banach algebra, socle

MSC 2000: 46H05, 46H10, 47B48

In a recent result, Brešar, Fošner, and Šemrl [3] extended Sourour's result [4] on the form of linear bijection, which preserve invertibility, from $\mathcal{B}(X)$ to arbitrary semisimple Banach algebras with 'large socle' (see also Aupetit and Mouton [2]). The present note was motivated by Sourour's question in [4]: Is a linear, unital surjection $\Phi: \mathcal{B}(X) \rightarrow \mathcal{B}(Y)$, which preserves invertibility, necessarily injective? We show below, with help of [3], that the answer is affirmative when 'invertibility' is replaced by 'noninvertibility'.

Before stating the result, we collect some terminology: If a is an element of a Banach algebra \mathcal{A} , we let $\text{Sp}(a)$ be its *spectrum* and $\text{soc } \mathcal{A}$ the *socle* of \mathcal{A} (see [1]). Recall that an ideal I of \mathcal{A} is called *essential* if it has a nonzero intersection with every nonzero ideal of \mathcal{A} ; in semisimple Banach algebras this is equivalent to $a \cdot I = 0 \Rightarrow a = 0$ for each $a \in \mathcal{A}$. As an example, if $\mathcal{A} = \mathcal{B}(X)$ then $\text{soc } \mathcal{A}$ equals the ideal of finite-rank operators, and is *essential*. Finally, a linear mapping Φ *preserves noninvertibility (in one direction)* if $\Phi(a)$ is not invertible whenever a is not invertible.

We will prove the following

Theorem 1. *Let $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ be a linear surjection that preserves noninvertibility between semisimple, unital, complex Banach algebras \mathcal{A} and \mathcal{B} (in one direction*

This work was supported by grants from the Ministry of Science of Slovenia.

only). Then it is bounded and bijective. Moreover, there exists an invertible $a \in \mathcal{A}$ such that $\Psi(x) := \Phi(ax)$ satisfies

$$(1) \quad \Psi(\Psi^{-1}(y^2) - \Psi^{-1}(y)^2) \cdot \text{soc } \mathcal{B} = 0 \quad \forall y \in \mathcal{B}.$$

In particular, if $\text{soc } \mathcal{B}$ is an essential ideal of \mathcal{B} then Ψ^{-1} , hence also Ψ , is a Jordan isomorphism. In this case, $\Phi(x) = \Psi(a^{-1}x) = b \cdot \Psi'(x)$, where $\Psi, \Psi': \mathcal{A} \rightarrow \mathcal{B}$ are Jordan isomorphisms, and $b \in \mathcal{B}$ is invertible.

P r o o f. We claim that Φ is injective:

Indeed, suppose $\Phi(n) = 0$ for some nonzero $n \in \mathcal{A}$. As \mathcal{A} is semisimple, we can then find some $c \in \mathcal{A}$ such that cn is not a quasinilpotent. By surjectivity, $\Phi(a) = \mathbf{1}$ for some a , which is necessarily invertible. Now, as $\text{Sp}[(1 - \xi)c + \xi a^{-1}n] \neq \{0\}$ at $\xi = 0$, the Scarcity Lemma (see [1, Theorem 3.4.25, and Corollary 3.4.18]) on the analytic multifunction $\xi \mapsto \text{Sp}[(1 - \xi)c + \xi a^{-1}n]$ implies that it differs from $\{0\}$ for any ξ off some subset $\Omega \subset \mathbb{C}$ with zero capacity. Observe that such Ω can contain no interval [1, Corollary A.1.27], and that a belongs to the open set of invertibles, while the mapping $x \mapsto x^{-1}$ is continuous at $x = a^{-1}$. Consequently, we may replace, if necessary, c by some $(1 - \xi)c + \xi a^{-1}$ to ensure that, in addition to $\text{Sp}(cn) \neq \{0\}$, the element c is invertible, and that the line-interval $[c^{-1}, a]$ contains solely invertible elements.

Let $b := c^{-1} - a$, and let $\mathcal{D} := \{\mu \in \mathbb{C}; (a + \mu b) \text{ is invertible}\}$ be an open subset, which contains $[0, 1]$. If $\mu \in \mathcal{D}$ is sufficiently small, the right-hand side of

$$\Phi(a + \mu b + \lambda n) = \Phi(a) + \mu\Phi(b) + 0 = \mathbf{1} + \mu\Phi(b),$$

is invertible for any λ . However,

$$(2) \quad a + \mu b + \lambda n = (a + \mu b) \cdot (\mathbf{1} + \lambda(a + \mu b)^{-1}n)$$

and the analytic function $\mu \mapsto (a + \mu b)^{-1}n$ has at least one nonzero spectral point at $\mu := 1$. By the Scarcity Lemma we may find arbitrarily small $\mu \in [0, 1]$, such that $(a + \mu b)^{-1}n$ is not a quasinilpotent. Consequently, for any of these small μ , the right-hand side of (2) is noninvertible for at least some λ , contradicting the fact that it is mapped into invertible $\Phi(a + \mu b + \lambda n) = \mathbf{1} + \mu\Phi(b)$. Thus, Φ is injective, hence also bijective.

Since a is invertible, the mapping $\Psi(x) := \Phi(ax)$ is also bijective. Its inverse is unital and preserves invertibility between semisimple Banach algebras. Obviously then, $\text{Sp}(\Psi^{-1}(y)) \subseteq \text{Sp}(y)$, so Ψ^{-1} is bounded by [1, Theorem 5.5.2]. The same holds for $\Phi: x \mapsto \Psi(a^{-1}x)$ by the Open Mapping Theorem. Eq. (1) now follows from [3, Main Theorem], which proves the first part.

Finally, if $\text{soc } \mathcal{B}$ is essential then, plainly, Ψ^{-1} and Ψ are Jordan. By [4, Proposition 1.3] such Ψ preserves invertibility in both directions. Hence, $b := \Phi(\mathbf{1}) = \Psi(a^{-1})$ is invertible, and the mapping $\Psi'(x) := b^{-1}\Phi(x)$ is a unital bijection, whose inverse preserves invertibility. As before we derive (1) for Ψ' in place of Ψ , and then conclude that Ψ' is Jordan. \square

References

- [1] *B. Aupetit*: A Primer on Spectral Theory. Springer-Verlag, New York, 1991. [Zbl 0715.46023](#)
- [2] *B. Aupetit, H. du T. Mouton*: Spectrum-preserving linear mappings in Banach algebras. *Stud. Math.* 109 (1994), 91–100. [Zbl 0829.46039](#)
- [3] *M. Brešar, A. Fošner, and P. Šemrl*: A note on invertibility preservers on Banach algebras. *Proc. Amer. Math. Soc.* 131 (2003), 3833–3837. [Zbl 1035.46036](#)
- [4] *A. R. Sourour*: Invertibility preserving maps on $\mathcal{L}(X)$. *Trans. Amer. Math. Soc.* 348 (1996), 13–30. [Zbl 0843.47023](#)

Author's address: Bojan Kuzma, Institute of Mathematics, Physics, and Mechanics, Department of Mathematics, Jadranska 19, 1000 Ljubljana, Slovenia, e-mail: bojan.kuzma@pef.upr.si.