

Long Miao

On complemented subgroups of finite groups

Czechoslovak Mathematical Journal, Vol. 56 (2006), No. 3, 1019–1028

Persistent URL: <http://dml.cz/dmlcz/128127>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON COMPLEMENTED SUBGROUPS OF FINITE GROUPS

LONG MIAO, Yangzhou

(Received August 14, 2004)

Abstract. A subgroup H of a group G is said to be complemented in G if there exists a subgroup K of G such that $G = HK$ and $H \cap K = 1$. In this paper we determine the structure of finite groups with some complemented primary subgroups, and obtain some new results about p -nilpotent groups.

Keywords: finite group, p -nilpotent group, primary subgroups, complemented subgroups

MSC 2000: 20D10, 20D20

1. INTRODUCTION

A subgroup H of a group G is said to be supplemented in G if there exists a subgroup K of G such that $G = HK$. Furthermore, a subgroup H of G is said to be complemented in G if there exists a subgroup K of G such that $G = HK$ and $H \cap K = 1$. It is obvious that the existence of supplements for some families of subgroups of a group gives a lot of information about its structure. For instance, Kegel [8], [9] showed that a group G is soluble if every maximal subgroup of G either has a cyclic supplement in G or if some nilpotent subgroup of G has a nilpotent supplement in G . Hall [6] proved that a group G is soluble if and only if every Sylow subgroup of G is complemented in G . Arad and Ward [1] proved that a group G is soluble if and only if every Sylow 2-subgroup and every Sylow 3-subgroup of G are complemented in G . More recently, A. Ballester-Bolinches and Guo Xiuyun [2] proved that the class of all finite supersoluble groups with elementary abelian Sylow subgroups is just the class of all finite groups for which every minimal subgroup is complemented.

This work is supported by NNSF of China (Grant No. 10471118).

The aim of this paper is to take these studies further. In fact, we analyze the finite group for which some primary subgroups are complemented. We obtain a series of new results for the p -nilpotency of finite groups.

All groups considered in this paper will be finite. Our notation is standard and can be found in [4] and [10]. We denote the fact that G is the semi-product of subgroups H and K by $G = [H]K$, where H is normal in G .

A subgroup H is called the second maximal subgroup of G , if H is the maximal subgroup of some maximal subgroup of G . A subgroup H is called the third maximal subgroup of G , if H is the maximal subgroup of some second maximal subgroup of G .

Let π be a set of primes. We say that $G \in E_\pi$ if G has a Hall π -subgroup. We say that $G \in C_\pi$ if any two Hall π -subgroups of G are conjugate in G . We say that $G \in D_\pi$ if $G \in C_\pi$ and every π -subgroup of G is contained in a Hall π -subgroup of G .

Let \mathcal{F} be a class of groups. \mathcal{F} is called Q -closed if G/N is in \mathcal{F} for any normal subgroup N of G when $G \in \mathcal{F}$. \mathcal{F} is called S -closed if any subgroup K of G is in \mathcal{F} when $G \in \mathcal{F}$. We call \mathcal{F} a formation provided that (1) if $G \in \mathcal{F}$ and $H \trianglelefteq G$, then $G/H \in \mathcal{F}$, and (2) if G/M and G/N are in \mathcal{F} , then $G/M \cap N$ is in \mathcal{F} for normal subgroups M and N of G . Each group has a smallest normal subgroup N such that $G/N \in \mathcal{F}$. This uniquely determined normal subgroup of G is called the \mathcal{F} -residual subgroup of G and denoted by $G^{\mathcal{F}}$. A formation \mathcal{F} is said to be saturated if $G/\Phi(G) \in \mathcal{F}$ implies that $G \in \mathcal{F}$. (cf. [4, Chapt. 2 and 3]). As we all know, the class of all p -nilpotent groups is a saturated formation.

2. PRELIMINARIES

For the sake of easy reference, we first give some basic definitions and known results from the literature.

Lemma 2.1 ([2, Lemma 1]). *Let G be a group and N a normal subgroup of G . Then the following statements hold.*

- (1) *If $H \leq K \leq G$ and H is complemented in G , then H is complemented in K .*
- (2) *If N is contained in H and H is complemented in G , then H/N is complemented in G/N .*
- (3) *Let π be a set of primes. If N is a π' -subgroup and A is a π -subgroup of G , then A is complemented in G if and only if AN/N is complemented in G/N .*

Lemma 2.2 (cf. [4, Theorem 1.8.17]). *Let N be a normal subgroup of a group G ($N \neq 1$). If $N \cap \Phi(G) = 1$, then the Fitting subgroup $F(N)$ of N is the direct product of minimal normal subgroups of G which are contained in $F(N)$.*

Lemma 2.3 ([3, Main Theorem]). *Suppose that a finite group G has a Hall π -subgroup where π is a set of primes not containing 2. Then all Hall π -subgroups of G are conjugate.*

Lemma 2.4. *Let G be a finite group and p be a prime divisor of $|G|$ such that $(|G|, p^2 - 1) = 1$. Assume that the order of G is not divisible by p^3 . Then G is p -nilpotent.*

Proof. Assume that the claim is false and choose G to be a counterexample of minimal order. Since every proper subgroup and every proper quotient group also satisfy the hypothesis of the lemma, the minimal choice of G implies that G is a minimal non- p -nilpotent group but every proper subgroup and every proper quotient group of are p -nilpotent. Therefore $G = [P]Q$ with Q cyclic [10, 9.1.9]. Since both $\Phi(P)$ and $\Phi(G)$ are in $Z(G) = 1$, we see that P is an elementary abelian Sylow p -subgroup and Q a cyclic group of order q . $Q \cong G/P$ and $N_G(P)/C_G(P)$ is isomorphic to a subgroup of $\text{Aut}(P)$. Hence q divides $p(p+1)(p-1)$ [10, 3.2.7]. Since $p \neq q$ and $(|G|, p^2 - 1) = 1$, G is p -nilpotent by the Burnside p -nilpotent Theorem, a contradiction.

The final contradiction completes our proof. □

Lemma 2.5 ([7, IV, 5.4, P₄₃₄]). *Suppose that G is a group which is not p -nilpotent but whose proper subgroups are all p -nilpotent. Then G is a group which is not nilpotent but whose proper subgroups are all nilpotent.*

Lemma 2.6 ([7, 5.2, P₂₈₁]). *Suppose that G is a group which is not nilpotent but whose proper subgroups are all nilpotent. Then*

- 1) G has a normal Sylow p -subgroup P for some prime p and $G/P \cong Q$, where Q is a non-normal cyclic q -subgroup for some prime $q \neq p$.
- 2) $P/\Phi(P)$ is a minimal normal subgroup of $G/\Phi(P)$.
- 3) If P is non-abelian and $p \neq 2$, then the exponent of P is p .
- 4) If P is non-abelian and $p = 2$, then the exponent of P is 4.
- 5) If P is abelian, then P is of exponent p .

Lemma 2.7 ([12, P₆₇]). *Let K be a subgroup of G . If $|G : K| = p$, where p is the smallest prime divisor of $|G|$, then $K \trianglelefteq G$.*

Lemma 2.8. *Let G be a finite group and p be the prime divisor of $|G|$ such that $(|G|, p^2 - 1) = 1$. If G/L is p -nilpotent and $p^3 \nmid |L|$, then G is p -nilpotent.*

Proof. By hypothesis and Lemma 2.4, we know that L is p -nilpotent and L has a normal p -complement $L_{p'}$. Since $L_{p'}$ char L and L is normal in G , we have $L_{p'} \trianglelefteq G$. Therefore $G/L \cong (G/L_{p'})/(L/L_{p'})$ is p -nilpotent. There exists a Hall p' -subgroup $(H/L_{p'})/(L/L_{p'})$ of $(G/L_{p'})/(L/L_{p'})$ and $H/L_{p'} \trianglelefteq G/L_{p'}$. By the Schur-Zassenhaus Theorem, we have $H/L_{p'} = [L/L_{p'}]H_1/L_{p'}$, where $H_1/L_{p'}$ is a Hall p' -subgroup of $H/L_{p'}$. Then by Lemma 2.4 we have $H_1/L_{p'} \trianglelefteq H/L_{p'}$ and so $H_1/L_{p'} \text{ char } H/L_{p'} \trianglelefteq G/L_{p'}$. Therefore $H_1/L_{p'} \trianglelefteq G/L_{p'}$ and hence $G/L_{p'}$ is p -nilpotent. Thus G is p -nilpotent. \square

Lemma 2.9 ([13]). *Let P be an elementary abelian p -group of order p^n , where p is a prime. Then $|\text{Aut}(P)| = k_n \cdot p^{n(n-1)/2}$, here $k_n = \prod_{i=1}^n (p^i - 1)$.*

Lemma 2.10 ([13]). *Let G be a group of order p^n , where p is a prime. Then $|\text{Aut}(G)|$ is the factor of the order of $\text{Aut}(P)$, where P is an elementary abelian p -group of order p^n .*

Lemma 2.11 ([7]). *Let G be a finite group and U any p -subgroup of G . If $N_G(U)/C_G(U)$ is a p -subgroup, then G is p -nilpotent.*

Lemma 2.12 ([5, Lemma 5]). *Let \mathcal{F} be an S -closed local formation and H a subgroup of G . Then $H \cap Z_{\mathcal{F}}(G) \subseteq Z_{\mathcal{F}}(H)$.*

3. MAIN RESULTS

Theorem 3.1. *Let G be a finite group and p a prime divisor of $|G|$ such that $(|G|, p^2 - 1) = 1$. If there exists a normal subgroup N in G such that G/N is p -nilpotent and each subgroup of N of order p^2 is complemented in G , then G is p -nilpotent.*

Proof. Assume that the claim is false and choose G to be a counterexample of minimal order.

By Lemma 2.8 and the hypothesis, we have $|N|_p > p^2$. Let L be a proper subgroup of G . We prove that conditions of the Theorem are inherited by L . Clearly, $L/L \cap N \cong LN/N \leq G/N$ implies that $L/L \cap N$ is p -nilpotent. If $|L \cap N|_p \leq p^2$, then L is p -nilpotent by Lemma 2.8. If $|L \cap N|_p > p^2$, then each subgroup of $L \cap N$ of order p^2 is complemented in G by Lemma 2.1 and hence is complemented in L , thus L is p -nilpotent by induction. Thus G is a minimal non- p -nilpotent group. Now Lemma 2.5 implies that G is a group which is not nilpotent but whose proper subgroups are all nilpotent. Then by Lemma 2.6, we have $G = PQ$, where P is

normal in G and Q is a non-normal cyclic Sylow q -subgroup of G . It is clear that $P/\Phi(P)$ is a minimal normal subgroup of $G/\Phi(P)$.

Clearly, $P \leq N$. Let $A \leq N$ and $|A| = p^2$. By hypothesis, there exists a subgroup K of G such that $G = AK$ and $K \cap A = 1$. By hypothesis, K is nilpotent and $K = K_p \times K_{p'}$. If $K_p = 1$, then $P = A$ and hence G is p -nilpotent by Lemma 2.4, a contradiction. If $K_p \neq 1$, then K_p is the second maximal subgroup of P . Then we consider the subgroup $N_G(K_p)$. Since $K \leq N_G(K_p)$, we have $|G : N_G(K_p)| = p$ or $N_G(K_p) = G$. If $|G : N_G(K_p)| = p$, then we have $N_G(K_p) < G$. Let $P_1 = P \cap N_G(K_p)$. Then $P_1 \trianglelefteq G$. If $P_1 \leq \Phi(P)$, then $P = P \cap AN_G(K_p) = A(P \cap N_G(K_p)) = A$, a contradiction. If $P_1 \not\leq \Phi(P)$, then $P_1\Phi(P)/\Phi(P) = P/\Phi(P)$ by Lemma 2.6. In this case, $P = P_1$ and hence $N_G(K_p) = G$, a contradiction. If $|G : N_G(K_p)| = 1$, then $K_p \trianglelefteq G$. Since G/K_p is p -nilpotent by Lemma 2.4, we have $P = K_p$, a contradiction.

The final contradiction completes our proof. □

Corollary 3.2. *Let G be a finite group and p a prime divisor of $|G|$ such that $(|G|, p^2 - 1) = 1$. If each subgroup of G of order p^2 is complemented in G , then G is p -nilpotent.*

Theorem 3.3. *Suppose that G is a group with a normal subgroup H such that G/H is p -nilpotent for some prime divisor p of $|G|$. If every cyclic subgroup of order 4 of H is complemented in G and every subgroup of H of order p is contained in $Z_{\mathcal{F}}(G)$, where \mathcal{F} is the class of all p -nilpotent groups, then G is p -nilpotent.*

Proof. Assume that the claim is false and choose G to be a counterexample of the smallest order.

The hypothesis is inherited by all proper subgroups of G . Let K be a proper subgroup of G . Then $K/K \cap H \cong KH/H \leq G/H$ implies that $K/K \cap H$ is p -nilpotent. Every cyclic subgroup of $K \cap H$ of order 4 is complemented in G and hence is complemented in K by Lemma 2.1. Every subgroup of $H \cap K$ of order p is contained in $K \cap Z_{\mathcal{F}}(G) \leq Z_{\mathcal{F}}(K)$ by Lemma 2.12. Thus G is a minimal non- p -nilpotent group. Now Lemma 2.5 implies that G is a group which is not nilpotent but whose proper subgroups are all nilpotent. Thus by Lemma 2.6, G has a normal Sylow p -subgroup P and $G/P \cong Q$, where Q is a non-normal cyclic Sylow q -subgroup of G , and $P/\Phi(P)$ is a minimal normal subgroup of $G/\Phi(P)$. We consider the following cases.

Case 1: P is abelian. By Lemma 2.6, P is an elementary abelian p -group. Since G/H is p -nilpotent, we have $P \leq H$. By hypothesis, every subgroup of H of order p is contained in $Z_{\mathcal{F}}(G)$, thus $P \leq Z_{\mathcal{F}}(G)$ and hence G is p -nilpotent, a contradiction.

Case 2: P is non-abelian and $p > 2$. By Lemma 2.6, the exponent of P is p and every subgroup of H of order p is contained in $Z_{\mathcal{F}}(G)$. Therefore $P \leq Z_{\mathcal{F}}(G)$ and we see that G is p -nilpotent, a contradiction.

Case 3: P is non-abelian and $p = 2$. Let A be a cyclic subgroup of H of order 4. By hypothesis, A is complemented in G and there exists a subgroup L of G such that $G = AL$ and $L \cap A = 1$. Since L is nilpotent, we have $L = L_p \times L_{p'}$. If $L_p = 1$, then $P = A$, a contradiction. Clearly, L_p is the 2-maximal subgroup of P . We consider $N_G(L_p)$. Since $L \leq N_G(L_p)$, we have $|G : N_G(L_p)| = 2$ or $|G : N_G(L_p)| = 1$. If $|G : N_G(L_p)| = 2$, then $N_G(L_p) \trianglelefteq G$ by Lemma 2.7 and hence G is 2-nilpotent, a contradiction. If $|G : N_G(L_p)| = 1$, then $L_p \trianglelefteq G$. Since $P/\Phi(P)$ is the minimal normal subgroup of $G/\Phi(P)$, we have $P = L_p$ or $L_p \leq \Phi(P)$. It is clear that $P = L_p$ is impossible. If $L_p \leq \Phi(P)$, then $P = AL_p = A$, a contradiction.

The final contradiction completes our proof. □

Corollary 3.4. *Let G be a group and p be the prime divisor of $|G|$. If every cyclic subgroup of order 4 is complemented in G and every subgroup of G of order p is contained in $Z_{\mathcal{F}}(G)$, where \mathcal{F} is the class of all p -nilpotent groups, then G is p -nilpotent.*

Theorem 3.5. *Let G be a group and $(|G|, 21) = 1$. If each subgroup of G of order 8 (if it exists) is complemented in G , then G is 2-nilpotent.*

Proof. Assume that the theorem is false and choose G to be a counterexample of the smallest order. Let 2^α be the order of a Sylow 2-subgroup P of G .

If $2 \nmid |G|$, then G is 2-nilpotent. If P is cyclic, then G is 2-nilpotent by [10, 10.1.9]. So we can suppose that P is not cyclic. Let L be a proper subgroup of G . We prove that L inherits the condition of the theorem. If $8 \nmid |L|$, then L is 2-nilpotent by Lemma 2.4. If $8 \mid |L|$, then each subgroup of L of order 8 is complemented in G and hence is complemented in L by Lemma 2.1, so L is 2-nilpotent by induction. Thus we may assume that G is a minimal non-2-nilpotent group. Now Lemma 2.5 implies that G is a group which is not nilpotent but whose proper subgroups are all nilpotent. Thus by Lemma 2.6, we have $G = PQ$, where P is normal in G and Q is a non-normal cyclic Sylow q -subgroup of G ($q \neq p$), and $P/\Phi(P)$ is a minimal normal subgroup of $G/\Phi(P)$.

Let H be a subgroup of G of order 8. By hypothesis, there exists a subgroup K of G such that $G = HK$ and $K \cap H = 1$.

First, we claim that $H \neq P$. Otherwise $P = H \trianglelefteq G$. If $\Phi(P) \neq 1$, then $G/\Phi(P)$ is 2-nilpotent by Lemma 2.4 and hence G is 2-nilpotent, a contradiction. It follows from $\Phi(P) = 1$ and Lemma 2.6 that P is an elementary abelian 2-group. Next we will consider $N_G(P)/C_G(P)$. It is clear that $N_G(P)/C_G(P)$ is isomorphic to a

subgroup of $\text{Aut}(P)$. By Lemma 2.9 we have $|\text{Aut}(G)| = 168$. Since $(|G|, 21) = 1$, we see that $N_G(P) = C_G(P)$. Then, by the Burnside Theorem (cf. [10]), we see that G is 2-nilpotent, a contradiction.

Since $H \cap K = 1$ and K is nilpotent, K_2 is the third maximal subgroup of P . We claim that $K_2 \not\trianglelefteq G$. Otherwise if $K_2 \trianglelefteq G$, then $P = K_2$ or $K_2 \leq \Phi(P)$. It is clear that $P = K_2$ is impossible. If $K_2 \leq \Phi(P)$, then $P = HK_2 = H$, a contradiction. Consider the subgroup $N_G(K_2)$. If $|G : N_G(K_2)| = 2$, then $N_G(K_2)$ is a nilpotent normal subgroup of G by Lemma 2.7 and so G is 2-nilpotent, a contradiction. If $|G : N_G(K_2)| = 4$, then we consider $N_G((N_G(K_2))_2)$. It is clear that $(N_G(K_2))_2$ is the second maximal subgroup of P . If $(N_G(K_2))_2 \trianglelefteq G$, then $G/(N_G(K_2))_2$ is 2-nilpotent by Lemma 2.4 and we have $P \leq (N_G(K_2))_2$, a contradiction. Thereby we may assume that $|G : N_G((N_G(K_2))_2)| = 2$. By Lemma 2.7 we know that $N_G((N_G(K_2))_2) \trianglelefteq G$. Since $N_G((N_G(K_2))_2)$ is a nilpotent normal subgroup of G , we know that G is 2-nilpotent, a contradiction.

The final contradiction completes the proof. □

Lemma 3.6. *Let G be a finite group with $(|G|, 21) = 1$. Assume that every third maximal subgroup (if it exists) of a Sylow 2-subgroup of G is complemented in G . Then $G/O_2(G)$ is 2-nilpotent.*

Proof. Assume that the claim is false and choose G to be a counterexample of minimal order. Let P be a Sylow 2-subgroup of G . Furthermore, we have:

(1) $O_2(G) = 1$.

If $O_2(G) = P$, then $G/O_2(G)$ is a 2'-group and of course it is 2-nilpotent, a contradiction. If $O_2(G) = P_1$, where P_1 is the maximal subgroup of P , then $G/O_2(G)$ is 2-nilpotent since $|G/O_2(G)|_2 = 2$, a contradiction. If $O_2(G) = P_2$ where P_2 is the second maximal subgroup of P , then $2^3 \nmid |G/O_2(G)|$. Hence $G/O_2(G)$ is 2-nilpotent by Lemma 2.4, a contradiction. If $1 < O_2(G) < P_2$, then $G/O_2(G)$ satisfies the hypothesis and the minimal choice of G implies that $G/O_2(G) \cong G/O_2(G)/O_2(G/O_2(G))$ is 2-nilpotent, a contradiction.

(2) $|G|$ is divisible by 2^4 .

If $2^3 \nmid |G|$ and $(|G|, 21) = 1$, then G is 2-nilpotent by Lemma 2.4, a contradiction. If $2^3 \mid |G|$ and $2^4 \nmid |G|$, then $|G_2| = 2^3$. Next we consider $N_G(U)/C_G(U)$, where U is any 2-subgroup of G . If $U = P$, then $N_G(P)/C_G(P)$ is isomorphic to a subgroup of $\text{Aut}(P)$. By Lemma 2.9 and Lemma 2.10, $N_G(P)/C_G(P)$ is a 2-subgroup. If $U \neq P$, it is easy to see that $N_G(U)/C_G(U)$ is also a 2-group according to Lemma 2.9 and Lemma 2.10. Then by Lemma 2.11 it is clear that G is 2-nilpotent in this case, a contradiction.

(3) *For every third maximal subgroup P_3 of a Sylow 2-subgroup P of G , the complement of P_3 in G is 2-nilpotent.*

By the hypothesis, P_3 is complemented in G . There exists a subgroup K_3 of G such that $G = P_3K_3$ and $K_3 \cap P_3 = 1$. By (2), we know that K_3 is 2-nilpotent.

(4) G is 2-nilpotent.

Let $N = N_G((K_3)_{2'})$ and $K_3 = (K_3)_2(K_3)_{2'}$. By (3), $K_3 \leq N$. So we have $G = P_3K_3 = P_3N$. If $N = G$, then G is 2-nilpotent, a contradiction. Let $P_3 \leq P_2 \leq P_1 \leq P$ where P_2 is the second maximal subgroup of P and P_1 is the maximal subgroup of P . Hence $G = P_3K_3 = P_2K_3 = P_2N$. If $P_2 \leq N$, then G is 2-nilpotent, a contradiction. So we may assume $P_2 \cap N < P_2$. We may choose a maximal subgroup P_3^* of P_2 such that $P_2 \cap N \leq P_3^*$. It is clear that P_3^* is the third maximal subgroup of P . By (3), P_3^* is complemented in G and the complement K_3^* of P_3^* is 2-nilpotent. We denote $K_3^* = (K_3^*)_2(K_3^*)_{2'}$. Lemma 2.3 implies that $G \in C_{2'}$. Now both $(K_3)_{2'}$ and $(K_3^*)_{2'}$ are the Hall $2'$ -subgroups of G , these two subgroups are conjugate in G . Let $(K_3)_{2'} = ((K_3^*)_{2'})^g$. Since $G = P_3^*K_3^*$ and $K_3^* \leq N_G((K_3^*)_{2'})$, we may choose $g \in P_3^*$. We also note that $(K_3^*)^g$ normalizes $((K_3^*)_{2'})^g = (K_3)_{2'}$ and hence $(K_3^*)^g \leq N$. Now $G = G^g = (P_3^*K_3^*)^g = P_3^*N$. Therefore $P_2 = P_2 \cap P_3^*N = P_3^*(P_2 \cap N) = P_3^*$, contrary to the choice of G .

The final contradiction completes our proof. \square

Theorem 3.7. *Let G be a finite group with $(|G|, 21) = 1$. Assume that there exists a normal subgroup N of G such that G/N is 2-nilpotent and every third maximal subgroup of a Sylow subgroup of N is complemented in G . Then G is 2-nilpotent.*

Proof. Assume that the claim is false and choose G to be a counterexample of minimal order. Then

(1) G is soluble, G has a minimal normal subgroup $L \leq N$ and L is an elementary abelian r -group for some prime r .

By hypothesis, every third maximal subgroup of every Sylow subgroup of N is complemented in G , thus is complemented in N by Lemma 2.1. By the choice of G and Lemma 3.6, N is soluble and hence G is soluble. Let L be a minimal normal subgroup of G which is contained in N . Then L is an elementary abelian r -group for some prime r .

(2) G/L is 2-nilpotent and L is the unique minimal normal subgroup of G which is contained in N . Furthermore, $L = F(N) = C_N(L)$.

In fact, $(G/L)/(N/L) \cong G/N$ is 2-nilpotent and $(|G/L|, 21) = 1$. Let R_1/L be a third maximal subgroup of the Sylow r -subgroup of N/L . Then R_1 is a third maximal subgroup of the Sylow r -subgroup R of N . By hypothesis of the theorem R_1 is complemented in G . By Lemma 2.1, R_1/L is complemented in G/L . Set Q_1/L be a third maximal subgroup of the Sylow q -subgroup of N/L , where $q \neq r$. It is clear that $Q_1 = Q_1^*L$, where Q_1^* is a third maximal subgroup of the Sylow q -subgroup

of N . By the hypothesis Q_1^* is complemented in G . Hence Q_1^*L/L is complemented in G/L by Lemma 2.1. We have proved that G/L satisfies the hypothesis of the theorem. Hence G/L is 2-nilpotent by the minimal choice of G . Since the class of all 2-nilpotent groups is a saturated formation, we can easily prove that L is the unique minimal normal subgroup of G which is contained in N , $L \not\leq \Phi(G)$. By Lemma 2.2, $F(N) = L$. The solubility of N implies that $L \leq C_N(F(N)) \leq F(N)$ and so $C_N(L) = F(N) = L$.

(3) L is a Sylow 2-subgroup of N .

By (1), we know that G is soluble. If $2 \nmid |N|$, then G is 2-nilpotent since G/N is 2-nilpotent, a contradiction. If $2 \neq r$, then G is 2-nilpotent by (2), a contradiction. Therefore L is an elementary abelian 2-subgroup of G which is contained in N . Let D be a Hall $2'$ -subgroup of N ; then it is clear that LD/L is a Hall $2'$ -subgroup of N/L . We have $LD/L \trianglelefteq N/L$ since N/L is 2-nilpotent. So $LD \trianglelefteq N$. Let P be a Sylow 2-subgroup of N . Then $L < P$ and $PD = PLD$ is a subgroup of N . Note that every third maximal subgroup of a Sylow subgroup of PD is complemented in G and hence is complemented in PD by Lemma 2.1. Therefore PD satisfies the hypothesis for G . If $PD < G$, the minimal choice of G implies that PD is 2-nilpotent, in particular, $D \trianglelefteq PD$. Hence $LD = L \times D$ and $D \leq C_N(L) = L$, a contradiction.

Now we assume that $G = PD = N$ and $L < P$. Since N/L is 2-nilpotent, $LD \trianglelefteq N = G$. By the Frattini argument, $G = LN_G(D)$. Note that $L \cap N_G(D) = 1$ since L is the unique minimal normal subgroup of G which is contained in N and D is not normal in G in this case. Therefore $G = [L]N_G(D)$. Let P_2 be a Sylow 2-subgroup of $N_G(D)$. Then LP_2 is a Sylow 2-subgroup of G . Choose a third maximal subgroup P_3 of LP_2 such that $P_2 \leq P_3$. Otherwise, if P_2 is the maximal subgroup of LP_2 , then $|L| = 2$ and hence G is 2-nilpotent by Lemma 2.8, a contradiction. If P_2 is the second maximal subgroup of LP_2 , then $|L| = 2^2$ and hence G is 2-nilpotent by Lemma 2.8 and (2), a contradiction. Clearly, $L \not\leq P_3$ and hence $(P_3)_G = 1$. By our hypothesis, P_3 is complemented in G . There exists a subgroup K of G such that $G = P_3K$ and $K \cap P_3 \leq (P_3)_G = 1$. It follows that K has a normal 2-complement which is in fact a Hall $2'$ -subgroup D_1 of G in this case. By the hypothesis and Lemma 2.3, there exists an element $g \in L$ such that $D_1^g = D$. Since $P_2 \leq P_3 < P_2^* < P_1 < LP_2$, where P_1 is a maximal subgroup of LP_2 which contains P_2^* , and P_2^* is a second maximal subgroup of LP_2 which contains P_3 , we have $G = P_3K = P_1K = (P_1K)^g = P_1K^g$. Since $K^g \cong K$ has a normal 2-complement D and $D = D_1^g \leq K^g$, it follows that $K^g \leq N_G(D)$. Since $LP_2 = LP_2 \cap G = LP_2 \cap P_1K^g = P_1(LP_2 \cap K^g)$, we have that $LP_2 \cap K^g \not\leq P_2$. Otherwise $LP_2 \leq P_1P_2 = P_1$, a contradiction. Therefore P_2 is a proper subgroup of $P_4 = \langle P_2, LP_2 \cap K^g \rangle$ where P_4 is a subgroup of the Sylow 2-subgroup LP_2 . Now both P_2 and K^g are contained in $N_G(D)$ and we see that P_4 is a 2-subgroup of $N_G(D)$

which contains a Sylow subgroup P_2 as a proper subgroup, a contradiction. Hence L is a Sylow 2-subgroup of N .

(4) G is 2-nilpotent.

If $|L| \leq 4$, then G is 2-nilpotent by Lemma 2.8 and (2), a contradiction. If $|L| = 2^3$, then G is 2-nilpotent by Lemma 3.6. So we may assume $|L| > 8$. Let L_1 be a nontrivial third maximal subgroup of L . Then L_1 is complemented in G . There exists a subgroup K of G such that $L_1K = G$ and $K \cap L_1 = 1$. Therefore $L = L_1(L \cap K)$ and $L \cap K \trianglelefteq G$. We have $L \cap K = L$, otherwise, $L \cap K = 1$ and $L = L_1$, a contradiction. Therefore $K = G$, a contradiction.

The final contradiction completes our proof. \square

Corollary 3.8. *Let G a finite group with $(|G|, 21) = 1$. If every third maximal subgroup of every Sylow subgroup of G is complemented in G , then G is 2-nilpotent.*

References

- [1] *Z. Arad, M.B. Ward*: New criteria for the solvability of finite groups. *J. Algebra* **77** (1982), 234–246.
- [2] *A. Ballester-Bolínches, X. Guo*: On complemented subgroups of finite groups. *Arch. Math.* **72** (1999), 161–166.
- [3] *F. Gross*: Conjugacy of odd order Hall subgroup. *Bull. London Math. Soc.* **19** (1987), 311–319.
- [4] *W. Guo*: *The Theory of Classes of Groups*. Kluwer Academic Publishers, Beijing-New York-Dordrecht-Boston-London, 2000.
- [5] *W. Guo*: The influence of minimal subgroups on the structure of finite groups. *Southeast Asian Bulletin of Mathematics* **22** (1998), 287–290.
- [6] *P. Hall*: A characteristic property of soluble groups. *J. London Math. Soc.* **12** (1937), 188–200.
- [7] *B. Huppert*: *Endliche Gruppen I*. Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [8] *O.H. Kegel*: On Huppert’s characterization of finite supersoluble groups. In: *Proc. Internat. Conf. Theory Groups*, Canberra, 1965. New York, 1967, pp. 209–215.
- [9] *O.H. Kegel*: Produkte nilpotenter gruppen. *Arch. Math.* **12** (1961), 90–93.
- [10] *D.J. Robinson*: *A Course in the Theory of Groups*. Springer-Verlag, Berlin-New York, 1993.
- [11] *Y. Wang*: Finite groups with some subgroups of Sylow subgroups c -supplemented. *J. Algebra* **224** (2000), 467–478.
- [12] *M. Xu*: *An Introduction to Finite Groups*. Science Press, Beijing, 1999. (In Chinese.)
- [13] *Y. Zhang*: *The Structure of Finite Groups*. Science Press, Beijing, 1982. (In Chinese.)

Author’s address: Department of Mathematics, Yangzhou University, Yangzhou 225002, P.R. China, e-mail: miaolong714@sohu.com.