

David Cruz-Uribe; Alberto Fiorenza

Weighted endpoint estimates for commutators of fractional integrals

*Czechoslovak Mathematical Journal*, Vol. 57 (2007), No. 1, 153–160

Persistent URL: <http://dml.cz/dmlcz/128162>

## Terms of use:

© Institute of Mathematics AS CR, 2007

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

WEIGHTED ENDPOINT ESTIMATES FOR COMMUTATORS  
OF FRACTIONAL INTEGRALS

D. CRUZ-URIBE, SFO, Hartford, A. FIORENZA, Napoli

(Received November 9, 2004)

*Abstract.* Given  $\alpha$ ,  $0 < \alpha < n$ , and  $b \in \text{BMO}$ , we give sufficient conditions on weights for the commutator of the fractional integral operator,  $[b, I_\alpha]$ , to satisfy weighted endpoint inequalities on  $\mathbb{R}^n$  and on bounded domains. These results extend our earlier work [3], where we considered unweighted inequalities on  $\mathbb{R}^n$ .

*Keywords:* fractional integrals, commutators, BMO, weights, Orlicz spaces, maximal functions

*MSC 2000:* 42B20, 42B25

1. INTRODUCTION

Given  $\alpha$ ,  $0 < \alpha < n$ , define the fractional integral operator  $I_\alpha$  by

$$I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-\alpha}} dy;$$

for  $b \in \text{BMO}$ , define the commutator  $[b, I_\alpha]$  by

$$[b, I_\alpha]f(x) = b(x)I_\alpha f(x) - I_\alpha(bf)(x) = \int_{\mathbb{R}^n} (b(x) - b(y)) \frac{f(y)}{|x - y|^{n-\alpha}} dy.$$

Commutators were first introduced by Chanillo [1] who proved  $L^p$  estimates,  $1 < p < \infty$ . In [3] we proved the following endpoint estimate:

$$|\{x \in \mathbb{R}^n : |[b, I_\alpha]f(x)| > t\}| \leq C\Psi\left(\int_{\mathbb{R}^n} B\left(\|b\|_{\text{BMO}} \frac{|f(x)|}{t}\right) dx\right),$$

where  $B(t) = t \log(e + t)$  and  $\Psi(t) = [t \log(e + t^{\alpha/n})]^{n/(n-\alpha)}$ .

We initially conjectured that the corresponding weighted inequality was

$$w(\{x \in \mathbb{R}^n : |[b, I_\alpha]f(x)| > t\}) \leq C \Psi \left( \int_{\mathbb{R}^n} B \left( \|b\|_{\text{BMO}} \frac{|f(x)|}{t} \right) w(x)^{1/q} dx \right),$$

where  $w \in A_1$  and  $q = n/(n - \alpha)$ . However, this conjecture proved to be false, and we gave a counterexample [3, Example 1.8].

At the time, we were not able to make a new conjecture. One difficulty we had was that locally, our original conjecture appeared to be true; the counterexample works because it exploits the decay of the weight at infinity. Careful consideration of this behavior yielded two results. First, we found the correct weighted endpoint inequality for the commutator on  $\mathbb{R}^n$ . Second, we showed that if we restrict ourselves to a bounded domain, then there is a sharper endpoint inequality which is even simpler than our original conjecture.

**Theorem 1.1.** *Given  $\alpha$ ,  $0 < \alpha < n$ , and a function  $b \in \text{BMO}$ , let  $B(t) = t \log(e + t)$ ,  $\Psi(t) = [t \log(e + t^{\alpha/n})]^{n/(n-\alpha)}$ ,  $\Theta(t) = t^{1-\alpha/n} \log(e + t^{-\alpha/n})$ , and  $q = n/(n - \alpha)$ . Then for each weight  $w \in A_1$ , there exists a constant  $C$  such that*

$$(1.1) \quad w(\{x \in \mathbb{R}^n : |[b, I_\alpha]f(x)| > t\}) \leq C \Psi \left( \int_{\mathbb{R}^n} B \left( \|b\|_{\text{BMO}} \frac{|f(x)|}{t} \right) \Theta(w(x)) dx \right).$$

But, given any bounded domain  $\Omega \subset \mathbb{R}^n$ ,

$$(1.2) \quad w(\{x \in \Omega : |[b, I_\alpha]f(x)| > t\}) \leq C \left( \int_{\Omega} B \left( \|b\|_{\text{BMO}} \frac{|f(x)|}{t} \right) w(x)^{1/q} dx \right)^q.$$

**Remark 1.2.** If we replace  $f$  by  $f\chi_\Omega$ ,  $\Omega$  unbounded, (1.1) yields a nominally more general result for unbounded domains. However, in both (1.1) and (1.2), the  $A_1$  weights are still defined on all of  $\mathbb{R}^n$ .

Theorem 1.1 is best understood by comparing inequalities (1.1) and (1.2) to the weighted endpoint inequality for the fractional integral operators due to Muckenhoupt and Wheeden [7]. They showed that if  $w \in A_1$ , then

$$(1.3) \quad w(\{x \in \mathbb{R}^n : |I_\alpha f(x)| > t\}) \leq C \left( \frac{1}{t} \int_{\mathbb{R}^n} |f(x)| w(x)^{1/q} dx \right)^q.$$

Intuitively, Theorem 1.1 shows that the commutator  $[b, I_\alpha]$  is more singular than  $I_\alpha$  itself, and that its singularity is worse at infinity.

Theorem 1.1 should also be compared to the analogous result for commutators of singular integral operators (formally corresponding to the case  $\alpha = 0$ ) due to Pérez [9]. If  $T$  is a singular integral operator,  $b \in \text{BMO}$  and  $w \in A_1$ , then

$$w(\{x \in \mathbb{R}^n : |[b, T]f(x)| > t\}) \leq C \int_{\mathbb{R}^n} B\left(\|b\|_{\text{BMO}} \frac{|f(x)|}{t}\right) w(x) dx.$$

The proof of Theorem 1.1 follows the same outline as the proof of the unweighted result on  $\mathbb{R}^n$  [3, Theorem 1.1]. The first step is to prove that the associated Orlicz fractional maximal operator,

$$M_{\alpha, B}f(x) = \sup_{Q \ni x} |Q|^{\alpha/n} \|f\|_{B, Q},$$

satisfies inequalities analogous to (1.1) and (1.2).

**Theorem 1.3.** *With the same notation and hypotheses as in Theorem 1.1, we have that*

$$(1.4) \quad w(\{x \in \mathbb{R}^n : M_{\alpha, B}f(x) > t\}) \leq C \Psi\left(\int_{\mathbb{R}^n} B\left(\frac{|f(x)|}{t}\right) \Theta(w(x)) dx\right)$$

and

$$(1.5) \quad w(\{x \in \Omega : M_{\alpha, B}f(x) > t\}) \leq C \left(\int_{\Omega} B\left(\frac{|f(x)|}{t}\right) w(x)^{1/q} dx\right)^q.$$

We prove Theorem 1.3 in Section 2 below; since we can do so with essentially no more work, we prove a generalization that holds for a large class of Young functions.

The proof of Theorem 1.1 now proceeds as in the unweighted case. Here we sketch the main steps, and we refer the reader to [3] for details. On the left-hand side of (1.1) (or (1.2)) we replace  $|[b, I_\alpha]f(x)|$  by  $M^d([b, I_\alpha]f)(x)$ , where  $M^d$  is the dyadic maximal operator. We then use the good- $\lambda$  inequality relating  $M^d$  and the sharp maximal operator  $M^\#$  (Lemma 6.1 in [3], which remains true in the weighted case since  $w \in A_1$ ) to replace this with  $M^\#([b, I_\alpha]f)(x)$ . Next, we apply the inequality

$$M^\#([b, I_\alpha]f)(x) \leq C \|b\|_{\text{BMO}} [I_\alpha f(x) + M_{\alpha, B}f(x)]$$

(Theorem 1.3 in [3]), which then reduces the estimate to endpoint inequalities for  $I_\alpha$  and  $M_{\alpha, B}$ . To complete the proof we apply inequality (1.3) and Theorem 1.3, and use the fact that  $w^{1/q} \leq \Theta(w)$ .

## 2. ENDPOINT RESULTS FOR ORLICZ FRACTIONAL MAXIMAL OPERATORS

In this section we state and prove two endpoint inequalities for the Orlicz fractional maximal operator. Theorem 1.3 will be an immediate consequence of these results. Hereafter we will assume that the reader is familiar with the basic facts about Orlicz spaces, maximal operators and Muckenhoupt  $A_p$  weights, and we refer the reader to [4], [5], [6], [10] for further information. Also, we will draw heavily on our work [3], and we urge the reader to consult that paper.

**Remark 2.1.** The conclusions of our main results in this section, Theorems 2.3 and 2.5, remain true if  $\alpha = 0$ . (See Pérez [8].) However, to avoid technical difficulties, we restrict ourselves to  $\alpha > 0$ .

To state our results, we need one definition.

**Definition 2.2.** Given an increasing function  $\varphi$ , define a function  $h_\varphi$  by

$$h_\varphi(s) = \sup_{t>0} \frac{\varphi(st)}{\varphi(t)}, \quad 0 \leq s < \infty.$$

The function  $h_\varphi$  could be infinite if  $s > 1$ , but if  $\varphi$  is doubling, then it is finite for all  $0 < s < \infty$ . (See Maligranda [6, Theorem 11.7].) If  $\varphi$  is submultiplicative, then  $h_\varphi \approx \varphi$ . Also, for all  $s, t > 0$ ,  $\varphi(st) \leq h_\varphi(s)\varphi(t)$ .

**Theorem 2.3.** Given  $\alpha$ ,  $0 < \alpha < n$ , let  $B$  be a Young function such that  $B(t)/t^{n/\alpha}$  is decreasing for all  $t > 0$ , and let  $w \in A_1$ . Then there exists a constant  $C$  depending only on  $B$  and the  $A_1$  constant of  $w$  such that for all  $t > 0$ ,  $M_{\alpha,B}$  satisfies the modular weak-type inequality

$$(2.1) \quad \Phi(w(\{x \in \mathbb{R}^n : M_{\alpha,B}f(x) > t\})) \leq C \int_{\mathbb{R}^n} B\left(\frac{f(x)}{t}\right) h_\Phi(w(x)) \, dx$$

for all non-negative  $f \in L_B(\mathbb{R}^n)$ , where

$$(2.2) \quad \Phi(s) = \begin{cases} 0 & \text{if } s = 0, \\ \frac{s}{h_B(s^{\alpha/n})} & \text{if } s > 0. \end{cases}$$

Inequality (1.4) follows easily from Theorem 2.3. Since  $B$  is submultiplicative,  $h_B \approx B$ , so

$$\Phi(t) \approx \frac{t^{1-\alpha/n}}{\log(e + t^{\alpha/n})}.$$

The function  $\Phi$  is invertible with

$$\Phi^{-1}(t) \approx \Psi(t) = [t \log(e + t^{\alpha/n})]^{n/(n-\alpha)}.$$

Thus, inequality (2.1) yields (1.4), provided that

$$h_\Phi(t) \leq C\Theta(t) = Ct^{1-\alpha/n} \log(e + t^{-\alpha/n}).$$

However, this follows from the definition: since  $h_B$  is submultiplicative and  $h_B \approx B$ ,

$$h_\Phi(s) = \sup_{t>0} \frac{\Phi(st)}{\Phi(t)} = s \sup_{t>0} \frac{h_B(t^{\alpha/n})}{h_B((st)^{\alpha/n})} \leq sh_B(s^{-\alpha/n}) \leq CsB(s^{-\alpha/n}) = C\Theta(s).$$

**Remark 2.4.** Note that if we let  $t = s^{-1}$  and use the fact that  $h_B(t) \geq cB(t)$ , we get  $h_\Phi(s) \geq c\Theta(s)$ . Hence,  $\Theta$  is the best possible function we can get by this means.

**Theorem 2.5.** *Given a bounded domain  $\Omega \subset \mathbb{R}^n$  and given  $\alpha$ ,  $0 < \alpha < n$ , let  $B$  be a Young function such that  $B(t)/t^{n/\alpha}$  is decreasing for all  $t > 0$ . Suppose further that there exists  $r$ ,  $1 \leq r \leq n/\alpha$ , such that  $h_B(t) \leq Ct^r$  for all  $t \leq 6 \operatorname{diam}(\Omega)^\alpha$ . Then for each  $w \in A_1$  there exists a constant  $C$ , depending on  $B$ ,  $\operatorname{diam}(\Omega)$  and the  $A_1$  constant of  $w$ , such that for all non-negative  $f \in L_B(\Omega)$ ,*

$$(2.3) \quad w(\{x \in \Omega : M_{\alpha,B}f(x) > t\})^{1-r\alpha/n} \leq C \int_{\Omega} B\left(\frac{f(x)}{t}\right) w(x)^{1-r\alpha/n} dx.$$

Inequality (1.5) is an immediate consequence of Theorem 2.5. Since  $B(t) = t \log(e + t)$ ,  $h_B \approx B$ , so if  $t \leq 6 \operatorname{diam}(\Omega)$ , then  $h_B(t) \leq C \log(e + 6 \operatorname{diam}(\Omega))t$ . Thus inequality (2.3) holds with  $r = 1$ , and this yields (1.5).

**P r o o f** of Theorem 2.3. Our proof is very similar to the proof of Theorem 3.3 in [3], and we refer the reader there for many lemmas and technical details.

Fix a non-negative function  $f$  and  $t > 0$ . Define  $E_t = \{x \in \mathbb{R}^n : M_{\alpha,B}f(x) > t\}$ .

For each  $x \in E_t$  there exists a cube  $Q_x \ni x$  such that

$$|Q_x|^{\alpha/n} \|f\|_{B,Q_x} > t.$$

The collection  $\{Q_x\}_{x \in E_t}$  covers  $E_t$ . Thus, by Lemma 3.14 in [3], there exists  $\beta > 0$  and a collection of disjoint dyadic cubes  $\{P_j\}$  such that  $E_t \subset \bigcup_j 3P_j$  and

$$|P_j|^{\alpha/n} \|f\|_{B,P_j} > \beta t.$$

By the properties of the Luxemburg norm on Orlicz spaces and by Definition 2.2,

$$\begin{aligned}
 (2.4) \quad 1 &< \frac{1}{|P_j|} \int_{P_j} B\left(\frac{|P_j|^{\alpha/n} f(x)}{\beta t}\right) dx \\
 &\leq \frac{Ch_B(|3P_j|^{\alpha/n})}{|3P_j|} \int_{P_j} B\left(\frac{f(x)}{t}\right) dx \\
 &= \frac{C}{\Phi(|3P_j|)} \int_{P_j} B\left(\frac{f(x)}{t}\right) dx.
 \end{aligned}$$

The growth conditions assumed on  $B$  imply (see Lemma 3.12 in [3]) that

$$\Phi(w(E_t)) \leq \Phi\left(\sum_j w(3P_j)\right) \leq \sum_j \Phi(w(3P_j)).$$

Hence, if we combine the two inequalities above and apply Definition 2.2, we get

$$\begin{aligned}
 \Phi(w(E_t)) &\leq C \sum_j \frac{\Phi(w(3P_j))}{\Phi(|3P_j|)} \int_{P_j} B\left(\frac{f(x)}{t}\right) dx \\
 &\leq C \sum_j h_\Phi\left(\frac{w(3P_j)}{|3P_j|}\right) \int_{P_j} B\left(\frac{f(x)}{t}\right) dx;
 \end{aligned}$$

since  $w \in A_1$  and the  $P_j$ 's are disjoint,

$$\begin{aligned}
 \Phi(w(E_t)) &\leq C \sum_j \int_{P_j} B\left(\frac{f(x)}{t}\right) h_\Phi(w(x)) dx \\
 &\leq C \int_{\mathbb{R}^n} B\left(\frac{f(x)}{t}\right) h_\Phi(w(x)) dx.
 \end{aligned}$$

This completes the proof. □

The proof of Theorem 2.5 requires two lemmas.

**Lemma 2.6.** *Given  $\alpha$ ,  $0 < \alpha < n$ , let  $B$  be a Young function such that  $B(t)/t^{n/\alpha}$  is decreasing for all  $t > 0$ . Then  $h_B(s) \leq s^{n/\alpha}$  for all  $s \geq 1$ .*

*Proof.* Fix  $s \geq 1$ . Then for all  $t > 0$ ,

$$\frac{B(st)}{(st)^{n/\alpha}} \leq \frac{B(t)}{t^{n/\alpha}}, \quad \text{or equivalently,} \quad \frac{B(st)}{B(t)} \leq s^{n/\alpha}.$$

Taking the supremum over all  $t$  we get the desired inequality. □

**Lemma 2.7.** *Given  $\alpha$ ,  $0 < \alpha < n$ , let  $B$  be a Young function such that  $B(t)/t^{n/\alpha}$  is decreasing for all  $t > 0$ . If  $Q$  and  $\bar{Q}$  are cubes and  $f$  is a function such that  $\text{supp}(f) \subset Q \subset \bar{Q}$ , then*

$$|\bar{Q}|^{\alpha/n} \|f\|_{B, \bar{Q}} \leq |Q|^{\alpha/n} \|f\|_{B, Q}.$$

*Proof.* Let  $s = |\bar{Q}|/|Q| \geq 1$ . Then by the definition of the Luxemburg norm and by Lemma 2.6,

$$\begin{aligned} |\bar{Q}|^{\alpha/n} \|f\|_{B, \bar{Q}} &= |Q|^{\alpha/n} \|s^{\alpha/n} f\|_{B, \bar{Q}} \\ &= |Q|^{\alpha/n} \inf \left\{ \lambda > 0: \frac{1}{|\bar{Q}|} \int_{\bar{Q}} B\left(\frac{s^{\alpha/n} |f(x)|}{\lambda}\right) dx \leq 1 \right\} \\ &\leq |Q|^{\alpha/n} \inf \left\{ \lambda > 0: \frac{1}{|Q|} \int_Q h_B(s^{\alpha/n}) B\left(\frac{|f(x)|}{\lambda}\right) dx \leq 1 \right\} \\ &\leq |Q|^{\alpha/n} \inf \left\{ \lambda > 0: \frac{1}{|Q|} \int_Q B\left(\frac{|f(x)|}{\lambda}\right) dx \leq 1 \right\} \\ &= |Q|^{\alpha/n} \|f\|_{B, Q}. \end{aligned}$$

□

*Proof of Theorem 2.5.* The proof of this result is nearly the same as the proof of Theorem 2.3. The major difference is that we must show that we can restrict the size of the cubes used to compute  $M_{\alpha, B}$ . Fix  $x \in \Omega$ , and let  $Q$  be any cube containing  $x$ . If  $\ell(Q) \geq \text{diam}(\Omega)$ , then we can maximize the quantity  $|Q|^{\alpha/n} \|f\|_{B, Q}$  by taking  $Q$  to be such that  $\Omega \subset Q$ . Further, by Lemma 2.7 we can increase this quantity by choosing  $Q$  such that  $\ell(Q) \leq \text{diam}(\Omega)$ . Consequently, in computing  $M_{\alpha, B}$  we can restrict ourselves to cubes whose sidelength is at most the diameter of  $\Omega$ .

Thus, in the proof of Theorem 2.3, when we cover the set  $E_t$  by cubes  $Q_x$ , we may assume that each of these cubes has sidelength bounded by  $\text{diam}(\Omega)$ . We then use Lemma 3.14 in [3] to replace these cubes by a collection of dyadic cubes  $P_j$ . But from the proof of this lemma (see [2], [8]) we see that the sidelength of each  $P_j$  is bounded by twice the largest sidelength of the  $Q_x$ 's; therefore  $\ell(P_j) \leq 2 \text{diam}(\Omega)$  for all  $j$ .

By assumption,  $h_B(t) \leq Ct^r$  for  $t \leq 6 \text{diam}(\Omega)^\alpha$ . Therefore,  $h_B(|3P_j|^{\alpha/n}) \leq |3P_j|^{r\alpha/n}$  for all  $j$ . Hence, we can modify the argument that yielded (2.4) to get

$$1 < \frac{C}{|3P_j|^{1-r\alpha/n}} \int_{P_j} B\left(\frac{f(x)}{t}\right) dx.$$

If we now let  $\Phi(t) = t^{1-r\alpha/n}$ , then  $h_\Phi(t) = \Phi(t)$  and we can repeat the remainder of the argument in the proof of Theorem 2.3 to get (2.3). □



### References

- [1] *S. Chanillo*: A note on commutators. *Indiana Math. J.* 31 (1982), 7–16. [Zbl 0523.42015](#)
- [2] *D. Cruz-Uribe, SFO*: New proofs of two-weight norm inequalities for the maximal operator. *Georgian Math. J.* 7 (2000), 33–42. [Zbl 0987.42019](#)
- [3] *D. Cruz-Uribe, SFO, A. Fiorenza*: Endpoint estimates and weighted norm inequalities for commutators of fractional integrals. *Publ. Mat.* 47 (2003), 103–131. [Zbl 1035.42015](#)
- [4] *J. Duoandikoetxea*: *Fourier Analysis*. Grad. Studies Math. Vol. 29. Am. Math. Soc., Providence, 2000. [Zbl 0969.42001](#)
- [5] *J. García-Cuerva, J. L. Rubio de Francia*: *Weighted Norm Inequalities and Related Topics*. Math. Studies Vol. 116. North Holland, Amsterdam, 1985. [Zbl 0578.46046](#)
- [6] *L. Maligranda*: Orlicz spaces and interpolation. *Seminars in Mathematics* 5, IMECC. Universidad Estadual de Campinas, Campinas, 1989. [Zbl 0874.46022](#)
- [7] *B. Muckenhoupt, R. Wheeden*: Weighted norm inequalities for fractional integrals. *Trans. Am. Math. Soc.* 192 (1974), 261–274. [Zbl 0289.26010](#)
- [8] *C. Pérez*: On sufficient conditions for the boundedness of the Hardy-Littlewood maximal operator between weighted  $L^p$ -spaces with different weights. *Proc. London Math. Soc.* 71 (1995), 135–157. [Zbl 0829.42019](#)
- [9] *C. Pérez*: Endpoint estimates for commutators of singular integral operators. *J. Funct. Anal.* 128 (1995), 163–185. [Zbl 0831.42010](#)
- [10] *M. M. Rao, Z. D. Ren*: *Theory of Orlicz Spaces*. Marcel Dekker, New York, 1991. [Zbl 0724.46032](#)

*Authors' addresses:* D. Cruz-Uribe, SFO, Dept. of Mathematics, Trinity College, Hartford, CT 06106-3100, USA, e-mail: [david.cruzuribe@mail.trincoll.edu](mailto:david.cruzuribe@mail.trincoll.edu); A. Fiorenza, Dipartimento di Costruzioni e Metodi Matematici in Architettura, Università di Napoli, Via Monteoliveto, 3, I-80134 Napoli, Italy, and Istituto per le Applicazioni del Calcolo “Mauro Picone”—sezione di Napoli, Consiglio Nazionale delle Ricerche, via Pietro Castellino, 111, I-80131 Napoli, Italy, e-mail: [fiorenza@unina.it](mailto:fiorenza@unina.it).