

Qu Han-Zhang

A topological space is strongly paracompact if and only if for any monotone increasing open cover of it there exists a star-finite open refinement

Czechoslovak Mathematical Journal, Vol. 58 (2008), No. 2, 487–491

Persistent URL: <http://dml.cz/dmlcz/128272>

Terms of use:

© Institute of Mathematics AS CR, 2008

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A TOPOLOGICAL SPACE IS STRONGLY PARACOMPACT IF AND ONLY IF FOR ANY MONOTONE INCREASING OPEN COVER OF IT THERE EXISTS A STAR-FINITE OPEN REFINEMENT

QU HAN-ZHANG, Xi'an

(Received April 6, 2006)

Abstract. We get the following result. A topological space is strongly paracompact if and only if for any monotone increasing open cover of it there exists a star-finite open refinement. We positively answer a question of the strongly paracompact property.

Keywords: paracompact property, strongly paracompact property

MSC 2000: 54C35, 54E32

In this paper we assume all spaces are T_2 .

$|u|$ is the cardinal number of a set u .

A set u which consists of some subsets of a topological space X is locally finite if for any $x \in X$ there exists an open subset U of X , such that $x \in U$ and $|\{V; V \cap U \neq \varnothing, V \in u\}|$ is a finite cardinal number.

A set u which consists of subsets of a topological space X is star-finite if for any $U \in u$, $|\{V; V \cap U \neq \varnothing, V \in u\}|$ is a finite cardinal number.

A set u which consists of subsets of a topological space X is star $< k$ (k is a cardinal number) if for any $U \in u$, $|\{V; V \cap U \neq \varnothing, V \in u\}| < k$.

A topological space is paracompact if for any open cover of it there exists a locally finite open refinement. A topological space is strongly paracompact if for any open cover of it there exists a star-finite open refinement.

On the paracompactness of a topological space there is the following theorem.

Theorem 1. *A topological space is paracompact if and only if for any monotone increasing open cover of it there exists a locally finite open refinement.*

Comparing the strong paracompactness with the paracompactness, there is the following question.

Question 1. If for any monotone increasing open cover of a topological space there exists a star-finite open refinement, is it strongly paracompact?¹

In the nineties of the last century, the following question on strong paracompactness was posed.

Question 2. If for any infinite open cover u of a topological space X there exists a star $< |u|$ open refinement, is it strongly paracompact?

Theorem 2. *If X is a topological space, the following conditions are equivalent.*

- (1) *For any infinite open cover u of X there exists a star $< |u|$ open refinement.*
- (2) *For any infinite monotone increasing open cover u of X there exists a star $< |u|$ open refinement.*
- (3) *For any infinite monotone increasing open cover u of X there exists a star-finite open refinement.*

Proof. (1) \implies (2). This is obvious.

(2) \implies (3). Let u be any infinite monotone increasing open cover of X . We make the proof by transfinite induction with respect to $|u| = k$.

(a) If $k = \omega$, a star $< |u|$ open refinement of u is a star-finite open refinement of u .

(b) If $k > \omega$, we assume that for any infinite monotone increasing open cover of u , if $|u| < k$, there exists a star-finite open refinement.

$|u| = k$. If $\text{cof}(k) < k$, then there is a subcover u_1 of u such that $|u_1| = \text{cof}(k)$. According to the assumption of the transfinite induction, there exists a star-finite open refinement of u_1 . It is a star-finite open refinement of u .

If $\text{cof}(k) = k$, there exists a star $< |u|$ open refinement v of u . In elements of v we introduce the equivalence relation as follows.

$V \approx V' \iff$ there exists a finite subset $\{V_i; 1 \leq i \leq n\}$ of v , such that $V \cap V_1 \neq \varnothing$, $V_n \cap V' \neq \varnothing$, $1 \leq i \leq n-1$, $V_i \cap V_{i+1} \neq \varnothing$. Let $v = \bigcup \{v_\alpha; \alpha < k'\}$, with v_α the equivalence classes of v .

For any $\alpha < k'$, $W_\alpha = \bigcup \{V; V \in v_\alpha\}$. W_α is closed and open. Because $W_\alpha = X - \bigcup \{W_\beta; \beta \neq \alpha, \beta < k'\}$. $|v_\alpha| < k$. if $|v_\alpha| = k$, $\text{cof}(k) = \omega < k$. This is a contradiction with $\text{cof}(k) = k$. That is, $k = k'$.

For any $\alpha < k$, $v_\alpha = \{V_{\alpha,\beta}; \beta < k_\alpha\}$, $k_\alpha < k$. $\beta = 0$, $V'^{\alpha,0} = V_{\alpha,0} \cup (X - W_\alpha)$; $0 < \beta < k_\alpha$, $V'^{\alpha,\beta} = (\bigcup \{V_{\alpha,\gamma}; \gamma < \beta\}) \cup (X - W_\alpha)$.

¹ The project was supported by NSFC (Grant No. 90207015) and the Science Research Foundation of the Education of Shaanxi Province of China (grant 06JK192).

$v_{\alpha'} = \{V'_{\alpha,\beta}; \beta < k_{\alpha}\}$ is a monotone increasing open cover of X and $|v_{\alpha}| < k$. According to the assumption of the transfinite induction there exists a star-finite open refinement w'_{α} of v_{α} . $w_{\alpha} = \{W' \cap W_{\alpha}; W' \in w'_{\alpha}\}$ is a star-finite open cover of W_{α} and a refinement of v_{α} . $w = \bigcup\{w_{\alpha}; \alpha < k\}$ is a star-finite open refinement of u .

(3) \implies (1) $u = \{U_{\alpha}; \alpha < k\}$ is an infinite open cover of X such that $|u| = k$. $V_0 = U_0; 0 < \alpha < k, V_{\alpha} = \bigcup\{U_{\beta}; \beta < \alpha\}$. $v = \{V_{\alpha}; \alpha < k\}$ is a monotone increasing infinite open cover of X . Then there exists a star-finite open refinement $w = \{W_{\beta}; \beta < k'\}$ of v such that there exists a function $f: k' \rightarrow k$ such that for any $\beta < k', W_{\beta} \subset V_{f(\beta)} \cdot \{U_{\alpha} \cap W_{\beta}; \alpha < f(\beta), \beta < k'\}$ is a star-finite open refinement of u . For any $U_{\alpha} \cap W_{\beta}$, because w is star-finite, $\{W; W \cap W_{\beta} \neq \varphi, W \in w\} = \{W_{\beta_1}, \dots, W_{\beta_n}\}$. Then $|\bigcup\{\{U_{\alpha} \cap W_{\beta_i}; \alpha < f(\beta_i)\}, i \leq n\}| \leq |f(\beta_1)| + \dots + |f(\beta_n)| = \max\{|f(\beta_i)|; i \leq n\} < k$. That is, it is star-finite.

According to Theorem 1 and Theorem 2 the following result is obtained.

Lemma 1. *If for any infinite open cover u of a topological space there exists a star-finite open refinement, it is paracompact.*

On the star-finite property, we have the following results.

Lemma 2. *Let u, v consist of some subsets of a topological space X . If u, v are star-finite, then*

- (1) $u \wedge v = \{U \cap V; U \in u, V \in v\}$ is star-finite;
- (2) $\{\bigcap \Phi; \Phi \subset u, |\Phi| < \omega\}$ is star-finite;
- (3) $\{\text{star}\{x, u\} = \bigcup\{U; x \in U, U \in u\}; x \in X\}$ is star-finite;
- (4) $\{\text{star}\{U, u\} = \bigcup\{V; V \cap U \neq \varphi, V \in u\}; U \in u\}$ is star-finite.

Theorem 3. *If for any infinite monotone increasing open cover of a topological space there exists a star-finite open refinement, it is strongly paracompact.*

Proof. Let X be a topological space which satisfies the condition of Theorem 3. Let $u = \{U_{\alpha}; \alpha < k\}$ ($|u| = k$) be any infinite open cover of X . We make the proof by transfinite induction with respect to $k = |u|$.

If $k = \omega, \alpha = 0, V_0 = U_0; 0 < \alpha < \omega, V_{\alpha} = \bigcup\{U_{\beta}; \beta < \alpha\}$. $v = \{V_{\alpha}; \alpha < \omega\}$ is a monotone increasing infinite open cover of X . There exists a star-finite open refinement $w = \{W_{\beta}; \beta < k'\}$ of v such that there exists a function $f: k' \rightarrow \omega$ such that for any $\beta < k', W_{\beta} \subset V_{f(\beta)} \cdot \{U_{\alpha} \cap W_{\beta}; \alpha < f(\beta), \beta < k'\}$ is a star-finite open refinement of u . For any $U_{\alpha} \cap W_{\beta}$, because w is star-finite, $\{W; W \cap W_{\beta} \neq \varphi, W \in w\} = \{W_{\beta_1}, \dots, W_{\beta_n}\}$. Thus $|\bigcup\{\{U_{\alpha} \cap W_{\beta_i}; \alpha < f(\beta_i)\}, i \leq n\}| \leq |f(\beta_1)| + \dots + |f(\beta_n)| = \max\{|f(\beta_i)|; i \leq n\} < \omega$. That is, it is star-finite.

If $k > \omega$, we suppose that for any infinite open cover of u , which satisfies the hypothesis of Theorem 3, if $|u| < k$, there exists a star-finite open refinement.

For any infinite open cover u of X , which satisfies the condition of Theorem 3, $|u| = k$, $\alpha = 0$, $V_0 = U_0$; if $0 < \alpha < k$, $V_\alpha = \bigcup\{U_\beta; \beta < \alpha\}$. $v = \{V_\alpha; \alpha < k\}$ is a monotone increasing infinite open cover of X . There exists a star-finite open refinement $w' = \{W'_\beta; \beta < k'\}$ of v such that there exists a function $f: k' \rightarrow k$ such that for any $\beta < k'$, $W'_\beta \subset V_{f(\beta)}$. According to Lemma 1, X is paracompact. Then there is an open cover $w = \{W_\beta; \beta < k'\}$ of w' such that for any $\beta < k'$, $\overline{W_\beta} \subset W'_\beta$.

For any $\beta < k'$, $\{W'_\beta \cap U_\alpha; \alpha < f(\beta)\} \cup \{X - \overline{W_\beta}\}$ is an open cover of X and $|\{W'_\beta \cap U_\alpha; \alpha < f(\beta)\} \cup \{X - \overline{W_\beta}\}| = |f(\beta)| < k$. According to the hypothesis of the transfinite induction, there exists a star-finite open refinement w'_β of it. Set $w_\beta = \{W; W \cap \overline{W_\beta} \neq \varnothing, W \in w'_\beta\}$.

For any $x \in \overline{W_\beta}$, $O_{\beta,x} = \bigcap\{W; x \in W, W \in w_\beta\}$. w_β is star-finite. $|\{W; x \in W, W \in w_\beta\}| < \omega$. $O_{\beta,x}$ is an open subset of X . According to Lemma 2, $\{O_{\beta,x}; x \in \overline{W_\beta}\}$ is star-finite.

$O_x = \bigcap\{O_{\beta,x}; x \in \overline{W_\beta}\}$. w is star-finite. $|\{O_{\beta,x}; x \in \overline{W_\beta}, \beta < k'\}| < \omega$. O_x is an open subset of X .

$G_x = X - \bigcup\{\overline{W_\beta}; x \notin \overline{W_\beta}, \beta < k'\}$. w' is star-finite. $\bigcup\{\overline{W_\beta}; x \notin \overline{W_\beta}, \beta < k'\}$ is closed. That is, G_x is open.

According to Lemma 2 $\{\text{star}\{x, \{\overline{W_\beta}; \beta < k'\}\}; x \in X\}$ is star-finite. For any $x \in X$, $G_x \subset \text{star}\{x, \{\overline{W_\beta}; \beta < k'\}\}$. So $\{G_x; x \in X\}$ is star-finite.

$\{G_x \cap O_x; x \in X\}$ is star-finite.

For any $x \in X$, if $y \notin \bigcup\{\overline{W_\beta}; x \in \overline{W_\beta}, \beta < k'\}$, $G_x \cap G_y = \varnothing$. That is, $(G_x \cap O_x) \cap (G_y \cap O_y) = \varnothing$.

Because $\{\overline{W_\beta}; \beta < k'\}$ is star-finite, $|\{\overline{W_\beta}; x \in \overline{W_\beta}, \beta < k'\}| < \omega$. That is, $\{\overline{W_\beta}; x \in \overline{W_\beta}, \beta < k'\} = \{\overline{W_{\beta_1}}, \dots, \overline{W_{\beta_n}}\}$. For any $1 \leq i \leq n$, $\{O_{\beta_i,y}; y \in \overline{W_{\beta_i}}\}$ is finite. So for any $1 \leq i \leq n$, $\{O_{\beta_i,y}; O_{\beta_i,x} \cap O_{\beta_i,y} \neq \varnothing; y \in \overline{W_{\beta_i}}\}$ is finite. $\{O_y; O_x \cap O_y \neq \varnothing, y \in \overline{W_{\beta_1}} \cup \dots \cup \overline{W_{\beta_n}}\}$ is finite. $\{G_y; G_x \cap G_y \neq \varnothing, y \in \overline{W_{\beta_1}} \cup \dots \cup \overline{W_{\beta_n}}\}$ is finite. $\{O_y \cap G_y; O_x \cap G_x \cap O_y \cap G_y \neq \varnothing, y \in \overline{W_{\beta_1}} \cup \dots \cup \overline{W_{\beta_n}}\}$ is finite.

That is, $\{G_x \cap O_x; x \in X\}$ is star-finite. It is a star-finite open refinement of u .

According to the principle of the transfinite induction, for any infinite open cover of u , there exists a star-finite open refinement.

That is, X is strongly paracompact.

According to Theorem 2, Theorem 3, we can get following.

Theorem 4. *If for any infinite open cover u of a topological space there exists a star $< |u|$ open refinement, it is strongly paracompact.*

Theorem 5. *If for any monotone increasing open cover of a topological space there exists a star-finite open refinement, it is strongly paracompact.*

We thus positively answer a question on the strong paracompactness of a topological space.

References

- [1] *Ryszard Engelking*: General Topology. Panstwowe Wydawnictwo Naukowe, 1977.
- [2] *Yoshikazu Yasui*: Generalization Paracompactness. Chapter 13 of Topics in General topology (Editor by Kitti Morit and Jun-iti Nagata), Elsevier Science Publishers B.V., 1989.
- [3] *Wang Shu-tang, Dai Jin-sheng and Wang Shang-zhi*: Principles of General Topology. Science and Technique Press of Shannxi, China, 1985.
- [4] *Jiang Ji-guang*: Especial Topics of General Topology. Education Press of Sichuan, China, 1991.

Author's address: Qu Han-Zhang, Department of Applied mathematics and Physics, Xi'an Institute of Post and Telecommunications, 710121, Xi'an, China, e-mail: QHZ_002@163.com.